

094242

GEOLOGICAL REPORT

on

the

RUSTY SPRINGS PROPERTY

Connor 1-9, Diduck 1-4, Matt 1-4, Trog 1-4, 18, 20, 29, 50, 55, 57 Claims

Yukon Territory

N.T.S. 116 K/8 and 116 K/9

Latitude 66° 30' N, Longitude 140° 25' W

prepared for

EAGLE PLAINS RESOURCES LTD.  
2720 17<sup>th</sup> St. S  
CRANBROOK, BRITISH COLUMBIA  
VIC 6Y6

by

C.C. Downie, P.Geol.  
Eagle Plains Resources Ltd.  
122 13<sup>th</sup> Ave. S  
Cranbrook, B.C. V1C 2V5

January 30<sup>th</sup>, 2002

YUKON ENERGY, MINES  
& RESOURCES LIBRARY  
P.O. BOX 2703  
WHITEHORSE, YUKON Y1A 2C6



**TABLE OF CONTENTS**

	<b><u>PAGE NO.</u></b>
<b>INTRODUCTION.....</b>	<b>3</b>
<b>LOCATION AND ACCESS .....</b>	<b>5</b>
<b>PROPERTY TENURE .....</b>	<b>6</b>
<b>HISTORY OF EXPLORATION .....</b>	<b>8</b>
<b>REGIONAL GEOLOGICAL SETTING .....</b>	<b>11</b>
<b>PROPERTY GEOLOGY .....</b>	<b>11</b>
<b>2001 PROGRAM AND RESULTS .....</b>	<b>20</b>
<b>CONCLUSIONS AND RECOMMENDATIONS .....</b>	<b>22</b>
<b>REFERENCES.....</b>	<b>24</b>

### LIST OF TABLES

TABLE 1:SUMMARY OF EXPLORATION WORK.....	after page 8
TABLE 2:EXPLORATION METHODS .....	after page 8

### LIST OF FIGURES

FIGURE 1: PROPERTY LOCATION MAP.....	after page 5
FIGURE 2: REGIONAL GEOLOGY;CLAIM LOCATION MAP;2001 SAMPLE LOCATION .....	in pocket
FIGURE 3: GEOLOGY OF THE CORE OF THE RUSTY SPRINGS ANTICLINE, MIKE AND ORMA HILLS; 1999 DRILLHOLE LOCATIONS.....	in pocket
FIGURE 4: CROSS SECTION MIKE AND ORMA HILLS.....	in pocket

### LIST OF APPENDICES

APPENDIX I: CERTIFICATES OF QUALIFICATION
APPENDIX II: STATEMENT OF EXPENDITURES
APPENDIX III: ANALYTICAL RESULTS
APPENDIX IV: 2001 SAMPLE DESCRIPTIONS

## INTRODUCTION

Despite many years of exploration and relatively limited success, the Rusty Springs prospect retains considerable potential for a large-tonnage deposit. The property lies within the east-vergent Taiga-Nahoni foldbelt, occurring in the core of a structural culmination exposing host Lower and Middle Devonian Ogilvie Formation dolostones. Mineralization occurs in stratabound and discordant zones along the contact with the overlying Devono-Mississippian Unnamed shale. Various deposit models, ranging from Mississippi Valley-type to epithermal vein-type have been employed. Poor exposure and relatively deep weathering resulting from the lack of Pleistocene glaciation account for the lack of consensus with regard to genesis, and accumulating evidence points to the potential for a high-temperature, carbonate-hosted massive sulphide deposit (manto-chimney complex). The great extent of mineralized and altered rocks, together with their stratabound nature, common significant thicknesses, local high grades, and potential for supergene enrichment suggest that Rusty Springs remains an attractive drill-oriented exploration target.

The Rusty Springs Property area has seen sporadic exploration since 1975, when rusty ground seeps were recognized during regional oil and gas exploration programs. Subsequent ground examination revealed silver-lead-zinc mineralization nearby. Staking of the area by Rio Alto Exploration followed, with systematic exploration programs carried out over the years by various operators.

High-grade mineralization was discovered in the Orma Hill area in 1978, and the focus of exploration efforts were concentrated in this area. Virtually all drilling was aimed at the Orma Vein since this time. Preliminary work, previous to the Orma discovery however, outlined anomalous soil geochemical values in the Mike Hill area. Limited drilling was carried out to define the nature of this mineralization, but met only limited success.

In 1992, the final core claims comprising the Rusty Springs Property were allowed to lapse. They were subsequently restaked, and optioned to Eagle Plains Resources, who now retain a 100% interest in the property.

Bulldozer trenching of the Mike Hill area in 1994 resulted in the discovery of high grade silver-lead-zinc mineralization within silicified carbonate material. Drilling carried out during 1995 was aimed at evaluating the mineralized zones exposed on the Mike Hill. Trenching and soil geochemical sampling was completed at the Big Onion area to follow-up geochemical work initiated during 1994.

In 1996, a 15 hole diamond drill program defined highly anomalous base metal values over significant widths within an apparently stratabound – stratiform horizon at the Ogilvie - Hart River contact. The 1997 program employed a reverse circulation drill in an attempt to improve penetration problems related to the highly abrasive cap rocks overlying the mineralized horizon. The drilling confirmed the presence of strata bound mineralization over a large area.

The 1998 program consisted of a combined shallow seismic and gravity geophysical survey. The survey defined a coincident positive Bouger gravity anomaly and seismic reflection profile interpreted to be related to a shallow sulphide body at the same stratigraphic horizon as sulphide mineralization defined in 1996 - 97.

Evaluation of the Rusty Springs Property continued in 1999 with a \$273,001.81 diamond drilling and geological mapping program undertaken by the Eagle Plains Resources / CanAustra Resources joint venture. CanAustra has an option to earn a 60% interest in the Rusty Springs property by completing \$2,000,000 in exploration expenditures, and making \$70,000 in cash payments to Eagle Plains by 2003. Diamond drilling was directed toward testing geophysical anomalies defined by the 1998 combined seismic and gravity surveys and geological targets generated by 1999 mapping. A total of 616.9 meters (2024 feet) of diamond drilling was completed in three holes. None of the holes were completed to target depth due to drilling problems. One of the holes, RS9901, intersected significant base metal mineralization.

Charlie Greig, a noted structural geologist, was retained in 1999 to compile a detailed structural map of the Rusty Springs property and to define a regional framework for the Rusty Springs mineralization. His work forms the basis for much of this report and was published in 2000 as part of the Yukon Exploration and Geology, 1999; Department of Indian and Northern Affairs.

2001 work on the Rusty Springs Property consisted of a short reconnaissance geological program. The total cost of the 2001 program was \$16,500.08.

Further work is recommended for the property and surrounding region. A \$374,000.00, 1500m diamond drilling program utilizing a modified diamond drill designed to address the difficult drilling conditions encountered at Rusty Springs is recommended for 2002.

## LOCATION AND ACCESS

The Rusty Springs Ag/Pb/Zn/Cu prospect is situated in the north-western part of the Yukon Territory at approximately 66° 30' North latitude and 140° 25' West longitude on N.T.S. mapsheet 116 K/8 and 116 K/9. The property is 8 km south of the Arctic Circle and 29 km east of the Alaska border, near the headwaters of the Salmon Fork of the Yukon River (see Figure 1; following). Relief in the Rusty Springs area is on the order of 1000 metres, with the highest point in the surrounding mountains at about 1500 metres. Summits and ridges are generally rounded and subdued, and the valleys are broad as the area lies in the part of the Yukon that was not glaciated during Pleistocene time.

Access to the property is via wheel or ski-equipped aircraft or by winter road. An all-weather, 600m (2000') airstrip was completed in 1996. Supply centres are located at Dawson City, Yukon (274km), Circle, Alaska (175km), or Fairbanks, Alaska (365km). Airstrip staging areas to Rusty Springs are available along the Dempster Highway at Eagle Plains (164kms), or from the "150 Mile" airstrip (137km).

Road access has been previously developed for winter haulage from Mile 123 (Ogilvie Crossing) on the Dempster Highway over a distance of 193 km. The Dempster Highway is a government-maintained all-weather road providing access from the south. The winter road access traverses gently sloping terrain without any major topographic obstacles.

BEAUFORT SEA

PROPERTY  
LOCATION

Porcupine R.

Old Crow

Inuvik

Arctic Circle



Chance  
Gas Field

Eagle Plains

Circle

Dempster Hwy.

Prairie Creek  
Deposit N.W.T

YUKON

Fairbanks

Yukon R.

ALASKA

Dawson City


Keno Hill

Mayo

U.S.A.  
CANADA

Whitehorse

Ross River



**EAGLE PLAINS RESOURCES  
RUSTY SPRINGS PROJECT**

*LOCATION MAP*

NTS Reference: 116K/8,116K/9

Rev. Date: Jan./98

TOKLAT RESOURCES INC.

Fig: 1

**PROPERTY TENURE**

The total property area consists of 81 quartz-claims, staked in accordance with existing Yukon Quartz Mining Act regulations.

Pertinent tenure is summarized below. Claim and post locations are shown on Figure 2 in pocket.

<u>Claim Name</u>	<u>Record No.</u>	<u>Units</u>	<u>Location Date</u>	<u>Expiry Date</u>
Eric 1	YB41182	1	July 29, 1992	Dec. 10, 2008
Eric 2	YB41183	1	July 29, 1992	Dec. 10, 2008
Eric 3	YB41184	1	July 29, 1992	Dec. 10, 2008
Eric 4	YB41185	1	July 29, 1992	Dec. 10, 2008
Eric 5	YB41186	1	July 29, 1992	Dec. 10, 2008
Eric 6	YB41187	1	July 29, 1992	Dec. 10, 2008
Eric 7	YB48768	1	June 10, 1994	Dec. 10, 2009
Eric 8	YB48769	1	June 10, 1994	Dec. 10, 2009
Jessica 1	YB41188	1	July 29, 1992	Dec. 10, 2008
Jessica 2	YB41189	1	July 29, 1992	Dec. 10, 2008
Jessica 3	YB41190	1	July 29, 1992	Dec. 10, 2008
Jessica 4	YB41191	1	July 29, 1992	Dec. 10, 2008
Jessica 5	YB41192	1	July 29, 1992	Dec. 10, 2008
Jessica 6	YB41193	1	July 29, 1992	Dec. 10, 2008
Jessica 7	YB48750	1	June 10, 1994	Dec. 10, 2009
Jessica 8	YB48751	1	June 10, 1994	Dec. 10, 2009
Shelly 1	YB48752	1	June 10, 1994	Dec. 10, 2009
Shelly 2	YB48753	1	June 10, 1994	Dec. 10, 2009
Shelly 3	YB48754	1	June 10, 1994	Dec. 10, 2009
Shelly 4	YB48755	1	June 10, 1994	Dec. 10, 2009
Shelly 5	YB48756	1	June 10, 1994	Dec. 10, 2009
Shelly 6	YB48757	1	June 10, 1994	Dec. 10, 2009
Shelly 7	YB48758	1	June 10, 1994	Dec. 10, 2009
Shelly 8	YB48759	1	June 10, 1994	Dec. 10, 2009
Shelly 9	YB48760	1	June 10, 1994	Dec. 10, 2009
Shelly 10	YB48761	1	June 10, 1994	Dec. 10, 2009
Shelly 11	YB48762	1	June 10, 1994	Dec. 10, 2009
Shelly 12	YB48763	1	June 10, 1994	Dec. 10, 2009
Shelly 13	YB48764	1	June 10, 1994	Dec. 10, 2009
Shelly 14	YB48765	1	June 10, 1994	Dec. 10, 2009
Shelly 15	YB48766	1	June 10, 1994	Dec. 10, 2009
Shelly 16	YB48767	1	June 10, 1994	Dec. 10, 2009
Joel 1	YB52722	1	Aug. 27, 1994	Dec. 10, 2005
Joel 2	YB52723	1	Aug. 27, 1994	Dec. 10, 2005
Joel 3	YB52724	1	Aug. 27, 1994	Dec. 10, 2005
Joel 4	YB52725	1	Aug. 27, 1994	Dec. 10, 2005
Joel 5	YB53897	1	July 2, 1995	Dec. 10, 2006
Joel 6	YB53898	1	July 2, 1995	Dec. 10, 2006
Joel 7	YB53899	1	July 2, 1995	Dec. 10, 2006
Joel 8	YB53900	1	July 2, 1995	Dec. 10, 2006
Glen	YB53901	1	July 2, 1995	Dec. 10, 2006
Calli	YB53902	1	July 2, 1995	Dec. 10, 2006
Marlo	YB53903	1	July 2, 1995	Dec. 10, 2006
Katie	YB53904	1	July 2, 1995	Dec. 10, 2006
Alecia	YB53905	1	July 2, 1995	Dec. 10, 2006
Kelsey	YB53906	1	July 2, 1995	Dec. 10, 2006
Lauren	YB53907	1	July 2, 1995	Dec. 10, 2006
Tyler	YB53908	1	July 2, 1995	Dec. 10, 2006

<u>Claim Name</u>	<u>Record No.</u>	<u>Units</u>	<u>Location Date</u>	<u>Expiry Date</u>
Casey	YB53909	1	July 2, 1995	Dec. 10, 2006
Lane	YB53910	1	July 2, 1995	Dec. 10, 2006
Kayla	YB53911	1	June 16, 1995	Dec. 10, 2008
Ben	YB53912	1	June 16, 1995	Dec. 10, 2008
Trevor	YB53913	1	June 16, 1995	Dec. 10, 2007
James	YB53914	1	June 16, 1995	Dec. 10, 2007
Connor 1	YB54257	1	Sept. 7, 1995	Dec. 12, 2003
Connor 2	YB54258	1	Sept. 7, 1995	Dec. 12, 2003
Connor 3	YB54259	1	Sept. 7, 1995	Dec. 12, 2003
Connor 4	YB54260	1	Sept. 7, 1995	Dec. 12, 2003
Connor 5	YB54261	1	Sept. 7, 1995	Dec. 12, 2003
Connor 6	YB54262	1	Sept. 7, 1995	Dec. 12, 2003
Connor 7	YB54263	1	Sept. 7, 1995	Dec. 12, 2003
Connor 8	YB54264	1	Sept. 7, 1995	Dec. 12, 2003
Connor 9	YB54265	1	Sept. 7, 1995	Dec. 12, 2003
Matt 1	YB54266	1	Sept. 7, 1995	Dec. 12, 2003
Matt 2	YB54267	1	Sept. 7, 1995	Dec. 12, 2003
Matt 3	YB54268	1	Sept. 7, 1995	Dec. 12, 2003
Matt 4	YB54269	1	Sept. 7, 1995	Dec. 12, 2003
Diduck 1	YB54270	1	Sept. 7, 1995	Dec. 12, 2003
Diduck 2	YB54271	1	Sept. 7, 1995	Dec. 12, 2003
Diduck 3	YB54272	1	Sept. 7, 1995	Dec. 12, 2003
Diduck 4	YB54273	1	Sept. 7, 1995	Dec. 12, 2003
Trog 1-4	YB88193-196	4	July 29, 1996	Dec. 10, 2005
Trog 18	YB88210	1	July 29, 1996	Dec. 10, 2005
Trog 20	YB88212	1	July 29, 1996	Dec. 10, 2005
Trog 29	YB88221	1	July 29, 1996	Dec. 10, 2005
Trog 50	YB88242	1	July 29, 1996	Dec. 10, 2005
Trog 55	YB88247	1	July 29, 1996	Dec. 10, 2005
Trog 57	YB88249	1	July 29, 1996	Dec. 10, 2005

Total: 81 units

**HISTORY OF EXPLORATION (Table 1, Table 2; following)**

The Rusty Springs property was first staked in 1975, after investigation of deep red-orange springs and seeps in the valley of Carrol Creek led to the discovery of nearby silver, lead, zinc, and copper mineralization; the rusty seeps were first noted during petroleum exploration in the area. Since the discovery, the property has been the focus for nearly \$5,000,000.00 of exploration, including ten separate drill campaigns in two major phases (1975-83 and 1994-96) totaling over 10,000 metres of drilling in 123 holes (Table I).

Exploration has mainly targeted high-grade silver, lead, copper, and zinc mineralization within brecciated and quartz- and carbonate-cemented and veined dolomite, and has been based on several genetic models, developed in part by geology students employed on the property and working on Bachelors theses (e.g., Schoel 1978, Hansen 1979, Bankowski 1980a). At various stages of exploration, models used to help guide exploration include: Mississippi Valley-type (MVT); Irish Plains-type (carbonate-hosted exhalative, Bankowski 1980a); epithermal-type (veins and(or) hydrothermal replacement along a karsted surface, with supergene enrichment); and manto-chimney-type (high-temperature, carbonate-hosted massive sulphides). Direct targeting of drillholes utilized various techniques, including prospecting, geologic mapping, geochemistry and geophysics (see Table II). Many of the drill programs were plagued by drilling problems, such as poor recoveries in the strongly oxidized and leached mineralized intervals, or loss of water pressure in blocky brecciated zones with abundant open space. Drilling was often slow and costly in resistant siliceous 'chert' horizons that cap the mineralized stratigraphy. Trenching also met with varying success, mainly because of the deep permafrost and the deep, soliflucted overburden which predominates in unglaciated parts of the Yukon.

During the fall of 1975, while investigating an oolitic iron formation, a rusty spring-seep was observed by M.N. Chernoff. Upon investigation, the spring was found to be associated with high-grade silver, lead, zinc, and copper mineralization. A total of 92 quartz claims and 15 iron claims were staked during the fall and winter seasons.

During the 1976 summer season, a preliminary investigation of the property was conducted by Rio Alto Exploration Ltd., under the supervision of M.N. Chernoff. Exploration completed included helicopter-supported geological mapping, prospecting, sampling of mineralized float, and limited soil geochemical sampling. This work established the structural setting, confirmed the presence of high-grade silver values, and demonstrated the usefulness of soil geochemistry. The mineral occurrences were considered to be hydrothermal vein systems with supergene enrichment possibilities.

Based on encouraging results from this preliminary reconnaissance, a follow-up field program consisting of geological mapping, soil geochemical sampling, and 975 metres (3200 feet) of diamond drilling was conducted in 1977. Again, the results were considered positive, even though poor drill core recoveries were obtained. Additional ground was staked to give a total of 380 quartz claims and 15 iron claims.

A geological thesis by G. Schoel concluded that the mineralization was probably Mississippi Valley type.

During the winter of 1978, fuel, drill equipment, and supplies were ferried to the property by tractor train. That summer, two picket grids (totalling 67 line km) were established over the claims. Further geological mapping, soil geochemical

**TABLE 1 : Summary of Exploration Work, Rusty Springs Property**

Year	Work done	Company	Interpretations	Drilling	Significant results	Expenditure	Reference
1976	staking, prospecting, mapping, limited soil sampling, hand-pitting	Rio Alto Exploration Ltd.	intrusive-related hydrothermal vein systems with supergene enrichment possibilities		Chip samples of float from several localities with 30-40% Zn, 5-15% Cu, and variable Pb and Ag; grab samples commonly averaged 10-70 opt Ag	\$150,000	Chernoff (1976)
1977	prospecting, mapping, grid soil sampling, diamond drilling, staking, metallurgical sampling	Rio Alto Exploration Ltd.	precious metal enriched Mississippi Valley type (MVT) model adopted	3,200 ft. (975 m) in 8 holes	High Ag and Pb values in one hole (123 ft. averaging 33.27 opt Ag, 4.72% Pb, 2.36% Cu) but with poor recoveries	\$187,000	White (1978); Schoel (1978)
1978	extensive linecutting and soil geochemistry, prospecting, diamond drilling, mapping, construction of winter road and airstrip	Rio Alto Exploration Ltd.	mineralized zones on Orma hill follow low angle "fault"; MVT model still accepted	6035 ft. (1,840 m) in 30 holes	stratigraphic control noted on anomalous soil geochem zones following "chert"-dolomite contacts: Cu-Pb-Ag±Zn on Orma hill; Zn±Cu±Pb±Ag on Mike hill; poor recoveries in drilling	\$555,000	Beck (1978)
1979	Induced Polarization and gravity surveys, linecutting, prospecting, mapping, soil sampling, hand pitting, trenching	Rio Alto Exploration Ltd.	MVT model still accepted		extent of upper Ogilvie Formation (mineralized showings or float found throughout) and contacts with overlying siliciclastic rocks established	\$300,000	Hansen and Bankowski (1979), White (1979)
1980	diamond drilling, cat trenching, detailed mapping	E&B Explorations Inc. and Rio Alto Exploration Ltd. joint venture	mineralization considered to be of hydrothermal origin; Ogilvie-Hart River contact still considered a karsted horizon channeling mineralizing solutions	6,000 ft. (1829 m) in 27 holes	poor recoveries in upper parts of holes; numerous cm- to decimetre thick tetrahedrite-tennantite veins intersected and which commonly yielded high Ag, Pb, and Cu values; mineralization on Orma hill in part appears to be vein-related	\$1,200,000	Bankowski (1980), Liedtke (1980)
1982	soil geochemistry, VLF-EM surveys, mapping, trenching, diamond drilling	Kenton Natural Resources Corporation	epithermal veins	1673 ft. (510 m) in 7 holes	common WNW, NW, and NNW trending EM conductors outlined; Orma hill vein systems defined	\$116,000	Davis and Aussant (1982)
1983	fill-in soil geochemistry and VLF-EM surveys, diamond drilling	Kenton Natural Resources Corporation	epithermal veins	1600 ft. (488 m) in 2 holes	focused on Orma Hill vein systems	\$350,000	Aussant (1983)
1986	diamond drilling	Kenton Natural Resources Inc.		1326 ft. (404m) in 2 holes	tested (unsuccessfully) IP anomalies between Orma and Mike hills	\$96,000	Chamberlain (1986)
1992	restaking						

1994	regional reconnaissance; trenching, airstrip and road construction; clean-up	Eagle Plains Resources Ltd.	epithermal veins, MVT		vein mineralization on 040 trend discovered using soil geochem and trenching on Mike Hill; new showings discovered SW of Mike Hill	\$190,000	Downie (1994)
1995	trenching, diamond drilling, soil geochemistry, staking, airstrip and road construction, GPS survey, claim staking	Eagle Plains Resources Ltd.	manto-chimney type carbonate hosted deposits	5440 ft. (1658 m) in 21 holes	15.1 oz/ton Ag, 3% Cu, and 1.3% Zn over 50 ft. (15.3 metres) on Mike Hill	\$539,000	Termuende (1996)
1996	diamond drilling; airstrip extension, road construction, staking	Eagle Plains Resources Ltd.	carbonate-hosted manto type deposits; stratabound hydrothermal mineralization along Ogilvie-Hart River Formation contact	7610 ft (2320 m) in 15 holes	highly anomalous base metal values over significant widths along Ogilvie-Hart River Formation contact	\$560,000	Termuende and Downie (1997)
1997	reverse-circulation drilling, surface mapping, prospecting, road and drill pad construction, improvements to airstrip	Eagle Plains Resources Ltd. and Canastra Resources Ltd.	stratabound hydrothermal mineralization along Ogilvie-Hart River Formation contact	1351 feet (412 m) in 8 holes	two widely spaced holes drilled through Ogilvie-Hart River Formation contact, confirming presence of stratbound mineralization; affirmation of distribution of chert and shale, including in low-lying areas (may cap mineralization preserved beneath the water table)	\$356,000	Termuende and Downie (1998), Hodder (1997)
1998	gravity and seismic reflection surveys, property reconnaissance prospecting and mapping	Eagle Plains Resources Ltd. and Canastra Resources Ltd.	stratabound hydrothermal mineralization along Ogilvie-Hart River fm contact, below present and paleo-water tables		continuation of prospective stratigraphy at shallow depths northeast of Orma hill; coincident with gravity anomalies	\$54,000	Power (1998)
1999	diamond drilling, property-scale mapping, regional reconnaissance mapping, prospecting, and sampling; clean-up	Eagle Plains Resources Ltd. and Canastra Resources Ltd.	stratabound hydrothermal mineralization along Ogilvie-Hart River fm contact, below present and paleo-water tables	1040 ft. (317 m) in 3 holes	drillhole north of Orma Hill intersects disseminated sphalerite in Devonian-Mississippian shale microbreccia overlying the Ogilvie Formation	\$273,000	Downie and Greig 2000
				total drill footage: 35,280 ft. (10,750 m) in 123 holes		total expenditures: \$4,927,000	

**TABLE 2 : Exploration Methods Employed on the Rusty Springs Property**

Method	Aim of survey/application	Results and comments	Recommendations
prospecting	locating mineralization	successful in locating silica-hosted vein-type mineralization	useful for following-up geochem
soil geochemistry	to locate potential mineralized zones and target trenches and drillholes	in spite of thick overburden and permafrost, effective in outlining near-surface mineralization	target top of Ogilvie Fm on remaining unsampled parts of property
stream geochemistry	location of new drill targets	creek sampling led to discovery of new showings local to property; geochemically anomalous drainages present in region	regional stream sediment sampling, targeting Ogilvie Fm. and overlying shale
trenching	to reach bedrock	mixed success with cat trenching: bedrock exposure not guaranteed; may require 2 seasons; environmental degradation problems	any further trenching may be more successful using an excavator
geophysics	targeting drillholes	most geophysical anomalies tested were coincident anomalies	
IP	targeting sulphides	resistivity anomalies outlined, but drill testing unsuccessful	not recommended without sound geologic framework
VLF-EM	targeting conductive sulphide horizons	many conductors outlined, but drill testing unsuccessful; may outline water-filled gougy fault zones	not recommended without sound geologic framework
magnetometer	??was this done??		not recommended without sound geologic framework
gravity	targeting more dense sulphides	anomalies outlined, but drill testing unsuccessful	several anomalies untested; not recommended without sound geologic framework
seismic	determining depth to favourable stratigraphic contact	unsuccessful, possibly imaged permafrost horizon	not recommended without sound geologic framework
drilling			
diamond drilling		reasonable drilling and recovery in oxidized mineralized zones using modern equipment and drilling techniques; drilling slow in resistant siliceous zones	recommended for future work; need high-powered rig, plenty of casing, mud, bits, core barrels, and patience
RC drilling		difficult drilling in oxidized mineralized zones; good drilling in resistant siliceous zones	not recommended

sampling, diamond drilling (1840 metres), and metallurgical sampling were also completed. Poor drill core recoveries once again hampered the effectiveness of the program.

A geological thesis was undertaken by D. Hansen, again emphasising a Mississippi Valley type model for the mineralization.

Exploration during the period 1975 to 1978 inclusive was funded by Rio Alto Exploration.

In 1979, detailed geological mapping, a soil geochemical survey, an Induced Polarization survey, and a gravity survey were completed. Joint funding of this work was by Rio Alto and E & B Explorations Ltd. of Calgary, Alberta.

A geological thesis by J. Bankowski indicated a hydrothermal exhalative nature.

In 1980, E & B Explorations Ltd. as operator, focused on the widespread mineralization discovered on the Orma Hill. Their program saw 1830 metres (6000 feet) of diamond drilling, bulldozer trenching, and some detailed geological mapping completed. Core recoveries were not significantly improved over previous years.

In 1982, Taiga Consultants Ltd. was contracted by Kenton Natural Resources to carry out a geological evaluation of the property and subsequently a comprehensive mineral exploration and diamond drilling program. During this period, 510 metres (1673 feet) of diamond drilling was completed, as well as a soil geochemical survey, a geophysical (VLF-EM) survey, detailed geological mapping of the property, and six trenches dug in order to define the style of mineralization.

The most recent research work, carried out by Jill Kirker (April 1982), strongly supports a hydrothermal origin for the mineralization.

In 1983, additional geophysical surveying and geochemical sampling were completed by Taiga Consultants Ltd. to detail geophysical conductors and geochemical zones previously outlined. During the fall of 1983, 488 metres (1600 feet) of diamond drilling were completed.

In 1986, Kenton Natural Resources Inc., as operator, drilled two holes in the valley bottom between the Mike and Orma Hills in order to test an I.P. anomaly delineated in 1979 by previous operators. This program consisted of 404m (1326') of drilling, and failed to intersect any significant mineralization. The drill was removed from the property following this short program.

The claims were gradually allowed to lapse, and in the spring of 1992, all claims comprising the property had expired. R.W. Termuende restaked the core area of the property on July 29th, 1992. 12 quartz claims were recorded, consisting of the Eric 1-6 and Jessica 1-6 claims.

A \$190,000 exploration program was completed during the 1994 season. The focus of the two-stage program was to carry-out further systematic exploration in the Mike Hill area, as well as undertake initial reconnaissance work in the region

surrounding the claim area. A total of 531 soil, 67 rock, and 36 silt samples were taken, over two separate control grids that were established on the property, covering the Mike Hill and Big Onion areas. Concurrent with the geological program, efforts were made to improve the infrastructure of the property, and included construction of a 530m (1800') airstrip, a 3.4km permanent road connecting the airstrip and camp areas, and 10km of drill-tote trails throughout the property. Environmental work was also undertaken in the Orma Hill area, with 8 man days spent collecting some 140 used fuel drums, refuse-burning, and general clean-up activities in areas of past development.

A two-phase trenching and diamond drilling program was carried out during 1995. Twenty-one drillholes totalling 1658 meters (5440 feet) were completed in the Mike and Orma hill areas, and a total of 400m of bulldozer trenching carried out in the Big Onion area. In addition, a 339-sample soil geochemistry survey was undertaken proximal to the Big Onion showing. A further 35 claim units were added to the existing property, bring the total area to 71 units. In addition, improvements were made to the airstrip, and an all-weather road network was completed to access all areas of the property. The total cost of the 1995 program was \$539,000. The most impressive mineralized interval intersected in 1995 occurred in hole RS95-M7, where a 15.3m interval from a hole drilled on the Mike Hill assayed 15.1 oz/ton silver, 3% copper, and 1.3% zinc, from 28.6-43.9m.

A 15-hole, 7600' (2320m) diamond drilling program was carried out on the property in 1996 at a total cost of \$560,000. The program was designed to test for the presence of deep-seated manto-type mineralization, which was interpreted to lie beneath high-grade "chimney" veins exposed on surface in the Mike and Orma Hill areas. In addition to geological work, significant improvements were made to property infrastructure, with three km of new roadwork completed, and the airstrip extended to 2000' (600m). Significant to the 1996 program was the discovery of stratabound mineralization, apparently over much of the property area, and beyond. As a result of the new interpretation, 478 quartz claim units were staked in the region, covering all favourable stratigraphy in the immediate area.

The \$355,000 1997 program utilized a reverse circulation drill in an attempt to mitigate drilling problems associated with the highly abrasive cap rocks overlying the mineralized horizon. While the drill was performed better in the siliceous ground, there were problems with recovery and sample contamination within the mineralized zone. Two of the holes confirmed the presence of stratabound mineralization at the Hart River – Ogilvie Formation contact over a large area. During 1997, R. W. Hodder, Ph.D., P.Eng., visited the property and examined existing drill core, outcrop, trenches and technical data. He concluded that "*The limonitic interval at Rusty Springs is a resource of hundreds of millions of tons, but of very subeconomic amounts of base or precious metals... the limonitic interval and it's enclosed quartz veins and lamellae are however vital symptoms that ore forming processes existed for major deposits of silver-lead-zinc and that deposits of this type cluster in districts of enormous potential*". Hodder also recommended focusing on locating sulphides below the present and paleo water table.

The \$54,000.00 1998 program involved a combined shallow seismic and gravity geophysical survey. The surveys were run from the northeast flank of the airstrip east across the low lying swampy area. The survey defined a coincident positive Bouger gravity anomaly and seismic reflection profile interpreted to be related to a shallow sulphide body at the same stratigraphic horizon as sulphide mineralization defined in 1996 - 97.

Eagle Plains Resources continued mineral exploration at the Rusty Springs property continued in 1999 with a \$273,000.00 field program. A three hole helicopter supported, diamond drill program to test for stratabound mineralization in the

area of Orma Hill was carried out concurrently with a property and regional scale geologic mapping program under the direction of Charlie Greig.

Results from the 616.9 m (2024 feet) 1999 diamond drilling program at the Rusty Springs property were largely inconclusive because none of the holes could be completed to target depth. Two of the three holes intersected quartz carbonate crackle breccia within a strongly silicified (cherty?) black mudstone that overlies the mineralized horizon elsewhere on the property. While no massive sulphide or Katshat horizon was intersected, a mineralized breccia zone was encountered between 229.2 and 264.9 meters in hole RS99-01. Mineralization consisted of finely disseminated to patchy orange – red sphalerite associated with fine quartz crackle breccia and coarser collapse type breccia. The host rock was silicified black mudstone. The best mineralized intervals were 226.8 to 233.4 meters which averaged 2819 ppm zinc over 6.6 meters, and 249.8 to 264.9 meters which averaged 3100 ppm zinc over 15.1 meters. The brecciation, strong silicification and mineralization are consistent with the nature of the cap rocks associated with the Katshat horizon elsewhere on the property. Casing was left in all three of the drillholes to facilitate future deepening of the holes to intersect the mineralized horizon.

## **REGIONAL GEOLOGICAL SETTING**

The area mapped lies within the northernmost part of the Cordilleran orogenic belt, known locally as the Taiga -Nahoni foldbelt, where Precambrian to Cretaceous predominantly sedimentary rocks of the eastward and northward tapering North American miogeocline were deformed in latest Cretaceous to Tertiary time (Norris 1996, Lane 1998). The area was first mapped by Norris (1981), who outlined a structural culmination, in part coincident with his Porcupine Anticline, cored by rocks of the Lower and Middle Devonian Ogilvie Formation. Norris (1981) shows stratigraphically lower rocks of Early Paleozoic, Cambrian, and Proterozoic age bounding the west side of the culmination and brought up by mainly west vergent contractional faults.

## **PROPERTY GEOLOGY (Fig.2, Fig.3)**

Nine map units, ranging in age from Proterozoic to Cretaceous, correspond largely with those mapped by previous workers (e.g., Chernoff 1976, Kirker 1980a, Tempelman-Kluit 1981; Fig. 2). Ages of the map units were taken mainly from Norris (1981, 1996). Exposure is generally poor near the valley bottom and consequently the focus for property-scale geologic mapping was on the rocks underlying surrounding ridges. The geology in the immediate vicinity of the mineralized and altered zones at Rusty Springs, which crop out at lower elevations in the vicinity of two lower hills, named the Mike and Orma hills, was examined briefly.

### Lower to Upper Proterozoic rocks

Rusty weathering fine-grained sandstone (quartzite), interbedded with maroon and local green siltstone and silty mudstone (siltite), occurs in a northerly trending belt in the southwesternmost corner of the area mapped. The siliciclastic rocks, which were only briefly examined, appear to be conformable with steeply east dipping Lower Paleozoic dolostone and quartz-rich sandstone to their east.

### Lower Paleozoic rocks

Like the older rocks which they appear to overlie conformably, rocks of probable Late Cambrian through Early Devonian age occur in a northerly trending belt along the west margin of the map area. The Lower Paleozoic rocks consist of white weathering dolostone, rusty weathering quartz-rich sandstone (quartzite), and siliceous fine grained clastic rocks, including green and maroon siltstone and silty mudstone (siltite). Rocks of similar general appearance occur to the north, but were neither examined nor differentiated from the older siliciclastic rocks. The Lower Paleozoic rocks are inferred to be in thrust contact with younger Paleozoic and Mesozoic rocks to the west, although a down-to-the-east normal fault was mapped along trend to the south by Norris (1981). The presence of inferred thrust is supported by the marked easterly vergence of folds in the area.

## Lower and Middle Devonian Ogilvie Formation

Pale grey weathering dark grey dolostone and subordinate limestone and argillaceous rocks of the Ogilvie Formation underlie the central part of the Rusty Springs property in the core of the Porcupine-Rusty Springs anticlinorium (Figs. 7, 8). They form common talus slopes on the flanks of Orma and Mike hills, but outcrop is scarce, even on roads and cat trails. Dolostone is fetid, and commonly brecciated, veined, and(or) vuggy. Breccia cements consist mainly of dolomite and sparry calcite with local quartz; vugs are commonly lined with calcite and quartz, and veinlets are of similar mineralogy. Another common constituent of Ogilvie Formation breccias is pyrobitumen—it is commonly intergrown with dolomite cements and always associated with quartz and(or) calcite spar (Kirker 1982); it also locally coats vugs. Dolomite crystals in dolostone are typically fine- to medium-grained and locally coarse-grained, with coarser-grained varieties typically weathering a paler grey colour. Locally, weakly dolomitized limestone contains recognizable brachiopods, ostracods, corals, and crinoids (Hansen 1979, Davis and Aussant 1982), although no diagnostic fossils have been reported. Float boulders and the few outcrops of the Ogilvie Formation suggest that it is not well stratified, but bedding is more apparent in diamond drill core, particularly where brecciation is less intense, and bedding to core axis angles typically suggest that the strata in the vicinity of Mike and Orma hills are gently dipping. Mainly on the basis of their contained fauna, Hansen (1979) interpreted the dolostones of the Ogilvie Formation as a shallow water “reefal” unit, while Kirker (1982) suggested a shallow water shelf environment. The base of the Ogilvie Formation at Rusty Springs is not exposed, but a drill hole between Mike and Orma hills penetrated about 210 metres (probable true thickness) of dolostone, with local interbedded shale and rare limestone and quartzite (Chamberlain 1986).

At the top of the Ogilvie Formation at Rusty Springs is the informally named “Katshat unit”, a recessive, gossanous oxide- and clay-rich unit which corresponds to a significant degree with the mineralized zones on the property. In general the unit appears to be stratabound, separating the dolostone from overlying siliciclastic rocks, but in detail its contacts are highly irregular. The Katshat unit most likely represents altered and mineralized Ogilvie Formation limestone—it is discussed in more detail below.

## Devono-Mississippian fine-grained siliciclastic rocks

Disconformably overlying the Ogilvie Formation are siliceous mudstone, slate, shale, siltstone, and rare limestone of probable Devono-Mississippian age. The rocks were assigned by Norris (1981) to the Hart River Formation (Early and Late Carboniferous age), but they are more likely correlative with fine grained clastic rocks, such as the Upper Devonian Canol Formation, the Unnamed shale, the Upper Devonian and Lower Carboniferous Ford Lake shale (Norris 1981, 1996), and the Kayak Formation (Richards et al. 1996), because the Hart River consists mainly of limestone (Norris 1981, 1996). Herein the rocks have been assigned to the Unnamed shale.

The lowermost rocks in the sequence, best exposed on Orma and Mike hills and referred to locally as black ‘chert’, are perhaps more accurately referred to as a silicified and(or) siliceous mudstone. Thin laminations and recrystallized radiolaria are locally preserved (Hansen 1979). The siliceous rocks are up to 40 metres thick (Hodder 1997) and are commonly veined and brecciated; veins and breccia matrices consist mainly of quartz, calcite, and dolomite. The brecciated siliceous rocks appear in most places to cap the mineralized Katshat unit of the uppermost Ogilvie Formation, and black siliceous(?) fragments are locally

a common component of the dolostone breccias that commonly comprise upper Ogilvie Formation rocks beneath the Katshat unit.

Up-section from the siliceous rocks, and comprising the bulk of the rocks assigned to the Unnamed shale, are relatively recessive pyritic, carbonaceous shale, mudstone, silty mudstone, and local thin- to medium bedded, poorly sorted fine grained litharenite. They are generally thinly bedded, and typically siliceous, although local calcareous shale was also noted. Local true slate and rare dark grey, fetid and laminated algal limestone occur not far above its contact with the Ogilvie Formation. Erosion of this part of the unit, which is as much as 500 metres thick, has led to the broad and open drainage basin within which the Rusty Springs property sits.

The transition of the fine grained clastic sequence to the overlying mixed carbonate and clastic unit is commonly marked by the presence of thin to medium bedded siliceous fine sandy siltstone or fine grained sandstone. These rocks are typically pale grey and locally rusty weathering up close, but appear very dark from a distance because of a common covering of black lichen.

#### Upper Carboniferous and Permian(?) limestone and fine grained calcareous and siliceous clastic rocks

Medium bedded, pale grey weathering, medium to dark grey sandy and locally pebbly fetid limestone and rare dolostone characterize this unit. The limestone commonly contains irregular dark grey chert nodules and occurs in several(?) horizons of amalgamated beds that are up to several tens of metres thick. They form many of the better outcrops in the area and because of their resistant character, they underlie many of the ridges surrounding the broad upper drainage basin of Carrol Creek. The upper limit of the map unit is defined by presence of the uppermost continuous limestone sequence, while the transition from the underlying siliciclastic sequence is commonly marked by scattered float blocks of pebbly limestone. The pebbles are typically round to sub-round and are dominantly chert. Pebbly lithologies are more common to the southwest, whereas to the east, sandy limestone is more common and pebbly limestone occurs only locally. In addition, a limestone horizon containing abundant *in situ* corals was noted in the east but not to the south or southwest, and composite limestone horizons appear somewhat thicker (up to 50-60 metres) and may contain thicker-bedded to massive layers of up to 15 metres thickness. In spite of the predominance in outcrop of pebbly and cherty limestone, a significant portion of the map unit consists of relatively recessive, variably calcareous fine grained clastic rocks. They include dark weathering, thin bedded and laminated siliceous or calcareous silty mudstone, and calcareous to siliceous shale, as well as local fine grained siliceous sandstone and siltstone. The total thickness of the limestone and associated clastic units is about 550-700 metres.

The rocks of this sequence have been included previously in the Upper Carboniferous Ettrain Formation, but Pennsylvanian and Permian fossils have been reported from within the area mapped, and so it is probably longer-ranging and likely includes rocks mapped as Jungle Creek Formation by earlier workers. If so, it is difficult to distinguish Ettrain from Jungle Creek in the field.

#### Jurassic and Lower Cretaceous dark weathering siliciclastic rocks

Lying conformably above the sequence containing the resistant grey carbonates is a dark weathering package of shale,

silty mudstone, and sandstone of approximately 600 metres thickness. Included in this map unit are rocks that Norris (1981) assigned to the Jurassic and Lower Cretaceous Kingak, Porcupine River, and Husky formations. The lower part in the Rusty Springs area consists of common pale to medium brown weathering silty mudstone with local buff-weathering carbonate layers, and dark brown weathering shale. Near the east-central part of the area mapped, near its base, the sequence includes a thick (up to 46 metres, Chernoff 1976) oolitic hematite-magnetite siliceous iron formation. Several kilometres along strike to the north, and at the same stratigraphic level, the base of the unit is marked by massive black carbonaceous and siliceous mudstone and silty mudstone. Similarly resistant siliceous rocks mark the upper part of the unit, which underlies many of the highest ridges in the south and east parts of the area mapped. They are very dark weathering and consist mainly of blocky weathering, medium grained feldspathic cherty quartz arenite and carbonaceous fine grained siliceous litharenite.

#### Lower Cretaceous shale, siltstone, and quartz arenite

The two units bounding the east side of the map area were taken from the mapping of Chernoff (1976), who shows numerous overturned beds within their bounds. He assigned the shale, siltstone, and quartz arenite comprising the units to the Cretaceous Marten Creek and Goodenough (*sic*) formations. Norris (1981) assigned them a Lower Cretaceous age, and included them in his "KWC" unit and the Mount Goodenough Formation.

#### Structural geology

Folds are the dominant structural feature in the map area, and wavelengths of the typical east vergent open to tight and locally overturned folds are on the order of 1-5 kilometres. The folds occur across the crest of an approximately 20 kilometre wide, northerly trending and doubly-plunging anticlinorium centered on the mineralized showings at Rusty Springs. The east side of this domal feature corresponds to the Porcupine Anticline of Norris (1981). Brittle faults are common on the property, and have been intersected in drillholes and interpreted from geophysical surveys and surface features (such as linear stream patterns), but none of these faults appears to offset map units at the property scale. The plunge reversal that corresponds with the mineralized area and which has been interpreted by some (e.g., Chernoff 1976) to have been associated with a brittle fault, appears, from the map patterns, to be fold-related and the consequence of some deeper-level structure, such as a lateral ramp.

Several property-scale cross-sections have been prepared previously, beginning with that of Chernoff (1976), and followed by Kirker (1980) and Tempelman-Kluit (1981). Chernoff (1976) shows a large-scale easterly-overturned antiform which is centered on the Rusty Springs showings and which he interprets as being cored by intrusive rocks and floored by north-trending, east-directed thrust faults. In contrast, Kirker (1980) and Tempelman-Kluit show inferred, north-trending faults, but interpret them as west-vergent contractional faults. They also show related folds with generally open geometries (Kirker 1980, Tempelman-Kluit 1981). Our cross-sections, based on improved bedding control compiled in part from previous work and benefitting from drillhole control, suggests that the structural setting is somewhat more akin to that shown by Chernoff (1976), in that the transport direction across the anticlinorium is toward the east. An east-directed transport direction is also more in accord with the regional sense of vergence.

Speculatively, the area may be floored by a large-scale east-vergent contractional fault, in part as envisioned by Chernoff (1976). Key to this interpretation are the steeply dipping and overturned Cretaceous rocks along the east side of the

area mapped by Chernoff (1976). They may represent the eastern, overturned limb of the northern Porcupine Anticline, and may be floored by an inferred southern continuation of an east vergent contractional fault shown by Norris (1981) as bounding a panel of Upper Proterozoic to Lower Paleozoic rocks on their east side about 15-20 kilometres to the north-northeast. If this is the case, the doubly plunging anticlinorium underlying the Rusty Springs area may reflect the influence of a deep-seated feature, such as a lateral ramp, along the inferred contractional fault.

## Mineralization

Although exploration models utilized at Rusty Springs have tended to exclusively target either stratabound or discordant styles of mineralization (e.g., Mississippi Valley-type or Irish Plains-type for the former, hydrothermal veins for the latter), there appears to be good evidence for both styles on the property, and they appear to be genetically related. Both styles of mineralization are found almost exclusively in the upper Ogilvie Formation and in the vicinity of the Mike and Orma (Hansen and Bankowski 1979), and their spatial association, similar geochemical signatures, and their association with similar brecciated and dolomitized zones, suggests a genetic link. Potential rests mainly with the stratabound mineralization, which may have greater thickness, much greater continuity, and can be much more readily explored for.

### Vein-type mineralization: the Orma zone

Mineralization at the Orma zone, on the northwest flank of Orma hill, has been the focus for the bulk of the exploration work at Rusty Springs. Up to the 1990's, virtually all of the drilling on the property occurred there. The zone has yielded many of the highest grades in grab samples, trenches, and drill core (e.g., DDH80-1: 583 gm/t Ag, 8.23 % Pb, 1.48 % Cu over 6.5 metres) and trenching and drilling have confirmed that it is a discontinuous vein and vein stockwork zone which trends northwest and dips steeply. Vein-type mineralization also appears to be present locally at Mike Hill, with the difference that relatively high Zn and trace Au values commonly accompany the Ag, Pb, and Cu common to mineralization at the Orma zone (Downie 1994; e.g., DDH95-07: 518 gm/t Ag, 0.77 % Pb, 3.0 % Cu, and 1.3% Zn over 15.3 metres).

Veins consist of massive galena-tetrahedrite (tennantite?, as is suggested by elevated As:Sb ratios in some assays, Liedtke (1980)), locally up to 1.0 m thick, which assay roughly 10-50 ounces per ton Ag. The veins are contained within a broader, commonly oxidized mineralized and altered zone (in part a vein stockwork) of up to 6 or 7 metres thickness. The altered zone typically assays 30 to 60 grams ounces per ton Ag (Davis and Aussant 1982). Alteration within Ogilvie Formation carbonates, as described by Bankowski (1980b), is characterized by silica replacement, dolomitization, local brecciation, sanding (silicic alteration?), and decomposition (supergene alteration), and is manifest in part as a darker grey colour of the host rocks. The margin of the altered zone has a northwest trend, subparallel to that of the mineralized zone, and it appears to terminate, or turn bedding-parallel, to the southeast at the contact with overlying siliciclastic rocks (Bankowski 1980b). Minerals identified from the oxidized zones include smithsonite, cerussite, malachite, azurite, aurichalcite, pyrolusite, hemimorphite, plumbojarosite, gibbsite, valentinite, and natroalunite (Hansen 1979, Kirker 1980b); sphalerite and pyrite are also preserved locally with galena and tetrahedrite in siliceous vein and vein-breccia material.

### Stratabound mineralization: the Katshat unit

Near the end of the 1996 exploration program, stratabound mineralization along the contact between the Ogilvie Formation and overlying Devonian-Mississippian siliciclastic rocks became the principal exploration target (Termuende and Downie 1997). Almost all holes drilled in footwall Ogilvie Formation dolostone had essentially been barren, and with relatively thick oxidized mineralization cored at the contact in several previous drillholes that were collared in hangingwall siliciclastic rocks, it was realized that substantial potential existed for stratabound mineralization. It was also recognized that the most extensive geochemical anomalies on the property coincided with the contact, and that many drillholes targeting them had been collared in the strongly oxidized mineralized material—these holes had been plagued by poor core recoveries.

The oxidized material common to the upper contact of the Ogilvie Formation was referred to locally as the Katshat unit. It consists of strongly leached, porous limonitic to kaolinitic material with an earthy, gougy consistency, and is similar in appearance to the oxidized material surrounding discordant mineralization. It is typically 20 to 40 metres thick, and although it appears stratabound at the property scale, in detail it is irregular and discordant. Many of the minerals noted above as occurring in the Orma zone are also common in the Katshat unit. X-ray diffraction studies indicate that much of the Katshat material consists of granular Fe, Mn, Ag, Pb, An, Cu, Ba, Al, P, and V oxide, carbonate, sulphate, and silicate mineral species, as well as quartz veinlets and laminae locally containing sulphides and sulphosalts like those in Orma zone veins and vein stockworks (Hodder, 1997). The Katshat unit is invariably overlain by brecciated and veined siliceous or silicified mudstone and chert of probable Devonian-Mississippian age, which caps and in part has protected it from erosion. It is underlain by Ogilvie Formation dolostone, also typically brecciated and veined. The Katshat unit is strongly anomalous in Ag, Cu, Pb, and Zn over broad intervals and across a wide area (for e.g., 1.1 gm Ag, 881 ppm Cu, 139 ppm Pb, 3301 ppm Zn over 19.1 m in hole RS96-04 from the southwest part of Mike hill, and 1.6 gm Ag, 1475 ppm Cu, 1321 ppm Pb, and 2701 ppm Zn over 22.2m in hole RS96-14 from the south end of the airstrip on Orma hill;). Results such as these suggest the possibility of tremendous continuity and potential, but the oxidized nature of the mineralization and the subeconomic grades also suggest that the preferred target be unoxidized portions of the horizon below the present and(or) paleo- water table (Hodder 1997). Unoxidized Katshat unit was the target of the latest drill program, which attempted to test the upper Ogilvie Formation to the east and south of Orma hill (Fig.3) Results were mixed. Because of problems penetrating the very resistant siliceous and brecciated rocks which overlie the upper Ogilvie Formation and cap the Katshat horizon, the mineralized horizon was never reached. However, the presence of the siliceous rocks suggests that a strong stratabound mineralizing system existed well away from the surface exposures on Mike and Orma hills, and as such, the new information confirms that the Rusty Springs system is very large, and that it has significant potential remaining to be tested.

### Timing of mineralization

The interpretation that Rusty Springs is a Mississippi Valley-type deposit related to karsting along the upper Ogilvie Formation contact suggests that the mineralizing event was likely bracketed by the ages of the Middle Devonian rocks below and the Upper Devonian to Mississippian rocks above. On the other hand, the discordant nature of mineralization and alteration at Rusty Springs indicates that it postdates deposition of the Lower to Middle Devonian Ogilvie Formation and at least the lowermost part of the overlying Devonian-Mississippian section. In addition, one can argue that evidence such as the lack of

obvious cleavage development in the Ogilvie Formation dolostones, which contrasts sharply with that common to most rocks across the property, including other carbonates, suggests that the mineralizing event may even have postdated much of the latest Cretaceous to Tertiary deformation affecting the area (alternatively, it is possible that this may reflect a contrast in competency between the more competent silica-altered and dolomitized rocks associated with mineralization and other less competent lithologies, or that a more subtle stylonitic cleavage exists in the dolostones—further study is needed). The parallelism of the Orma zone with structural trends (a fold axial plane?) and localization of Katshat-style mineralization in anticlinal hinge zones at Orma and Mike hills may also support the hypothesis that mineralization post-dated deformation. A relatively young age is also supported by the rare occurrence of discordant metre-scale vein-breccia bodies of quartz or Fe-carbonate at higher stratigraphic levels (Carboniferous to Permian) in the area surrounding Rusty Springs, and by limited Pb isotope data suggesting which that approximate those of Cordilleran Ag vein deposits of Late Mesozoic age (Kirker 1982).

### Genesis

As mentioned above, several deposit models, including those for MVT and hydrothermal replacement along a karsted surface, have been employed in an effort to aid exploration at Rusty Springs. Poor exposure and consequent lack of local bedding control has hindered the collection of evidence with which to evaluate the various models, as has leaching and oxidation of the mineralized zones and dolomitization of footwall rocks. However, discussion of some of the existing evidence is worthwhile so that some models may be critically evaluated and perhaps ruled out, and others put forward in the hope that they aid exploration.

### Mississippi Valley-type

Few, if any, of the textural features distinctive of MVT type deposits (e.g., Leach and Sangster 1993) have been positively identified on the property. For example, although the breccias common on the property have been interpreted as solution collapse features (e.g., Hansen 1979, Hodder 1997), cements and infillings of carbonate and local quartz are either massive or encrusted symmetrically around breccia fragments (e.g., Kirker 1982). There is no evidence for infilling by internal sediment, which would be strongly suggestive of a karst environment. Stratigraphic evidence also appears to argue against a karst environment. No regolith or is preserved along the contact between the Ogilvie Formation and the overlying siliciclastic rocks that would indicate subaerial exposure, and even evidence for uplift, such as the presence of coarse grained clastic rocks, is lacking. According to Liedtke (1980), very little relief exists on the contact, and if anything, subsidence is indicated: the stratigraphic transition is from a shallow water environment in which platformal carbonate was deposited, to a deeper water environment in which basinal shales were deposited.

Differences from classic MVT deposits also exist in the geochemistry and mineralogy at Rusty Springs, as has been noted by many previous workers. The high copper and silver contents, as well as low Zn:Pb ratios are generally atypical of MVT deposits (Leach and Sangster 1993), as are locally very high As and Sb values and the high Al values occurring in the Katshat unit (Termuende and Downie 1997). A geochemical fingerprint such as this is more consistent with an epithermal origin for metals within the host unit. Similar arguments can be made on mineralogic grounds, with the siliceous character of alteration, particularly in the hangingwall, and the common presence of tetrahedrite and argentiferous galena, which are more diagnostic of vein rather than stratabound Ag-Pb-Zn deposits, in the mineralized zones. Fluid inclusion and sulphur isotope data from quartz,

calcite, and sphalerite at Rusty Springs are also more comparable to those from epithermal deposits than from those of MVT (Kirker 1982).

Regionally, the evidence also argues against an MVT setting. As Hodder (1997) notes, it is significant that the Ogilvie Formation at Rusty Springs is comprised largely of dolostone in an area in which limestone generally predominates. Even within the Ogilvie Formation itself, the regional dolomitization common to MVT districts appears to be absent--Norris (1996) describes only local dolomite beds in the lower part of the Ogilvie Formation in measured sections farther south in the Ogilvie Mountains.

In spite of the arguments against the presence of MVT mineralization, it remains possible that the mineralization and alteration evident on the Rusty Springs property may simply be the distal expression of a more typical MVT system origins lie in a hydrothermal karst system rather than a meteoric or meteoric-hydrothermal one (c.f. Leach and Sangster 1993).

#### High-temperature, carbonate-hosted massive sulphides: manto-chimney complexes

The mineralizing system at Rusty Springs bears some of the features of high-temperature, carbonate-hosted massive sulphide deposits (Titley 1993), which are also commonly referred to as manto-chimney complexes, and are rich sources of base and precious metals. This type of deposit, although occurring in quite varied structural or stratigraphic settings, is typically wholly or partially stratabound, commonly contains abundant pyrite, and contains Pb and significant Ag. Copper and Au can be present but are less common than Ag-Pb-Zn, and enrichment in one or the other of Cu-Pb-Zn can be variable. The deposits are generally thought to occur by replacement processes, initiated by hot fluids and(or) gases, above or near centres of thermal activity, and so intrusions are commonly (though not always) spatially associated. Vein, skarn, and even porphyry copper deposits may be closely associated the manto-chimney ores, and it is generally accepted that all are genetically related to the associated intrusions (Titley 1993).

The potential for manto-chimney deposits at Rusty Springs was initially recognized by Termuende (1996). The few preserved hypogene ore minerals recognized at Rusty Springs, such as galena and tetrahedrite, are common in the manto-chimney class, and the silica alteration common on the property is also commonly peripheral to ore or in this deposit type, or at least to districts in which such deposits occur. In addition, dolomitization is known to play a role in the formation of many high-temperature, carbonate-hosted deposits, and breccia bodies are also common to these systems (Titley 1993). The apparent controls on mineralization at Rusty Springs, such as the overlying impermeable fine grained siliceous shale cap, and perhaps the anticlinal fold hinges at Mike and Orma hills, also bear similarities to some manto-chimney deposits (e.g., Tombstone, Arizona; Titley 1993). This factor of predictability is an important advantage in exploration for manto-chimney ores, since they are known to be difficult to explore for. One of the main arguments against the application of the manto-chimney model at Rusty Springs is the lack of direct evidence for intrusive rocks, either on the property or in the region, although Chernoff (1976) shows an inferred intrusion at depth below the domal core of the Rusty Springs antiform. The nearest known plutons to Rusty Springs are Devonian(?) in age and outcrop to the north in the vicinity of Old Crow ( Woodsworth et al. 1991).

#### Other economic potential in the vicinity of Rusty Springs

Little in the way of significant mineralization has been found in the immediate area around Rusty Springs, but recent work and a reevaluation of work done previously indicates that some potential exists and that it should be tested. For example, in

the most recent drilling, an interval approximately 40 metres thick within the Devono-Mississippian pyritic shales that overlie the Ogilvie Formation was highly anomalous in zinc; it included intersections of 7 and 15 metres which returned nearly 3000 ppm Zn. Although the hole did not reach its target, it is estimated that the Zn-rich zone lies approximately 100-150 metres up-section from the Ogilvie Formation, at about 250 metres depth. The zone occurs within a siliceous or weakly silicified carbonaceous pyritic mudstone, and sphalerite occurs as fine to medium grained honey brown disseminations, both within mudstone clasts, and within matrix host rocks to zones of quartz or quartz-carbonate microbreccia. The pyritic and locally zinc-rich shales may be the source for the gossanous springs near the base of the north end of Mike hill which lend their name to the Rusty Springs property and, in fact, sediment issuing from the springs themselves was highly anomalous in zinc (Chernoff 1976). This suggests further that the recessive shale package may have potential for hosting Zn deposits, either similar in character to Rusty Springs, or perhaps of the sedex type, much as rocks of similar age, character, and tectonic setting farther southward in the Cordillera do (e.g., Macmillan Pass area, Y.T.; Gataga district, B.C.; Dawson et al. 1991). One might begin to evaluate this potential immediately south-southeast of the area mapped, where rusty creeks and springs, similar in appearance to those at the Rusty Springs property, were noted in the drainage that lies in the recessive core of Norris' (1981) Porcupine Anticline. The springs likely emanate from rocks correlative with the recessive and pyritic Devono-Mississippian rocks that overlie the Ogilvie Formation in the area mapped.

With regard to other possibilities, rare iron carbonate breccia and siliceous veins and vein-breccias were noted in outcrop or float while mapping the surrounding ridges, but none bore visible sulphides, appeared extensive or was accompanied by significant alteration. About 40 kilometres farther south, however, at the Pama (Bern) occurrence, which lies just inside the western boundary of the proposed Fishing Branch Protected area, an impressive, steeply dipping, north-northwest trending quartz-carbonate breccia zone that is hosted by carbonates can be traced for greater than two kilometres. It is outlined by a broad and intense soil geochemical anomaly (O'Donnell 1974) and near its southern end it contains tetrahedrite, copper oxides, and zinc and lead sulphates that bear some similarities to mineralization at Rusty Springs. The Pama property has never been drill-tested, yet smithsonite-rich samples yield assays of up to 47.80% Zn. Although it is hosted in carbonates and has at least some mineralogic similarities to Rusty Springs, no truly convincing evidence was found at the Pama that was suggestive of a significant element of stratigraphic control to mineralization. The breccia zone is hosted by limestone that is probably correlative with the uppermost limestones in the vicinity of Rusty Springs (Upper Carboniferous and Permian(?); considerably younger than the Ogilvie Formation). The breccia appears to dip steeply to the east-northeast, and lies subparallel to the steeply dipping eastern limb of what appears to be a gently southerly plunging, asymmetric, east vergent antiform. The breccia appears to be hosted entirely within limestone, and the limestone is only very locally dolomitized, which is in sharp contrast to Rusty Springs, where the better part of the Ogilvie Formation is dolomitized. Overlying the limestone is a sequence of relatively recessive, fine grained black carbonaceous rocks that appear to be capped by more resistant siliceous sandy beds. The sequence is similar in appearance to the Jurassic and Lower Cretaceous rocks along the east margin of the area mapped at Rusty Springs.

## 2001 PROGRAM AND RESULTS

Mineral exploration at the Rusty Springs property continued in 2001 with a \$16,500.08 field program. Geological work consisted of helicopter reconnaissance and rock sampling in areas of interest identified by past programs. A total of 11 samples were collected. All samples were shipped to Eco-Tech Laboratories in Kamloops, B.C. where they were analyzed for 30 element ICP using aqua-regia digestion. High-grade samples were further fire-assayed. The samples were also analyzed for Germanium content. All samples were collected, handled, catalogued and prepared for shipment by Toklat Resources and Eagle Plains Resources staff.

Personnel and equipment were mobilized to the Rusty Springs camp on June 10, 2001 by Fireweed Helicopters out of Dawson City using a Bell 206 Jet Ranger. Extra fuel was hauled to the airstrip at 150 Mile on the Demster Highway.

The results of the 2001 program continue to confirm the spectacular nature of the mineralization found throughout the Rusty Springs property area. Samples TTRS01R02 and TTRS01R08 returned Bonanza silver grades from float samples collected in the area of the Trog 2 and Conner 4 claims respectively. Two other samples collected in the area of the Trog 1 – 4 claims also returned highly anomalous values. Sample TTRS01R01, a sample of carbonate breccia float with massive coarse grained galena returned values of 98.2 g/t Ag and 3.22% Pb. Sample TTRS01R03 returned values of 62.3 g/t Ag, 3.07% Cu, 13.2% Pb and 1.44% Sb. Both of these samples also contained As values in excess of 10000ppm. Sample TTRS01R04, a sample of massive oolitic iron collected from the Iron Formation in the area of the Trog 50 claim returned anomalous Zn(454ppm) and Ba(315ppm) values. TTRS01R05, a sample of dolomite float with malachite replacement malachite returned values of 88.9 g/t Ag and 40.7% Cu. TTRS01R06, collected in the area of the Trog 20 claims, returned anomalous Ba values(1490ppm) from a piece of limonite breccia float. TTRS01R07, a sample of rounded, transported Katshat material collected from a drainage on the Conner 1 claims, returned values of 1.2 g/t Ag and 159ppm Zn. TTRS01R10, a sample of dolomite breccia float with aurichalcite collected on the Matt 4 claims returned Zn values in excess of 10000ppm. TTRS01R11, a sample of zinc mineralized crackle breccia hosted by silicified black shale from DDH 99-01 at a depth of 256.4m returned values of 7.4 g/t Ag, 698ppm Pb and 823ppm Zn. None of the samples returned anomalous Ge values.

## CONCLUSIONS AND RECOMMENDATIONS

In spite of many years of sound exploration work, the genesis of the Rusty Springs prospect remains incompletely understood, yet its considerable potential for a large-tonnage Ag-Pb-Zn-Cu deposit remains inadequately tested. The extent of the mineralized and altered rocks at Rusty Springs suggests that the hydrothermal system is large and, in fact, limited to the altered and brecciated zones in the upper Ogilvie Formation have yet to be established, with the possible exception of on the northeast (Bankowski 1980b). The size of the mineralizing system, together with its apparent stratabound nature, its commonly significant but subeconomic thicknesses, its local high grades, and its potential for supergene enrichment, indicates that Rusty Springs remains an attractive exploration target. Future exploration should be drill-oriented and should target the uppermost Ogilvie Formation beneath Devonian-Mississippian fine grained clastic rocks to the south and southeast of Mike and Orma hills.

Results from the 1999 diamond drilling program at the Rusty Springs property were largely inconclusive because none of the holes could be completed to target depth. The abrasive and generally fractured nature of the rocks overlying the mineralized Katshat horizon have been an obstacle to drill testing throughout the history of exploration on the property. The objective of the 2002 program is to complete the three holes drilled during 1999 using a specialized drill and specific hole conditioning techniques designed to optimize recovery and depth limits in abrasive and fractured ground. Hytech Drilling Limited, a drill contractor based in Smithers, B.C. has a heliportable customized hydraulic diamond drill that has been built specifically for diamond drilling in bad ground conditions. In concert with this drill, it is further recommended that cement should be used to stabilize the drillhole walls to prevent hole collapse. All holes should be collared using the maximum combination of casing and large diameter (HQ2) drill pipe. After advancing through bad ground conditions (sand faults, rubble zones), the drill rods should be removed, the hole cemented, and then recored with smaller diameter drill rod (NQ2 or BQ2). The cement will form a stable wall and prevent the incursion of black sand and other rubble around the rods and drill string. Drilling tools designed for abrasive ground conditions should be used, including hexagonal core barrels.

A 1500 meter diamond drilling program is recommended for the Rusty Springs property. The three holes collared in 1999 should be redrilled and completed to projected Katshat Horizon target depths defined by past drill programs and the 1999 mapping results. The 1500 meter proposal includes footage to follow up stratabound massive sulphide mineralization with a single step out hole. A proposed budget for the recommended work follows :

PROPOSED BUDGET FOR 2002 RUSTY SPRINGS EXPLORATION PROGRAM

Diamond Drilling :1500m @ \$100.00/m .....	\$150,000.00
Personnel / Contractors.....	\$35,000.00
Consultants .....	\$10,000.00
Air Charter (Helicopter) .....	\$20,000.00
Air Charter (Fixed Wing).....	\$45,000.00
Mob/Demob.....	\$6,000.00
Airfare/Accommodation .....	\$6,000.00
Analytical.....	\$10,000.00
Meals/Grocery .....	\$10,000.00
Truck/Equipment Rentals .....	\$10,000.00
Fuel (Diesel, Gasoline, Propane).....	\$10,000.00
Field Supply/Camp Materials.....	\$8,000.00
Communications .....	\$3,000.00
Miscellaneous.....	\$10,000.00
Report/Reproduction.....	<u>\$7,000.00</u>
	Sub-Total : \$340,000.00
	10% Contingency : <u>\$34,000.00</u>
	TOTAL: \$374,000.00

NOTE : all values in \$CAN

**REFERENCES**

- Aussant, C., 1983. Geological Report on the Rusty Springs Property, Yukon Territory. Unpublished report for Kenton Natural Resources Corporation, 12 p.
- Bankowski, J., 1980a. Genesis of Base Metal Sulphide Occurrence, Rusty Springs Prospect, Ogilvie Mountains, Yukon Territory. Unpublished B.Sc. thesis, University of Western Ontario, London, 70 p.
- Bankowski, J., 1980b. Report of Exploration Programme conducted 19 May–1 August, 1980 for Rio Alto Exploration Ltd., Rusty Springs Prospect, N.T.S. Map Sheet 116K/8 and 9, Porcupine Ranges, Yukon Territory. Unpublished report for Rio Alto Exploration Ltd. 16 p.
- Beck, F.M., 1978. Rusty Springs Prospect, Yukon Territory, 1978 Exploration Summary. Unpublished report for Rio Alto Exploration Ltd. 16 p.
- Chamberlain, J.A. 1986. Drill logs for holes 86-1 and 86-2, Rusty Springs Property. Unpublished report for Kenton Natural Resources Corp., 5 p.
- Chernoff, M.N., 1976. Geology of the Rusty Springs Mineral Prospect, Porcupine Ranges, Yukon Territory. Unpublished report for Rio Alto Exploration Ltd. 14 p.
- Davis, J.W., and Aussant, C.H., 1982. Report on Geochemical, Geophysical, Geological, and Trenching programs on the Rusty Springs Property, Yukon Territory. Unpublished report for Kenton Natural Resources Corporation, 27p.
- Dawson, K.M., Panteleyev, A., Sutherland-Brown, A., and Woodsworth, G.J., 1991. Regional Metallogeny, Chapter 19 of Geology of the Cordilleran Orogen in Canada, H. Gabrielse and C.J. Yorath (ed.); Geological Survey of Canada, Geology of Canada, no.4, p.707-768.
- Downie, C.C., 1994. Geological Report on the Rusty Springs Property, Yukon Territory. Unpublished report for Eagle Plains Resources Ltd., 24 p.
- Downie, C.C. and Greig, C.J. 2000. Diamond Drilling Report on the Rusty Springs Property. Unpublished report for Eagle Plains Resources Ltd., 24 p.
- Hansen, D., 1979. A Geological Model of the Rusty Springs Prospect, Porcupine Range, Yukon Territory. Unpublished B.Sc. thesis, University of Western Ontario, London, 32 p.
- Hansen, D., and Bankowski, J., 1979. Report of Geological Program, Rusty Springs Prospect, Porcupine Ranges, Yukon Territory. Unpublished report for Rio Alto Exploration Ltd., 21 p.

- Hodder, R.W., 1997. Rusty Springs Prospect, Yukon Territory, Observations and Interpretations. Unpublished report for Eagle Plains Resources Ltd., 27 p.
- Kirker, J., 1980a. Petrology and Ground Preparation of Rusty Springs Base Metal Deposit, Yukon Territory: a Thesis Proposal. Unpublished report for Rio Alto Exploration Ltd., 8 p.
- Kirker, J., 1980b. Summary of '80 Field Work , Rusty Springs, Yukon Territory. Unpublished report for Rio Alto Exploration Ltd., 3 p.
- Kirker, J.K., 1982. Geology, Geochemistry and Origin of Rusty Springs Lead-Zinc-Silver Deposit, Yukon Territory. Unpublished M.Sc. thesis, University of Calgary, Calgary, 159 p.
- Lane, L.S., 1998. Latest Cretaceous-Tertiary Tectonic Evolution of Northern Yukon and Adjacent Arctic Alaska. American Association of Petroleum Geologists Bulletin, v.82, no.7, p. 1353-1371.
- Leach, D.L., and Sangster, D.F., 1993. Mississippi Valley-type lead-zinc deposits. In Kirkham, R.V., Sinclair, W.D., Thorpe, R.I., and Duke, J.M., eds., Mineral Deposit Modeling: Geological Association of Canada, Special Paper 40, p. 289-314.
- Liedtke, G.J., 1980. Report on Exploration Results, 1980, Rusty Springs Prospect, Yukon Territory. Unpublished report for E & B Explorations Inc., 31 p.
- Norris, D.K., 1981. Geology, Porcupine River, Yukon Territory. Geological Survey of Canada, Map 1522A, scale 1:250,000.
- Norris, D.K. (Editor), 1996. The Geology, Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie. Geological Survey of Canada, Bulletin 422, 401 p.
- O'Donnell, J.R. 1974. Geological, Geochemical and Geophysical Report on the Mink Claim Group. Unpublished report for Inexco Mining Corporation, 53 p.
- Power, M.A., 1998. Gravity and Reflection Seismic surveys on the Rusty Springs Property, Northern Yukon Territory. Unpublished report for Eagle Plains Resources Ltd., 23 p.
- Richards, B.C., Bamber, E.W., and Utting, J., 1996. In The Geology, Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie. Geological Survey of Canada, Bulletin 422, p. 201-251.
- Schoel, G., 1978. Geology and Genesis of the Rusty Springs Zn-Pb-Cu-Ag Prospect, Porcupine Range, Yukon Territory. Unpublished B.Sc. thesis, University of Western Ontario, London.
- Tempelman-Kluit, D.J., 1981. Termuende (Rusty Springs), in: Yukon Geology and Exploration, 1979-80. Geology Section, Department of Indian and Northern Affairs, Whitehorse, pp. 301-304.

Termuende, T.J., 1996. Diamond Drilling Report on the Rusty Springs Property, Yukon Territory. Unpublished report for Eagle Plains Resources Ltd., 24 p.

Termuende, T.J., and Downie, C.C., 1997. Diamond Drilling Report on the Rusty Springs Property, Yukon Territory. Unpublished report for Eagle Plains Resources Ltd., 23 p.

Termuende, T.J., and Downie, C.C., 1998. Reverse-circulation Drilling Report on the Rusty Springs Property, Yukon Territory. Unpublished report for Eagle Plains Resources Ltd., 19 p.

Titley, S.R., 1993. Characteristics of High-temperature, Carbonate-hosted Massive Sulphide Ores in the United States, Mexico, and Peru. In Kirkham, R.V., Sinclair, W.D., Thorpe, R.I., and Duke, J.M., eds., Mineral Deposit Modeling: Geological Association of Canada, Special Paper 40, p. 585-516.

White, P.S., 1978. Report of 1977 Exploration of the Rusty Springs Mineral Prospects, Porcupine Ranges, Yukon Territory. Unpublished report for Rio Alto Exploration Ltd., 10 p.

White, P.S., 1979. Report of 1979 Exploration of the Rusty Springs Mineral Prospect, Porcupine Ranges, Yukon Territory. Unpublished report for Rio Alto Exploration Ltd., 11 p.

Woodsworth, G.J., Anderson, R.G., and Armstrong, R.L., 1991. Plutonic regimes, Chapter 15 in Geology of the Cordilleran Orogen in Canada, H. Gabrielse and C.J. Yorath (ed.); Geological Survey of Canada, Geology of Canada, no.4, p.491-531.

APPENDIX I

Certificate of Qualification

## CERTIFICATE OF QUALIFICATION

I, Charles C. Downie of 122 13<sup>th</sup> Ave. S. in the city of Cranbrook in the Province of British Columbia hereby certify that:

- 1) I am a Professional Geoscientist registered with the Association of Professional Engineers and Geoscientists of British Columbia (#20137).
- 2) I am a graduate of the University of Alberta (1988) with a B.Sc. degree and have practiced my profession as a geologist continuously since graduation.
- 3) This report is supported by data collected during fieldwork as well as information gathered through research.
- 4) I hold 125,000 shares of Eagle Plains Resources; I Hold an option to purchase a further 100,000 Common Shares of Eagle Plains at \$0.25 per share.

Dated this 30th day of January, 2002 in Cranbrook, British Columbia.

A circular stamp containing a handwritten signature in cursive script, which appears to be 'CD'. The stamp is slightly faded and has a dotted border.

Charles C. Downie, P. Geo.

**APPENDIX II**

**Statement of Expenditures**

## STATEMENT OF EXPENDITURES : 2001 RUSTY SPRINGS EXPLORATION PROGRAM

The following expenses were incurred on the Rusty Springs property, Dawson Mining Division, for the purpose of mineral exploration between the dates of June 1st to September 30<sup>th</sup>, 2001.

### Personnel:

Tim J. Termuende, P.Geo.: 5.0 days x \$450.00/day.....	\$2,250.00
Charles C. Downie, P.Geo.: 5.0 days x \$300.00/day.....	\$2,250.00
K. Scholtz, geological assistant: 1.5 days x \$250.00/day .....	\$375.00

### Equipment Rental:

4WD Vehicle; 2.0 days x \$50.00/day.....	\$100.00
Mileage: 1154 km x \$.20/km.....	\$1,938.20
Hand-held Radios (2): 2.0 days x \$20.00/day.....	\$40.00
Field Supply : 6 man-days @ \$30.00/day.....	\$180.00

### Other:

Meals/Accom:.....	\$1,512.68
Groceries:.....	\$63.61
Materials: .....	\$4.28
Airfare:.....	\$953.43
Air Charter(Fireweed Helicopters): .....	\$3,897.21
Shipping:.....	\$63.25
Fuel:.....	\$206.62
Filing Fees:.....	\$285.00
Drafting/Repro:.....	\$4.28
Misc. GST: .....	\$272.62
Repairs: .....	\$13.10
Analytical (Northern Analytical – 30 element ICP):.....	\$313.08
Miscellaneous:.....	\$54.48
Report (estimate): .....	\$2,500.00
Handling Fees (Toklat Resources):.....	<u>\$744.92</u>

**TOTAL:       \$16,500.08**

**APPENDIX III**  
**Analytical Results**

6-Jul-01

ECO-TECH LABORATORIES LTD.  
10041 Dallas Drive  
KAMLOOPS, B.C.  
V2C 6T4

ICP CERTIFICATE OF ANALYSIS AK2001-146

TOKLAT RESOURCES INC.  
2720-17th STREET SOUTH  
CRANBROOK, B.C.  
V1C 6Y6

Phone: 250-573-5700  
Fax : 250-573-4557

ATTENTION: TIM TERMUENDE

No. of samples received: 11  
Sample Type: Rock  
Project #: RS01  
Shipment #: RS01-01  
Samples submitted by: T. Termuende

Values in ppm unless otherwise reported

Et #.	Tag #	Ag	Al %	As	Ba	Bi	Ca %	Cd	Co	Cr	Cu	Fe %	La	Mg %	Mn	Mo	Na %	Ni	P	Pb	Sb	Sn	Sr	Ti %	U	V	W	Y	Zn
1	TTRS01R01	>30	0.03	>10000	215	<5	0.11	35	2	274	4517	1.15	20	0.04	48	8	<0.01	10	<10	>10000	615	<20	<1	<0.01	<10	3	<10	4	108
2	TTRS01R02	>30	<0.01	95	<5	<5	0.01	12	<1	78	96	0.58	<10	<0.01	13	<1	<0.01	5	20	>10000	1655	<20	<1	<0.01	<10	<1	<10	<1	<1
3	TTRS01R03	>30	0.02	>10000	115	<5	0.13	53	4	205	>10000	0.37	<10	0.02	87	9	<0.01	13	<10	>10000	>10000	<20	<1	0.01	30	2	<10	3	244
4	TTRS01R04	<0.2	2.63	85	315	<5	1.26	2	8	422	86	>10	20	0.25	73	<1	0.02	65	7140	310	<5	<20	273	<0.01	<10	451	<10	<1	454
5	TTRS01R05	>30	0.18	>10000	65	*	0.09	130	16	14	>10000	0.83	20	0.03	474	6	<0.01	28	*	6600	*	*	<1	<0.01	*	*	*	83	4711
6	TTRS01R06	0.6	0.03	65	1490	<5	0.02	<1	3	170	498	0.44	<10	<0.01	55	8	<0.01	26	40	132	20	<20	13	<0.01	<10	13	<10	<1	50
7	TTRS01R07	1.2	0.10	35	130	<5	0.73	<1	8	259	59	1.41	<10	0.35	66	6	<0.01	28	220	26	10	<20	2	<0.01	<10	7	<10	<1	159
8	TTRS01R08	>30	0.03	>10000	135	<5	0.39	589	10	56	>10000	2.10	<10	0.02	180	<1	<0.01	6	>10000	>10000	>10000	<20	15	<0.01	<10	<1	170	<1	3336
9	TTRS01R09	9.2	0.03	35	160	<5	>10	1	16	9	72	1.64	<10	>10	383	<1	0.02	59	130	32	<5	220	95	<0.01	<10	10	<10	24	140
10	TTRS01R10	0.8	0.10	30	65	<5	>10	6	8	21	88	0.53	<10	1.29	224	<1	0.01	1338	370	102	<5	<20	24	<0.01	<10	12	20	9	>10000
11	TTRS01R11	7.4	0.17	25	65	<5	1.20	4	7	270	68	1.67	<10	0.47	304	12	0.01	51	480	698	20	<20	27	<0.01	<10	27	<10	10	820

NOTE: \*= unable to report due to massive copper interference.

QC/DATA:

Resplit:

1	TTRS01R01	>30	0.03	>10000	220	<5	0.12	37	2	328	4432	1.29	20	0.03	53	9	<0.01	12	40	>10000	670	<20	<1	<0.01	<10	3	<10	3	82
---	-----------	-----	------	--------	-----	----	------	----	---	-----	------	------	----	------	----	---	-------	----	----	--------	-----	-----	----	-------	-----	---	-----	---	----

Repeat:

1	TTRS01R01	>30	0.03	>10000	220	<5	0.11	27	2	266	4763	1.12	20	0.04	46	8	<0.01	10	<10	>10000	605	<20	1	<0.01	<10	3	<10	4	82
---	-----------	-----	------	--------	-----	----	------	----	---	-----	------	------	----	------	----	---	-------	----	-----	--------	-----	-----	---	-------	-----	---	-----	---	----

Standard:

GEO'01		1.4	1.72	55	140	<5	1.56	<1	19	55	91	3.55	<10	0.94	677	<1	0.02	25	670	22	5	<20	62	0.10	<10	73	<10	4	72
--------	--	-----	------	----	-----	----	------	----	----	----	----	------	-----	------	-----	----	------	----	-----	----	---	-----	----	------	-----	----	-----	---	----

FP/kk  
df/150  
XLS/01Toklat  
Fax: 250-426-6899 & E-mail

  
**ECO-TECH LABORATORIES LTD.**  
 Frank J. Pezzotti, A.Sc.T.  
 B.C. Certified Assayer



**ASSAYING  
GEOCHEMISTRY  
ANALYTICAL CHEMISTRY  
ENVIRONMENTAL TESTING**

10041 Dallas Drive, Kamloops, B.C. V2C 6T4  
Phone (250) 573-5700 Fax (250) 573-4557  
email: ecotech@direct.ca

**CERTIFICATE OF ASSAY AK 2001-146**

**TOKLAT RESOURCES INC.**  
2720-17th STREET SOUTH  
**CRANBROOK, B.C.**  
V1C 6Y6

6-Jul-01

**ATTENTION: TIM TERMUENDE**

*No. of samples received: 11*

*Sample Type: Rock*

*Project #: RS01*

*Shipment #: RS01-01*

*Samples submitted by: T. Termuende*

ET #.	Tag #	Ag (g/t)	Ag (oz/t)	Cu (%)	Pb (%)	Sb (%)
1	TTRS01R01	96.8	2.82	-	3.22	-
2	TTRS01R02	2150.0	62.70	-	57.8	-
3	TTRS01R03	62.3	1.82	3.07	13.2	1.44
4	TTRS01R04	-	-	-	-	-
5	TTRS01R05	88.9	2.59	40.7	-	-
6	TTRS01R06	-	-	-	-	-
7	TTRS01R07	-	-	-	-	-
8	TTRS01R08	8460.0	246.72	5.69	4.56	9.77
9	TTRS01R09	-	-	-	-	-
10	TTRS01R10	-	-	-	-	-
11	TTRS01R11	-	-	-	-	-

XLS/01Toklat

Fax: 250-426-6899 & E-mail

  
**ECO-TECH LABORATORIES LTD.**  
Frank J. Pezzotti, A.Sc.T.  
B.C. Certified Assayer



ASSAYING  
GEOCHEMISTRY  
ANALYTICAL CHEMISTRY  
ENVIRONMENTAL TESTING

10041 Dallas Drive, Kamloops, B.C. V2C 6T4  
Phone (250) 573-5700 Fax (250) 573-4557  
email: ecotech@direct.ca

## CERTIFICATE OF ANALYSIS AK 2001-146

TOKLAT RESOURCES INC.  
2720-17th STREET SOUTH  
CRANBROOK, B.C.  
V1C 6Y6

10-Jul-01

ATTENTION: TIM TERMUENDE

No. of samples received: 11

Sample Type: Rock

Project #: RS01

Shipment #: RS01-01

Samples submitted by: T. Termuende

ET #.	Tag #	Ge (ppm)
1	TTRS01R01	<20
2	TTRS01R02	<20
3	TTRS01R03	<20
4	TTRS01R04	<20
5	TTRS01R05	<20
6	TTRS01R06	<20
7	TTRS01R07	<20
8	TTRS01R08	<20
9	TTRS01R09	<20
10	TTRS01R10	<20
11	TTRS01R11	<20

**QC/DATA:**

Repeat:

1 TTRS01R01 <20



ECO-TECH LABORATORIES LTD.

Frank J. Pezzotti, A.Sc.T.  
B.C. Certified Assayer

XLS/01Toklat

Fax: 250-426-6899 & E-mail

APPENDIX IV

2001 Rock Sample Descriptions

TTRS01R01:

Massive, coarse galena with tetrahedrite in Carbonate breccia. Galena as coarse-grained, euhedral crystals.

TTRS01R02:

Drusy-textured, siliceous katshat material with prominent scoroditic staining, minor smithsonite

TTRS01R03:

Malachite staining on quartz-flooded katshat material. Oxides coating much of surface (possibly smithsonite?)

TTRS01R04:

Oolitic iron material.

TTRS01R05:

Botryoidal malachite with some azurite stain.

TTRS01R06:

Barite crystals on brown-weathering, limonitic breccia.

TTRS01R07:

Rounded, transported Katshat material.

TTRS01R08:

High-grade silver, copper, arsenic material-Trench 8

TTRS01R09:

Dense, weakly limonitic dolomite. No visible mineralization.

TTRS01R10:

Smithsonite/aurichalcite-headed breccia with dolomite clasts.

TTRS01R11:

Sphalerite-mineralized shales from DDH 99-01, 256.4m




FIGURE #3 LOCATION

**Early Cretaceous**

- 9 shale and siltstone
- 8 quartz arenite

**Jurassic and Early Cretaceous**

- 7 dark weathering siliclastic rocks: shale, silty mudstone, and sandstone

**Late Carboniferous and Permian(?)**

- 6 Ettrian and Jungle Creek formations; limestone and fine grained calcareous and siliceous clastic rocks


**Devono-Mississippian fine-grained siliclastic rocks**

- 5 Unnamed shale; siliceous mudstone, slate, shale, siltstone; rare argillaceous limestone

**Early and Middle Devonian**

- 4 Ogilvie Formation; dolostone: fetid, vuggy, commonly brecciated; subordinate limestone and argillaceous rocks

— Claim Boundary



**Early Proterozoic to Early Paleozoic**

- 3 undivided dolomite and siliclastic rocks

**Late Cambrian to Early Devonian**

- 2 dolomite, local quartz-rich sandstone (quartzite) and siliceous siltstone (siltite)


**Early to Late Proterozoic**

- 1 rusty weathering siliclastic rocks: sandstone (quartzite), siltstone, and silty mudstone (siltite)



EAGLE PLAINS RESOURCES	
RUSTY SPRINGS PROJECT	
CLAIM LOCATION, REGIONAL GEOLOGY AND 2001 SAMPLE LOCATIONS	
GEOLOGY AFTER GREIG, 1999	
FIG.2	scale as shown
JANUARY 2002	
TOKLAT RESOURCES INC.	

\* TTRS01R02 2001 SAMPLE LOCATION



**Devono-Mississippian fine-grained siliciclastic rocks**  
 Unnamed shale



- 5b** shale, siltstone, siliceous mudstone, slate; rare argillaceous limestone
- 5a** brecciated black chert and siliceous mudstone, minor shale

**Early and Middle Devonian**

Ogilvie Formation

- 4e** Katshat unit: leached, mineralized and altered dolostone: porous limonitic to kaolinitic material containing quartz veinlets and laminae which locally contain sulphides and sulphosalts
- 4d** undifferentiated dolostone: fetid, vuggy, commonly brecciated; subordinate limestone and argillaceous rocks
- 4c** dolostone, brecciated, commonly siliceous
- 4b** dolostone, not brecciated
- 4a** limestone; locally fossiliferous

- road or cat trail
- mineral showing
- bulldozer trench
- vertical drillhole
- inclined drillhole
- inclined drillhole (>1)
- 1999 drillhole



**094242**

EAGLE PLAINS RESOURCES

**RUSTY SPRINGS PROJECT**


GEOLOGY OF THE RUSTY SPRINGS PROPERTY;  
1999 DRILLHOLE LOCATIONS

PROPERTY GEOLOGY AFTER C. GREIG 1999

FIG. 3	DATE: January 2000	1: 15,000
--------	-----------------------	-----------

TOKLAT RESOURCES INC.

99toklat\rusty\mike\_orma\_geol-i4.dwg



DDH96-4  
1.1 gm Ag, 881 ppm Cu,  
139 ppm Pb, 3301 ppm Zn  
over 19.1 m

DDH96-3  
0.22% Zn  
over 19.8 m

DDH97-5  
lost hole at 50.3 m  
in Katshat unit;  
7.6 m at EOH  
1811 ppm Zn, 21 ppm Ag,  
926 ppm Cu, 528 ppm Pb

DDH97-3  
Katshat unit: 38.1 m  
0.06 gm Ag, 475 ppm Cu,  
1137 ppm Zn, 475 ppm Cu

DDH97-6  
lost hole at 53 m  
in brecciated  
chert/mudstone

DDH97-7  
(projected 500 m to section)  
51.9 m with 2.66 gm Ag,  
1767 ppm Zn, 216 ppm Cu,  
163 ppm Pb

DDH99-1  
lost hole at 323 m;  
silicified and brecciated  
in bottom 80-95 m;  
locally high Zn: 3000 ppm  
over 7 and 15 m

DDH99-3  
lost hole at  
165 m in  
brecciated  
siliceous  
mudstone

DDH99-2  
lost hole at  
130 m in  
brittle fault

0 250 500 750 1000 m  
no vertical exaggeration

**Devono-Mississippian fine-grained siliciclastic rocks**  
Unnamed shale

- 5b** shale, siltstone, siliceous mudstone, slate; rare argillaceous limestone
- 5a** brecciated black chert and siliceous mudstone, minor shale

**Early and Middle Devonian**

- Ogilvie Formation
- 4e** Katshat unit: leached, mineralized and altered dolostone: porous limonitic to kaolinitic material containing quartz veinlets and laminae that locally contain sulphides and sulphosalts
  - 4d** undifferentiated dolostone: fetid, vuggy, commonly brecciated; subordinate limestone and argillaceous rocks

possible W-side-up fault approx.  
coincident with fold axial plane

094242

EAGLE PLAINS RESOURCES  
**RUSTY SPRINGS PROJECT**

---

GEOLOGIC CROSS SECTION MIKE AND ORMA HILLS  
C-C' LOCATION FIG. 3

---

PROPERTY GEOLOGY AFTER C. GREIG 1999

---

FIG. 4	DATE: January 2000	as shown
--------	-----------------------	----------

---

TOKLAT RESOURCES INC.

99tolat\rusty\mike\_orma\_sectn-14.dwg