ASSESSMENT REPORTS

Dawson M.D.

MAP No. 116B/3 1150/14 TYPE OF WORK: Geophys, Geological, Geochem

REPORT FILED UNDER	UKON Joint Venture 090274
DATE PERFORMED	May 30-June5, Sept.13-23 DATE FILED: 18 Jan. 1978 ~~Nov. 2-5, 1977~~
LOCATION – LAT.	64 01'N Hunker Creek Area, Yukon.
LONG.	139 04'W
CLAIM Nos.	Surprize 1-16, YA 9565-YA9580; Surprize 17-32, YA 10204-YA 10219; Surprize 33-95, YA 10681-YA10743
$6664.52.	
WORK DONE BY	Alan R. Archer, P. Eng.
WORK DONE FOR	
REMARKS	Water and soil geochemical surveys have outlined three areas anomalous in uranium. Source of uranium not yet discovered.

090274

STAR PRINTING, WHITEHORSE

JAN 13 1978
MINING RECORDER
LAND AGENT
DAWSON, Y.T.
I.A. & N.D.

Assessment Report

on

Surprize 1-95 Claims

Office Of The Supervising Mining Recorder
JAN 26 1977
WHITEHORSE
Yukon Territory

Dawson Mining District

Claim Sheets 116B/3 and 1150/14

Latitude 64°01'N, Longitude 139°04'W

December 23, 1977

RECEIVED
JAN 27 1978
INSP.
WHITEHORSE Y.T.

Alan R. Archer, P. Eng. Consulting Engineer

040274

This report has been examined by the Geological Evaluation Unit and is recommended to the Commissioner to be considered as representing assessment work to the amount of $6664.52

Recorder

Considered as prospecting/exploration work under Section 53 of the Quartz Mining Act.

B. R. BAXTER
Supervising Mining Recorder
for Commissioner of Yukon Territory

TABLE OF CONTENTS

Text	Page
Introduction	1
Property, Location and Access	1
Geology and Mineralization	2
1977 Exploration Program	3
Geology, Geochemical and Radiometric Surveys	3
Trenching	4
Conclusion and Recommendation	6

Figures in Text		Following Page
Figure U-UN3	Regional Water Sampling - Scale 1:50,000	4
Figure U-UN7	Trenching Soil Anomaly B - Scale 1:600	5
Figure U-UN8	Trenching Soil Anomaly A - Scale 1:300	5

Figures in Pocket

Figure U-UN4	Geology,Reconnaissance Geochemistry and Radiometrics - Scale 1:10,000
Figure U-UN5	Geochemistry and Geology - Scale 1:5,000
Figure U-UN6	Radiometrics and Geology - Scale 1:5,000

INTRODUCTION

The Unexpected property was staked as the Surprize 1-16 claims in 1976 to cover an area where highly anomalous uranium values were obtained from water in streams draining a schist-intrusive contact. Subsequent soil sampling located two specific areas of weakly anomalous uranium response near the creek headwaters.

The 1977 program was done in three stages under management by Archer, Cathro & Associates Ltd.; (a) May 30 to June 5 - claim staking, grid soil sampling and radiometric surveys, geological mapping and hand pitting; (b) September 13 to 23 - additional claim staking, reconnaissance soil sampling and geological mapping, and bulldozer trenching; and, (c) November 2 to 5 - additional bulldozer trenching. Field work was supervised by geologists M.P. Phillips and U. Schmidt.

Geochemical analyses for uranium was done at Chemex Labs Ltd., North Vancouver, B.C., by hot acid extraction of a minus 80 mesh fraction followed by fusion in a sodium fluoride-based flux and examination with a G.K. Turner fluorometer.

PROPERTY, LOCATION AND ACCESS

The Unexpected property consists of 95 contiguous Surprize claims that form a sub-rectangular block. The claims are recorded at Dawson as follows:

CLAIM NAME	GRANT NUMBERS	EXPIRY DATE
Surprize 1,3,5,7	YA 9565 - YA 9568	14 April, 1982
Surprize 10,12,14,16	YA 9569 - YA 9572	14 April, 1982
Surprize 2,4,6,8	YA 9573 - YA 9576	14 April, 1982
Surprize 9,11,13,15	YA 9577 - YA 9580	14 April, 1982
Surprize 17-32	YA10204 - YA10219	1 June, 1978
Surprize 33-95	YA10681 - YA10743	20 Sept., 1978

The property is located at 64°01'N and 139°04'W straddling claim sheets 1150/14 and 116B/3, 21 km (13 miles) by road east of Dawson. Access is via the all-weather Hunker Creek road which crosses the south end of the claim block.

GEOLOGY AND MINERALIZATION

The topography is typical of unglaciated terrain throughout the Dawson Range. Rounded subdued hills rise to elevations of 1200 m with local relief of up to 500 m. Streams occupy V-shaped valleys which have been modified by late Tertiary rejuvenation. Outcrop is rare and most hillsides are modified by a thin cover of residual till, soil and humus. Vegetation is characterized by open pine and aspen on south facing slopes, and thick moss with black spruce on north facing slopes where permafrost extends to surface. Surface leaching in similar terrain at the Casino property and the Keno Hill district reaches depths of up to 150 m.

The claims cover the western contact of a quartz-feldspar porphyry stock that intrudes a complex metamorphic assemblage called the Schist-Gneiss Unit by D.J. Tempelman-Kluit of the G.S.C. The stock consists of rounded, sometimes smoky, quartz eyes up to 4 mm wide, and subhedral feldspar phenocrysts up to 1 mm long in a white to buff felsic groundmass. It is thought to be contemporaneous with stocks elsewhere in the district that have been dated as Eocene (about 50 m y).

The Schist-Gneiss Unit is an undivided mixture of Klondike Schist and Nasina Quartzite. Most outcrops on the Surprize claims consist of dark to pale green or grey chlorite schist with some black carbonaceous phyllites, minor marble and foliated greenstone. These metasedimentary rocks are thought to correlate with less deformed Paleozoic rocks northeast of the Tintina fault in the Finlayson Lake district. The age of the metamorphism is believed to be Early or Middle Triassic.

Geochemical exploration in 1976 located two anomalies. Anomaly A consisted of a 400 m by 500 m area near the headwaters of Surprize Creek with soil values between 8 and 184 ppm U while anomaly B consisted of a small area of scattered anomalous soil values up to 184 ppm U near the headwaters of Tinhorn Gulch. No other evidence of uranium mineralization was found and rock assays of both intrusive and schist returned uranium values of 5 ppm or less.

1977 EXPLORATION PROGRAM

Geology, Geochemical and Radiometric Surveys

The 1977 mapping is illustrated on Figure U-UN4 in the pocket. Outcrop is virtually absent and the contact between the quartz porphyry stock and the Schist-Gneiss Unit has been defined by mapping surface float and rock fragments found in soil sample pits. The claims cover the western contact of the stock, which appears to be quite irregular although this may be merely an apparent irregularity caused by the distribution of float. A narrow linear band of Schist-Gneiss at the headwaters of Tinhorn Gulch is either a roof pendant within the stock or an unusually sharp contact irregularity, perhaps indicating an offset due to faulting.

Figure U-UN4 also illustrates results of reconnaissance geochemical sampling along drainage and claim lines of the Surprize 33-95 claims while Figure U-UN5 illustrates combined results of 1976 and 1977 grid soil sampling on the Surprize 1-32 claims. Figure U-UN3 on the following page shows assays of water samples collected within and around the property in 1976 and 1977. The highest 1977 value (95 ppb) was obtained from Surprize Creek, which drains the swampy bench containing 1976 soil anomaly A. The 1977 soil sampling program located a new area of weak response up to 59 ppm U (called anomaly C), near the headwaters of Catchup Gulch. All anomalies are found near the intrusive-porphyry contact.

Figure U-UN6 illustrates grid radiometric surveys in 1976 and 1977 on the Surprize 1-32 claims while Figure U-UN4 illustrates reconnaissance radiometric survey response along the claim lines of the Surprize 33 to 95 claims. All surveys were conducted with a Scintrex BGS-1SL broadband scintillometer (43 cc crystal). Backgrounds range from 40 to 150 cps over the porphyry and from 30 to 80 cps over the Schist-Gneiss Unit. No strongly anomalous areas were found. A response of twice background was obtained over soil anomalies B and C while soil anomaly A gave a slightly lower response of up to 50 per cent above background. Scintillometer readings of twice background with no associated geochemical response were also obtained from porphyry outcrops along the Hunker Creek road near Baseline A. Radiometric values along the claim lines tend to be slightly above background where the lines cross the porphyry contact.

Trenching

Initial trenching consisted of a hand pit followed by three trenches dug

with a D7 bulldozer, without ripper, at soil anomaly B. Attempts to trench soil anomaly A by hand and with the small bulldozer were unsuccessful because of frozen overburden. A ripper-equipped D8 bulldozer was obtained in early November and three trenches were cut at anomaly A without difficulty.

Figure U-UN7 (on the following page) illustrates the results of trenching at soil anomaly B. The initial hand pit was dug to a depth of 1.5 m and geochemical profiling returned an assay of 55 ppm U from decomposed chlorite schist bedrock at the bottom compared with only 1.5 ppm U from the overlying B soil horizon. Trench UN-B1, dug to a depth of 3 m at the same location with a bulldozer, returned bedrock assays that ranged from 35 ppm to 80 ppm U and averaged 58 ppm U and, as in the hand pit, demonstrated sharp increase in U content with depth. Bedrock appears to be unleached and consists of partially frost-shattered, grey chlorite schist containing minor vuggy, milky white quartz lenses. A second trench, located 30 m north, exposed similar chlorite schists and a sample profile returned assays ranging from 0.5 ppm U at surface to 28 ppm U at the base. Trench UN-B3, situated 60 m north of Trench UN-B1, exposed fresh quartz feldspar porphyry. A sample profile from this trench returned assays ranging from 0.5 ppm U at the top to 5.5 ppm U at the base. None of the trenches show any evidence of sulfides, sulfide leaching or secondary uranium oxides.

Figure U-UN8 (following Figure U-UN7 in the text) illustrates results of trenching at soil anomaly A. Two trenches were cut near the projected location of the quartz feldspar porphyry contact and one about 150 m northwest. All three trenches reached bedrock which consists of weathered, frost-broken chlorite schist with limonite-stained quartz lenses. Overburden ranges up to 2.3 m deep and




FIG. U-UNT


ARCHER, CATHRO AND ASSOCIATES

TRENCHING - SOIL ANOMALY B


SURPRIZE 1-95 CLAIMS

UKON JOINT VENTURE

Dec 23/77



TRENCH UN-A1



PROFILE A

cps	U ppm
130	10
140	6
150	4
150	2
160	3


0 – OVERBURDEN - BROWN CLAY, CHLORITE SCHIST QUARTZ - FELDSPAR PORPHYRY
1m – CHLORITE SCHIST - HIGHLY WEATHERED
2m

PROFILE B

cps	U ppm
130	57
140	4
160	6
180	14
200	38

0 – WEBBURDEN - BROWN CLAY, CHLORITE SCHIST, ABUNDANT QUARTZ FELDSPAR PORPHYRY NEAR BASE
1m
2m

TRENCH UN-A2



PROFILE A

cps	ppm U
170	34
200	11
210	8
220	7
220	8
230	8
250	8


0 – OVERBURDEN - BROWN CLAY, CHLORITE SCHIST, ABUNDANT QUARTZ FELDSPAR PORPHYRY FLOAT NEAR BASE
1m
2m – CHLORITE SCHIST - HIGHLY WEATHERED, LIMONITIC QUARTZ IS COMMON
3m

PROFILE B

cps	ppm U
150	
170	
190	
210	
230	
240	
240	
260	

0 – OVERBURDEN - BROWN SOIL, CHLORITE / SCHIST, OCCASIONAL PORPHYRY BOULDERS
1m
2m
3m – CHLORITE SCHIST - WEATHERED WITH MODERATE LIMONITE QUARTZ

TRENCH UN-A3



SCALE 1:300

PROFILE A


cps	ppm U
150	10
180	9
170	3
200	4
200	5
200	6

0
1m – OVERBURDEN - BROWN SOIL; CHLORITE SCHIST, PORPHYRY BOULDERS NEAR BOTTOM
2m – CHLORITE SCHIST - HIGHLY WEATHERED

PROFILE B

cps	ppm U
120	36
120	10
140	22
150	3
160	1
180	3
200	6
200	7

0
1m
2m – OVERBURDEN - BROWN SOIL; CHLORITE SCHIST AND PORPHYRY FRAGMENTS
3m – CHLORITE SCHIST - HIGHLY WEATHERED



1:5,000

Association of Professional Engineers of the Yukon
A. R. ARCHER
GEOLOGICAL ENGINEER

Dec 23/77

FIG. U-UN8

ARCHER, CATHRO AND ASSOCIATES LTD

TRENCHING - SOIL ANOMALY A

SURPRIZE 1-95 CLAIMS

UKON JOINT VENTURE

consists of partially frozen dark brown soil with chlorite schist and quartz-feldspar porphyry fragments. The porphyry float is most abundant near the bottom, often occurring as boulders up to 15 cm across. Geochemical profiling consistently returned higher uranium values (10 to 70 ppm U) from the near surface organic rich soil than from the base of the overburden (3 to 12 ppm U) or from bedrock (2 to 17 ppm U). A sample of water seeping from the bedrock-overburden contact in Trench UN-A2 assayed 710 ppb U while a surface seepage immediately east assayed 135 ppb U. A second water sample from the same surface seepage about 100 m south, just east of Trench UN-A1, assayed 245 ppb U.

CONCLUSIONS AND RECOMMENDATIONS

Unusually anomalous levels of uranium are found in water from some of the streams that drain a contact between quartz-feldspar porphyry and chlorite schist. Soil sampling located two anomalies, called A and B, which were explored by bulldozer trenching. Anomaly A proved to be caused by uranium concentrating in soil in an area where ground water seeps return intensely anomalous uranium values ranging from 135 to 710 ppb U. Anomaly B proved to be a tongue or linear roof pendant of unmineralized chlorite schist with a uranium content ranging between 35 and 80 ppm U. It has not been determined whether the uranium in the schist is a background component or has been precipitated from ground water.


The source of the uranium is uncertain. Rock samples of porphyry return normal background values (2 to 5 ppm U) and not all streams draining the

porphyry have anomalous water values. The implication is that only specific zones within the porphyry are liberating uranium into the geochemical cycle. These zones could be related to compositional differences, structures or unexposed uranium occurrences that are being actively leached. If such occurrences exist, they will be difficult to locate as leaching in similar unglaciated terrane elsewhere in Yukon reaches up to 150 m in depth.

Further work is warranted and should consist of a 300 m diamond drill hole across the tongue of chlorite schist at soil anomaly B. The hole should be angled in order to cut through the porphyry contact at a depth of 150 m or more below surface. Several days of bulldozer trenching should be concurrently conducted at soil anomaly A in attempt to expose the schist-porphyry contact.

Respectfully submitted,

ARCHER, CATHRO & ASSOCIATES LTD.,



ARA:jm

A.R. Archer, B.A.Sc., P. Eng.

SURPRIZE 33-95

SURPRIZE 1-32
(see Figs. U-UN5 and U-UN6)

BASELINE A
BASELINE B
BASELINE C
BASELINE D
BASELINE E
BASELINE F

Psn
eTqfp

Australian Hill
Gulch
Barcelau
Creek
Alki
Creek
Preido
Gulch
Unexpected
Gulch
Catchup
Gulch
Tinhorn
Gulch
Hunker
Creek
Surprize
Creek

powerline
powerline
powerline

tuff breccia
marble
marble

TO DAWSON
Hunker Road
Hunker Road

CAMP


SOIL ANOMALY A
SOIL ANOMALY B
SOIL ANOMALY C

UN-A1
UN-A2
UN-A3
UN-B1
UN-B2
UN-B3

1400
1600
1800
2000
2200
2400
2600

64° 00
139° 04

LOCATION MAP



LEGEND

eTqfp quartz feldspar porphyry, rounded smoky quartz eyes

Psn Schist-Gneiss Unit — dark green to grey chlorite schist, minor carbonaceous phyllite, marble and foliated greenstone

approximate geological contact

claim post location and claim numbers

bulldozer trench

(110) • 2·5 — (110)-total count at waist height using a Scintrex BGS-1SL broadband scintillometer (43 cc crystal)
2·5-soil sample value in ppm uranium
T-trace

42 — water sample value in ppb uranium

area of soil values in excess of 8 ppm uranium

• 7·5 — rock sample value in ppm uranium

Association of Professional Engineers of Ont. A. R. ARCHER GEOLOGICAL ENGINEER
Dec 22/77

Fig. U-UN4

ARCHER, CATHRO & ASSOCIATES LTD

GEOLOGY, RECONNAISSANCE GEOCHEMISTRY AND RADIOMETRICS


SURPRIZE 1-95 CLAIMS
UKON JOINT VENTURE

SCALE - 1:10,000

200 100 0 200 400 600 800
METRES

NTS 116 B/3

To accompany a report by A R Archer dated Dec 1977



LEGEND

- **eTqfp** — quartz feldspar porphyry, rounded smoky quartz eyes
- **Psn** — Schist-Gneiss Unit – dark green to grey chlorite schist, minor carbonaceous phyllite, marble and foliated greenstone
- soil sample value in ppm uranium
- rock sample value in ppm uranium
- water sample value in ppb uranium
- approximate geological contact
- claimpost location and claim numbers
- bulldozer trench

Association of Professional Engineers of Yukon
A. R. ARCHER
GEOLOGICAL ENGINEER
DEC 23/77

Fig. U-UN5

ARCHER, CATHRO & ASSOCIATES LTD

GEOCHEMISTRY, GEOLOGY

UNEXPECTED PROPERTY


SURPRIZE 1-32 CLAIMS

UKON JOINT VENTURE

SCALE – 1:5000

100 0 100 200 300 400
Metres

To accompany a report by A.R. Archer dated Dec. 1977



LEGEND

- ²Tqfp — quartz feldspar porphyry, rounded smoky quartz eyes
- Psn — Schist-Gneiss Unit – dark green to grey chlorite schist, minor carbonaceous phyllite, marble and foliated greenstone
- •/10 — total count at waist height using a Scintrex IGS-ISL broadband scintillometer (43 cc crystal)
- ANOMALY A — location of geochemical anomaly (see Fig. U-UN5)
- — approximate geological boundary
- claimpost location and claim numbers
- bulldozer trench

Association of Professional Engineers of the Yukon Territory
A. R. ARCHER
C. E. R. O.
GEOLOGICAL
ENGINEER
Dec 29/77

Fig. U-UN6

ARCHER, CATHRO & ASSOCIATES LTD

RADIOMETRICS, GEOLOGY

UNEXPECTED PROPERTY

SURPRIZE 1-32 CLAIMS

UKON JOINT VENTURE

SCALE – 1:5000

400 0 100 200 300 400

Metres

To accompany a report by A.R. Archer dated Dec. 1977.