

MOUNTAINEER MINES LTD. - PAN OCEAN OIL LTD.

JOINT VENTURE

GEOLOGICAL AND GEOCHEMICAL REPORT


ON THE

URSUS 1-24 MINERAL CLAIMS

N.T.S. 106-D-16

64°55'N 134°15'W

YUKON TERRITORY



by

D. A. Yeager - Geologist

R. K. Yorston - Geologist

C. K. Ikona - P.Eng.

090425

This report has been examined by the Geological Evaluation Unit and is recommended to the Commissioner to be considered as representation work in the amount of

\$ 24 00.00

D.B. Craig 26/2/79

Resident Geologist or
~~Resident Mining Engineer~~

Considered as representation work under
Section 53 (4) Yukon Quartz Mining Act.

[Signature]
B. R. BAXTER
Supervising Mining Recorder

[Signature]
Commissioner of Yukon Territory

TABLE OF CONTENTS

	<u>Page</u>
1.0 INTRODUCTION	1
2.0 LIST OF CLAIMS	1
3.0 LOCATION AND ACCESS	1
Figure 1 - Property Location Map	after page 1
Figure 2 - Claim Map	after page 1
4.0 TOPOGRAPHY AND VEGETATION	2
5.0 REGIONAL GEOLOGY	2
5.1 Structure	4
6.0 PROPERTY GEOLOGY	
6.1 Introduction	5
6.2 Lithology	6
6.3 Structure and Stratigraphy	8
6.4 Mineralization	9
Table 6.4.1 - Assays (1977)	11
7.0 GEOCHEMISTRY	
7.1 Introduction	13
7.2 Water Geochemistry	14
7.3 Stream Sediment Geochemistry	14
7.4 Soil Geochemistry	14
7.5 Rock Geochemistry	15
Table 7.5.1 - Rock Geochemistry Results	15
8.0 DISCUSSION AND CONCLUSIONS	17
9.0 RECOMMENDATIONS	18

LIST OF APPENDICES

APPENDIX I	Engineer's Certificate
APPENDIX II	- Assay Procedures
APPENDIX III	List of Personnel
APPENDIX IV	Affidavit of Expenditures
APPENDIX V	Statement of Expenditures
APPENDIX VI	- Assay Certificates
APPENDIX VII	- Figure 3 - Geology
APPENDIX VIII	- Figure 4 - Uranium in streams and soil

1.0 INTRODUCTION

The URSUS 1-24 mineral claims were staked on August 26, 1976 by Harman Management Ltd. to cover uranium and copper showings discovered by prospectors Doug Fulcher and Ward Harrison during a regional prospecting program carried out for Mountaineer Mines Ltd. A brief investigation of the property was conducted by Harman Management Ltd. subsequent to staking the ground. Pan Ocean Oil Ltd. of Calgary acquired majority interest in the claims in the fall of 1976.

During the period August 19 to August 25, 1977, a preliminary prospecting, geologic, and geophysical investigation was carried out on the property. The work program was done by Pamicon Developments Ltd. under the field supervision of R. Darney.

From June 9, 1978 to September 10, 1978, additional mapping and geochemical prospecting was done in the claims area by Pamicon Developments Ltd.

2.0 LIST OF CLAIMS

<u>CLAIM NAME</u>	<u>STAKING DATE</u>	<u>RECORDING DATE</u>	<u>GRANT NO.</u>
URSUS 1-24	August 26/76	September 16/76	YA6875-6890 incl. YA6997-7004 incl.

Claim posts examined by the author appear to conform with the Yukon Quartz Mining Act regulations.


3.0 LOCATION AND ACCESS

The URSUS claims are located on N.T.S. sheet 106-D-16 approximately 100 miles north-northeast of Mayo, Y.T. The property lies 15 miles south-southeast of Quartet Lakes, 14 miles northwest of Gillespie Lake and 6 miles north of

YUKON LOCATION MAP

URSUS GROUP

SCALE IN MILES



DRAWN


Altair


PROJECT

DATE

NOV. 1977

FIG. I





2	4	6	8	10	12
1	3	5	7	9	11
14	16	18	20	22	24
13	15	17	19	21	23

MOUNTAINEER-PAN OCEAN JOINT VENTURE			
URSUS CLAIMS			
CLAIM LOCATION MAP			
QUARTET LAKES AREA			
YUKON TERRITORY			
SCALE IN MILES			
1/2	1/4	0	1/2
DRAWN Altair	PROJECT	DATE NOV 1977	FIG. 2

the Bear River airstrip at approximately 64°55'N. latitude and 134°15'W. longitude.

Access to the property is by float equipped, fixed wing aircraft from Mayo to either Gillespie Lake or Quartet Lakes. Wheeled aircraft may use the serviceable Bear River airstrip. Helicopter transport from any of the three fixed wing landing locations is required to reach the property. Both helicopter and fixed wing aircraft as well as full expediting services are available in Mayo.

4.0 TOPOGRAPHY AND VEGETATION

Elevations on the property range from 3,400 feet to 6,200 feet A.S.L. and topography is rugged over most parts of the area. Bedrock exposure on the claim group is approximately 50% due to talus and overburden cover at the lower elevations. Outcrop is found mainly along ridge lines and eroded stream banks. The entire claim group lies above tree line so vegetation consists of low grasses and moss. There are some willow thickets in the larger stream beds.

5.0 REGIONAL GEOLOGY

The Quartet-Fairchild region lies in the Wernecke Mountains of the north eastern Yukon Territory. In the general area, the Werneckes consist of local ranges which include the Rackla Range, Bonnet Plume Range and Knorr Range. Topography is normally moderate to rugged with

elevations ranging from 2,000 to 6,500 feet. The major river valleys are broad, timbered and extensively overburden covered, while most mountain slopes present greater than 60% outcrop above the 4,000 foot level.

The entire area has been mapped by the Geological Survey of Canada and three separate publications are presented. The following memoir and open file reports give 1" = 4 miles geological coverage of the Nash Creek, Nadaleen River, Wind River and Snake River map areas.

- (1) Geology of Nash Creek, Larsen Creek and Dawson Map-Area, Yukon Territory by L.H. Green 1972 (Memoir 364).
- (2) Open File 205 (Geology of Nadaleen River and Bonnet Plume Lake Map sheets by S. Blusson) 1975.
- (3) Open File 279 (Geology of Snake River and Wind River sheets by D.K. Norris) 1975.

In the Quartet-Fairchild-Gillespie Lakes region Helikian rocks are exposed over an area of some 1,500 sq. miles in a roughly circular fashion centered near Longitude 134°00W and Latitude 65°00'N.

These rocks have been described as Units 1 & 2 by L. Green on the Nash Creek Sheet.

Recent G.S.C. stratigraphic work by Bell and Delaney (1976) has redesignated Units 1a, 1 and 2 (Green 1972) as Units A, B, and C respectively. The unit designations as established by Bell and Delaney will be used in this report.

Unit A, whose base is not exposed, is composed of a thick succession of moderately metamorphosed fine grained

clastic sediments with interbedded carbonates. The overlying Unit B consists of thinly interbedded slates and argillites with occasional quartzite beds.

Unit C, which conformably overlies the uppermost slate-quartzite section of Unit B, consists mainly of thickly bedded orange weathering dolomites. The base of the unit is marked by a series of transitional beds of alternating buff weathering dolomites and interbedded slates and quartzites.

Erratically distributed throughout the Proterozoic metasediments are irregularly shaped breccia bodes. The breccia zones vary from tens of feet to several thousand feet in size and appear as cross cutting pipe-like features at all levels in the stratigraphic column. Several varieties exist, but all exhibit an assortment of angular clasts derived from rock types common to the area. Hornfels margins observed at several localities indicate an intrusive origin.

A common association with many of the breccia bodes are zones of veining or locally pervasive feldspar alteration seen as internal features within the breccias or in host rocks adjacent to them.

The alteration zones are pink in colour due to either K-spar or strong hematization and in some instances contain varying amounts of specularite, chalcopyrite and minor uranium mineralization.

5.1 Structure

Two major periods of deformation have taken place within the Wernecke Mountain region. During the first period or

Racklan Orogeny, the Proterozoic rocks of Units A, B, and C underwent intense folding and faulting. Folds are tight to isoclinal with the development of strong axial plane cleavage and commonly an almost vertical foliation.

A major unconformity of Lower Hadrynian age forms the upper contact of Unit C. In many localities, erosion beneath this unconformity has resulted in the complete removal of Unit C and the strong angular relationship between the relatively flat lying Cambrian and younger rocks directly overlying Units A and B is apparent.

Further unconformities near the Upper Hadrynian, Lower Cambrian and Upper Cambrian margins leave Devonian carbonates directly over the Helikian section.

The second period of deformation, which involves both Paleozoic and Proterozoic strata, is weak compared to the first. This is particularly evident in the younger Carbonate sections to the west and southwest where deformation consists mainly of broad open folding and minor overthrusting.

6.0 PROPERTY GEOLOGY

6.1 Introduction

Preliminary mapping of the URSUS claims was carried out at a scale of 1 inch to 1,000 feet (see Fig. 3 this report). Assay samples were taken from any zones of mineralization encountered and locations and results are presented in Fig. 3 and Table 6.4.1 respectively. The property is underlain entirely by Units A and C. Faulting is very common

in the area and breccia intrusion with resultant folding, shearing and metasomatism was noted in many localities. Uranium and copper mineralization were found mainly in the breccias and their related alteration zones.

6.2 Lithology

The lowest rocks exposed on the claim group, Unit 1, are composed of thin to medium bedded, variably limonitic, grey-green siltstone. This unit is only exposed in the main creek cut that runs through the western portion of the property then bends and runs across the northern part of the claims. The unit in places has phyllitic partings and exhibits gradational contacts with the phyllites in the area.

Unit 2 is made up of a number of smaller sub-units of dark green siltstones, mudstones, argillites and related transitional rock types. Bedding thicknesses range from laminar banding to several inches. Several mineralized breccia bodies occur within Unit 2.

Unit 3 is composed of grey-green phyllites. Original bedding features are sometimes preserved in the phyllites as compositional layering, however generally only phyllitic cleavages could be measured. The phyllites exhibited kink banding and cleavage plane crenulations; minerals mainly noted were chlorite and sericite.

Unit 4 consists of thinly interbedded light green and purple siltstones. The unit mainly occurs at higher elevations on the property, close to the contact with the overlying Unit C.

Clasts of Unit 4 rocks appear in a number of the breccia bodies at higher elevations.

Unit 5 is a dark brown to grey weathering, medium to thick bedded dolomitic siltstone. The unit was only observed near the contact with the Unit C orange weathering dolomites and was the major rock type in most of the breccias mapped in that area. Many large blocks (up to 300 feet) of the dolomitic siltstone were seen as breccia fragments. The unit is variably metasomatized, resulting in obliteration of bedding features, migration and flooding of dolomite, and occasionally dissemination of dolomite in tiny rhombohedral porphyroblasts throughout the rock.

The overlying Unit C dolomites are grey to orange weathering, thin to thick bedded, and contained occasional sections with shale interbeds.

A large number of intrusive breccia bodies outcrop within the claims boundary. They were noted to intrude all the lithologic types found on the property with the exception of the Unit C dolomites. They are made up of clasts derived from stratigraphically nearby rock types and in many cases resulted from nothing more than a shattering and re-orientation of material virtually in situ. It appeared that the breccia bodies having a greater variety of size and composition of clasts (hence greater transport distances?) were accompanied by more obvious indications of hydrothermal activity. These indications included chloritic and/or siliceous alteration haloes in the country

rock around the bodies as well as feldspathization, hematization, silicification, and dolomitization within and immediately adjacent to the bodies themselves. Copper and uranium mineralization were noted in several of the breccia bodies or in altered rocks contacted by the breccias.

6.3 Structure and Stratigraphy

As is commonly the case in areas of intense breccia emplacement, extreme structural deformation has occurred in the rocks within the URSUS claims. The folding is very complex. In many cases the folds can be seen to be the result of warping and dragging along fault surfaces, however they more commonly appear to be a chaotic style of deformation related to the breccia intrusions. This folding has resulted in extreme fracturing in many locations on the property with resultant quartz, feldspar, and dolomite veining and flooding. Minor fold axes in the Unit C dolomites on the other hand tend to parallel the strikes of beds within the unit and are therefore thought to be drag folds related to the underlying thrust fault.

The thrust fault bringing the Unit C dolomites in contact with the Unit A siltstones is the most predominant fault feature in the area and extends for many tens of miles in either direction from the property. The prevailing fault direction in the Unit A rocks is 070° to 080° , roughly parallel to the large regional fault. A large number of faults and shears with other orientations were noted during the mapping, the most important of which are plotted in Figure 3.

The portion of Unit A found on the claims is thought to be stratigraphically high in the section as the dolomitic siltstones of unit 5 resemble those which lie just beneath Unit B.

6.4 Mineralization

A number of copper and uranium showings are located along a band of Unit 2 siltstones trending in an east-west direction through the central part of the property. The showings are associated with breccia bodies or their alteration haloes and mostly occur near the contact with the overlying Unit 3 phyllites. Assay results are presented in Table 6.4.1.

The East Showing is exposed on a talus covered knob near the middle of the eastern boundary of the property. Chalcopyrite and brannerite occur sporadically in a bleached metasiltstone bed within Unit 2. The bed has been significantly altered by silicification, feldspathization, and hematization to a white to pink weathering silicified siltstone resembling a quartzite in outcrop. The brannerite occurs along small shear zones and in associated fractures, as well as disseminated through the more heavily feldspathized parts of the siltstone bed. Disseminated chalcopyrite also occurs in the feldspathized uranium rich pods. Shattering has taken place along some of the shear zones in the area resulting in occasional breccia lenses in the siltstone. Chalcopyrite fills spaces in the breccia matrices up to several millimeters in size between breccia fragments

ranging from 2 m.m. to 15 m.m. in size. The mineralized pods were generally less than 5 feet by 5 feet in size and were spaced from 75 feet to 200 feet apart.

The Central Showing is located in the deeply incised cut of a northerly flowing stream crossing the central part of the property. This is approximately 2,400 feet west-northwest of the East Showing. Chalcopyrite was seen associated with quartz/feldspar flooding in an interlaminated siltstone/silty dolomite bed within Unit 2. The flooding has occurred along fractures and bedding planes and locally contains good grade copper mineralization. Immediately overlying the copper mineralized siltstone/silty dolomite bed, uranium mineralization occurs as fine grained brannerite particles in fracture controlled hematite stringers as well as discrete grains of brannerite up to 1 m.m. in size in and around a breccia body intruding a silicified siltstone. The mineralization is associated with quartz veining and feldspar flooding. Discrete grains of chalcopyrite up to 2 m.m. in size also occur in the feldspathized rocks. Pods of reddish brown barite up to 1 foot in size were noted in several localities in the breccia. The mineralized pods appeared to be in the order of 5 feet by 5 feet in size and were spaced approximately 150 feet apart.

Two other showings were sampled during the mapping: In the first, brannerite was found in a feldspathized zone at the edge of a breccia body approximately 2,000 feet west-northwest of the Central Showing. Three uraniferous pods, each

several square feet in size, were noted within a 20 foot by 20 foot alteration zone. In the second showing, chalcopyrite occurs in fractures in a breccia body as well as in the adjacent silicified siltstone. The mineralized zone was approximately 50 feet by 10 feet in size.

A number of other breccia related uranium and copper occurrences were noted during the mapping but were too limited in size and grade to warrant sampling.

Table 6.4.1. - Assays (1977)

<u>Sample No.</u>	<u>% U₃O₈</u>	<u>Description</u>
62611	-	Assays 1.07% Cu. Selected grab sample from talus of better grade material. Source in grey-brown weathering siltstone breccia and in silicified siltstone.
62614	0.011	Continuous chip sample across 3 feet of bedding in outcrop of white, silicified siltstone. Rock heavily fractured, exhibits hematized alteration haloes. Sample from East Showing.
62615	0.036	Random grab sample from sub-outcrop across 2 foot mineralized pod in silicified siltstone breccia. Located approx. 2,000 feet WNW of Central Showing.
62616	0.003	Continuous chip sample from outcrop along 3 feet of 2 inch to 6 inch thick silicified, hematized zone in metasomatized, breccia flooded siltstone. Approx. 2,000 feet WNW of Central showing.
62617	0.002	Continuous chip sample from outcrop across 4 foot wide hematized pod in siltstone breccia. Approx. 2,000 feet WNW of Central Showing.

Table 6.4.1 Continued..

<u>Sample No.</u>	<u>% U₃O₈</u>	<u>Description</u>
62618	0.008	Continuous chip sample across 4.5 feet of outcrop of silicified, hematized siltstone. Entire hematized area is approx. 8 feet in size. From East Showing.
62619	0.024	Continuous chip sample from outcrop across 1 foot by 1 foot pod of limonitic silicified siltstone. Abundant chlorite, specular hematite in surrounding rocks. Approx. 40 feet from 62618. From East Showing.
62620	0.016	Random grab sample from 10 foot by 10 foot talus train of abundantly hematized silicified siltstone. Approx. 200 feet NW of 62619. From East Showing.
62621	0.046	Selected grab sample of better grade material from sporadically mineralized talus train approx. 5 feet by 150 feet in size. Specular hematite and brannerite noted in silicified siltstone. Approx. 100 feet NW of 62620. From East Showing.
62622	0.020	Grab sample from talus from 10 foot by 100 foot area of pink, silicified siltstone. Mineralization associated with approx. 6 inch wide. From East Showing.
62623	0.042	Selected grab sample of better grade material from lower end of same talus train as 62622. From East Showing.
62624	0.007	Grab sample from talus train of tan to pink, hematized, silicified siltstone at west end of Central Showing. Brannerite with specular hematite on fractures.

Table 6.4.1 Continued ..

<u>Sample No.</u>	<u>% U₃O₈</u>	<u>Description</u>
62625	0.030	Grab sample from 40 foot long talus train at approx. mid-point of Central Showing. Siltstone has been progressively metasomatized, brecciated and specular hematite flooded.
62626	0.016	Grab sample from talus train of breccia boulders containing 6 inch wide dykelet of syenitic material. Rocks abundantly hematized. This from east end of Central Showing.
62627	0.212	Grab sample from mineralized pods in talus train of breccia boulders at extreme NE end of Central Showing. Visible brannerite in hematized pods.

7.0 GEOCHEMISTRY

7.1 Introduction

The geochemical program in the URSUS area consisted of silt and water sampling the main drainages in the area at approximately 500 foot intervals as well as any tributary streams depending on the availability of silt and water in those tributaries. In addition, two contour soil lines were run on either side of the valley. Soil samples were taken at approximately 300 foot intervals. Sample locations and values are presented in Figure 4 of this report. Complete descriptions of analysis procedures and certificates of analysis are included in Appendix II and Appendix VI respectively.

7.2 Water Geochemistry

Twenty-seven water samples were collected during the survey. Samples were taken in numbered, acid cleansed, plastic sample bottles and sent for analysis to Chemex Labs. Ltd. in North Vancouver, B.C. Upon receipt at Chemex, the samples were analysed for uranium using standard fluorometric procedures.

Values ranged from 1.7 to 17.8 parts per billion uranium. Using the sample classification scheme presented in Figure 4, seven samples are second order anomalous and twenty are first order anomalous. None of the samples taken fall within background levels.

7.3 Stream Sediment Geochemistry

Twenty-five silt samples were collected during the survey. Samples of silt size sediments were taken from active streams and placed in numbered kraft envelopes. The samples were dried in camp then sent for analysis to Chemex Labs. Ltd. in North Vancouver, B.C. Upon receipt at Chemex, the samples were analysed for uranium using standard fluorometric procedures.

Values ranged from less than 2.0 to 7.0 parts per million uranium. Using the classification scheme presented in Figure 4, two samples are first order anomalous and twenty-three fall within background levels.

7.4 Soil Geochemistry

One hundred and forty soil samples were collected during the survey. The samples were taken from the B and C soil horizons and placed in numbered kraft envelopes. They were dried in camp

then sent for analysis to Chemex Labs. Ltd. in North Vancouver, B.C. Upon receipt at Chemex the samples were analysed for uranium using standard fluorometric procedures.

Values ranged from less than 0.5 to 88 parts per million uranium. Using the classification scheme presented in Figure 4, three samples are second order anomalous, two samples are first order anomalous, and ninety-seven samples are within background limits. The remaining thirty-seven samples exhibited fluorescence quenching due to high levels of interfering ions.

7.5 Rock Geochemistry

Twenty-four rock samples were collected during the survey and analysed for uranium at Chemex Labs. Ltd. in North Vancouver, B.C. using standard fluorometric procedures. (See Figure 3 for sample locations.)

Values ranged from less than 0.5 to 3.0 parts per million uranium. As very few rock geochemistry samples have been taken in the area, no statistical breakdown of anomalous levels has been compiled.

Table 7.5.1 - Rock Geochemistry Results

<u>Sample No.</u>	<u>Unit Thickness</u>	<u>Description</u>
1	75'	Dark grey to black pyritic shale; rusty weathering; <0.5 ppm.
2	25'	Brown weathering dolomite with thin interbedded silicified light grey siltstone; some pyrite; <0.5 ppm
3	12'	Shear Zone, minor folding and kinking. Rock is dark grey green chloritized and slickensided minor rust; <0.5 ppm.

Table 7.5.1 - Continued..

<u>Sample No.</u>	<u>Unit Thickness</u>	<u>Description</u>
4	125'	Grey siltstone. Occasional dolomitic band (every 10'). Pyritic but not noticeable rusty weathering; 1.0 ppm.
5	200'	Dark grey shale to silty shale. Pyritic rusty weathering. Tight isoclinal folding; <0.5 ppm.
6	30'	Dark grey to black shale with partially developed slaty cleavage; 3.0 ppm
7	250'	Drab green phyllite. Rock grades to light green thin bedded altered siltstone small scale open folds are common; <0.5 ppm.
8	150'	Dark grey-brown weathering sandstone with thin interbeds of dark grey to black shaley material. Sandstone units reach 16' in thickness;<0.5 ppm.
9	20'	Med. to darker green fractured phyllite slightly talcy or serpentinized; <0.5 ppm.
10	300'	Light grey-green silicified siltstone thinly bedded; <0.5 ppm.
11	80'	Dark grey silicified siltstone;<0.5ppm
12	40'	Dark grey very thin bedded grey siltstone alternating with lighter thin bedded siltstone; <0.5 ppm.
13	20'	Dark green massively bedded pyritic and chloritized siltstone; <0.5 ppm.
14	100'	Light grey siliceous chloritic siltstone; weak schistosity; pyritic zones of up to 4' in thickness; <0.5 ppm.
15	50'	Increase in qtz and pyrite from previous sample; <0.5 ppm.
16	200'	Thick beds of sandstone and thin bands of dark grey silty shale; <0.5 ppm.

Table 7.5.1 - Continued..

<u>Sample No.</u>	<u>Unit Thickness</u>	<u>Description</u>
17	200'	Tan weathering thin bedded dolomitic siltstone; rock has slaty cleavage and parts along a very thin light green phyllite surface; <0.5 ppm.
18	350'	Dark grey to black very siliceous slate; some py; 1.0 ppm
19	150'	Very silicified siltstone. Some thin dolomite interbeds. Qtz flooding Increasing dolomite upsection; <0.5ppm.
20	400'	Dark grey to black thin bedded slate interbedded with lesser thinbedded dolomite; <0.5 ppm.
21	150'	Talus - dark grey-black silty shale; 1.0 ppm.
22	25'	Light grey very thin bedded to laminated siliceous siltstone; <0.5ppm
23	75'	Siliceous brown weathering dolomite with slaty partings; some pyrite; <4.0 ppm.
24	75'	Dark grey-black siliceous very fine grained shale; some pyrite; <0.5 ppm.

8.0 DISCUSSION AND CONCLUSIONS

All the second order anomalous water samples and the first order anomalous silt samples taken during the survey came from creeks running within the claims boundary. However, the best silt and water sample values were taken from streams entering the property from the north and northwest.

The soil and rock geochemistry program did not indicate any anomalous areas with the exception of a few spot highs.

It is felt that the degree of structural deformation, chemical alteration, and breccia intrusion noted during the mapping of the property has created a suitable geologic environment for uranium deposition. In addition, the promising stream geochemistry results indicates the need for further work in the area.

9.0 RECOMMENDATIONS

It is recommended that the URSUS 1-24 claims be retained on assessment credits and that a continuing program of close spaced silt and water sampling be carried out in streams draining the northern and northwestern claims area.

Respectfully submitted,


R. K. Yorston - Geologist

David A. Yeager

D. A. Yeager - Geologist

C. K. Ikona - P.Eng.

C. K. Ikona



CHEMEX LABS LTD.

212 BROOKSBANK AVE.
 NORTH VANCOUVER, B.C.
 CANADA V7J 2C1
 TELEPHONE: 985-0648
 AREA CODE: 604
 TELEX: 043-52597

• ANALYTICAL CHEMISTS • GEOCHEMISTS • REGISTERED ASSAYERS

CERTIFICATE OF ANALYSIS

TO: Pamicon Developments Ltd.,
 208 - 850 W. Hastings St.,
 Vancouver, B. C.

CERTIFICATE NO. 46235
 INVOICE NO. 28597
 RECEIVED Oct. 11/78
 ANALYSED Oct. 16/78

ATTN:

SAMPLE NO. :	PPM Uranium	
U 1	< 4.0	
2	0.5	
3	0.5	
4	< 4.0	
5	< 4.0	
6	< 0.5	
7	< 0.5	
8	< 0.5	
9	< 0.5	
10	< 0.5	
11	< 0.5	NOTE: Less than 4 detection limit due to fluorescence quenching caused by high concentration of the interfering metals.
12	< 0.5	
13	< 0.5	
14	< 0.5	
15	0.5	
16	< 0.5	✓
17	< 0.5	
18	0.5	
19	< 0.5	
20	0.5	
21	< 0.5	
22	< 0.5	
23	< 4.0	
24	< 0.5	
25	1.0	
27	< 4.0	
28	< 0.5	
29	< 0.5	
30	< 0.5	
31	< 0.5	
32	< 0.5	
33	< 0.5	
34	< 0.5	
35	0.5	
36	< 4.0	
37	< 0.5	
38	< 0.5	
39	4.5	
40	< 0.5	
U 41	< 0.5	
Std.	20	



MEMBER
 CANADIAN TESTING
 ASSOCIATION

CERTIFIED BY: 



CHEMEX LABS LTD.

212 BROOKSBANK AVE.
 NORTH VANCOUVER, B.C.
 CANADA V7J 2C1
 TELEPHONE: 985-0648
 AREA CODE: 604
 TELEX: 043-52597

• ANALYTICAL CHEMISTS • GEOCHEMISTS • REGISTERED ASSAYERS

CERTIFICATE OF ANALYSIS

TO: Pamicon Developments Ltd.,
 208 - 850 W. Hastings St.,
 Vancouver, B.C.

ATTN:

CERTIFICATE NO. 46236
 INVOICE NO. 28597
 RECEIVED Oct. 11/78
 ANALYSED Oct. 17/78

SAMPLE NO. :	PPM Uranium	
U 42	1.0	
43	< 0.5	
44	< 0.5	
45	< 0.5	
46	< 0.5	
47	< 4.0	
48	< 0.5	
49	< 0.5	
50	< 0.5	
51	< 0.5	
52	< 4.0	
53	< 4.0	
54	< 4.0	
55	< 4.0	
56	< 4.0	
57	< 0.5	Note: Less than 4 detection limit due to fluorescence quenching caused by high concentration of the interfering metals.
58	< 0.5	
59	< 4.0	
60	< 4.0	
61	< 4.0	
62	< 4.0	
63	< 4.0	
64	< 4.0	
65	< 0.5	
66	< 0.5	
67	< 0.5	
68	< 0.5	
69	< 0.5	
70	< 4.0	
71	< 4.0	
72	< 4.0	
73	< 0.5	
74	< 0.5	
75	< 0.5	
76	< 0.5	
77	< 0.5	
78	< 0.5	
79	< 0.5	
80	< 0.5	
U 81	< 0.5	
Std.	20	



MEMBER
 CANADIAN TESTING
 ASSOCIATION

CERTIFIED BY: *[Signature]*



CHEMEX LABS LTD.

212 BROOKSBANK AVE.
 NORTH VANCOUVER, B.C.
 CANADA V7J 2C1
 TELEPHONE: 985-0648
 AREA CODE: 604
 TELEX: 043-52597

• ANALYTICAL CHEMISTS • GEOCHEMISTS • REGISTERED ASSAYERS

CERTIFICATE OF ANALYSIS

CERTIFICATE NO. 46237
 INVOICE NO. 28597
 RECEIVED Oct. 11/78
 ANALYSED Oct. 16/78

TO: Pamicon Developments Ltd.,
 208 - 850 W. Hastings St.,
 Vancouver, B.C.

ATTN:

SAMPLE NO. :	PPM	
	Uranium	
U 82	< 0.5	
83	< 0.5	
84	< 0.5	
85	< 0.5	
86	< 4.0	
87	< 4.0	
88	< 4.0	
89	< 4.0	
90	< 0.5	
91	< 0.5	
92	< 4.0	
93	< 0.5	
94	0.5	
96	< 0.5	
97	< 0.5	
98	< 0.5	
99	< 0.5	
100	< 0.5	Note: Less than 4 detection limit due to fluorescence quenching caused by high concentration of the interfering metals.
101	< 4.0	
150	< 4.0	
152	< 0.5	
153	3.5	
154	< 0.5	
U 155	< 0.5	



MEMBER
 CANADIAN TESTING
 ASSOCIATION

CERTIFIED BY: *[Signature]*



CHEMEX LABS LTD.

212 BROOKSBANK AVE.
 NORTH VANCOUVER, B.C.
 CANADA V7J 2C1
 TELEPHONE: 985-0648
 AREA CODE: 604
 TELEX: 043-52597

• ANALYTICAL CHEMISTS • GEOCHEMISTS • REGISTERED ASSAYERS

CERTIFICATE OF ANALYSIS

TO: Pamicon Developments Ltd.,
 610 - 850 West Hastings Street,
 Vancouver, B.C.
 ATTN: V6B 1P1

cc: Mayo, Y. T.

CERTIFICATE NO. 43440
 INVOICE NO. 25177
 RECEIVED June 16, 1978
 ANALYSED June 21, 1978

SAMPLE NO. :	PPM
	Uranium
BRECCIA CREEK #1	<0.5
2	<0.5
3	<0.5
4	1.0
5	<0.5
6	3.0
7	<0.5
9	<0.5
9	<0.5
10	<0.5
11	<0.5
12	<0.5
13	<0.5
14	<0.5
15	<0.5
16	<0.5
17	<0.5
18	1.0
19	<0.5
20	<0.5
21	1.0
22	<0.5
23	<4.0
BRECCIA CREEK #24	<0.5



MEMBER
 CANADIAN TESTING
 ASSOCIATION

CERTIFIED BY: 



CHEMEX LABS LTD.

212 BROOKSBANK AVE.
NORTH VANCOUVER, B.C.
CANADA V7J 2C1
TELEPHONE: 985-0648
AREA CODE: 604
TELEX: 043-52597

• ANALYTICAL CHEMISTS •GEOCHEMISTS ••REGISTERED ASSAYERS

CERTIFICATE OF ANALYSIS

TO: Pamicon Developments Ltd.,
610 - 850 W. Hastings
Vancouver, B.C.
V6B 1P1

ATTN: David Yeager

cc: Mayo

CERTIFICATE NO. 43647
INVOICE NO. 26314
RECEIVED June 23/78
ANALYSED June 29/78


SAMPLE NO. :	PPM Uranium
BX 1	< 0.5
2	< 0.5
3	< 0.5
4	1.0
5	< 0.5
6	88
7	< 4.0
8	< 4.0
9	< 4.0
10	< 4.0
11	< 4.0
12	< 4.0
13	< 4.0
14	< 4.0
15	< 4.0
16	< 0.5
17	< 0.5
BX 18	< 0.5

Braccia ck

Note: Less than 4 detection limit due to fluorescence quenching caused by high concentration of the interfering metals.



CERTIFIED BY: *[Signature]*




CLASSIFICATION OF GEOCHEMICAL SAMPLES

	SOILS ●	SILTS ○	WATER □
BACKGROUND	< 0.8 ppm	< 5 ppm	< 13 ppb
1 ST ORDER ANOMALOUS	0.8 - 3.3 ppm	5 - 35 ppm	1.3 - 9.0 ppb
2 ND ORDER ANOMALOUS	> 3.3 ppm	> 35 ppm	> 9.0 ppb

MOUNTAINEER-PAN OCEAN JOINT VENTURE

URSUS I-24 CLAIM GROUP
 NTS 106-D-16
URANIUM IN STREAMS & SOIL
 YUKON TERRITORY



PAMICON DEVELOPMENTS LIMITED

DRAWN: Altair	PROJECT: Fairchild	DATE: JAN. 1979	FIGURE: 4
------------------	-----------------------	--------------------	--------------