

094206

**REPORT ON
ROCK AND SOIL SAMPLING AND ROAD RECLAMATION
on the**

**CANADIAN CREEK PROPERTY
(ANA, KOFFEE, AZTEC
MAYA AND ICE CLAIMS)**

DAWSON RANGE, YUKON TERRITORY

NTS: 115J/10

Latitude: 62° 44' N, Longitude: 138° 57' W

FOR

EASTFIELD RESOURCES LTD.


And

ALEXIS RESOURCES LTD.

by

J.W. Morton P.Geo.

May, 2000



This report has been examined by
the Geological Evaluation Unit
under Section 53 (4) Yukon Quartz
Mining Act and is allowed as
representation work in the amount
of \$ 12,000.

M. B. Buck
Regional Manager, Exploration and
Geological Services for Commissioner
of Yukon Territory.

TABLE OF CONTENTS

	PAGE NO.
INTRODUCTION	1
CLIMATE, TOPOGRAPHY AND VEGETATION	2
PROPERTY STATUS	2
GEOLOGY	3
RECOMMENDATIONS	4
COST STATEMENT	4
AUTHOR QUALIFICATIONS	5
GENERAL LOCATION MAP	FIGURE 1
CLAIM MAP	FIGURE 2
GEOLOGY AND ROCK SAMPLE LOCATIONS (ANA AREA)	FIGURE 4
GEOLOGY AND ROCK SAMPLE LOCATIONS (KOFFE AREA)	FIGURE 5
SOIL GEOCHEMICAL MAP KOFFEE AREA (Cu, Mo, Au)	FIGURE 6
SOIL GEOCHEMICAL MAP KOFFEE AREA (Pb, Zn, Ag, As)	FIGURE 7
ROCK SAMPLE DESCRIPTIONS	APPENDIX 1
TRENCH SAMPLE DESCRIPTIONS	APPENDIX 2
PETROGRAPHIC DESCRIPTION	APPENDIX 3
GEOCEMICAL CERTIFICAATE S	APPENDIX 4

INTRODUCTION


The Canadian Creek Property is located approximately 150 km south of Dawson City, in the Dawson Range, Yukon Territory. It is situated within the Whitehorse Mining District on NTS map sheets 115J/10, 11 and 14. The property comprises 262 contiguous quartz claims names the ANA, KOFFEE, AZTEC, MAYA, and ICE claims blocks. Eastfield is the recorded beneficial owner of all the claims. Breckenridge Resources Ltd. retains a 17.8 % working interest in the 54 ANA claims. Alexis Resources Ltd has an option to earn a 51% interest in the entire property.

Access to the property is available by using helicopters based in Carmacks and Dawson City and by fixed wing transportation from Whitehorse and Dawson using the Casino airstrip. Access is also available using a combination of river and overland transport employing a 65 ton barge based in Minto. The property road system, which extends from a barge-landing site on the Yukon River at the confluence with Britannia Creek, traverses the central region of the claims. A rough winter road, built in the early 1970's but used as recently as the late 1990's, extends from the property also to roads in the Mount Nansen and Freegold area.


The Canadian Creek property adjoins to the west of the Casino property that hosts a copper-gold-molybdenum porphyry deposit currently owned by Great Basin Gold Ltd. The Casino Deposit has published reserves exceeding 530 million tonnes grading 0.29 g/t gold, 0.26 % copper, and 0.025% MoS₂. Intrusives of the Casino complex, which host the Casino deposit, occur in several locations on the Canadian Creek property.

The 1999 field program consisting of rock and soil sampling and contouring road cuts and disturbed areas from activity that occurred in 1996 and 1997. The program was completed between September 19 and September 26, 1999. Samples originating from road construction in 1997 but which had not been analysed were retrieved and sent for analysis.

Fifty-two hours of bulldozer time (D-6) was expended repairing the road and completing contouring of areas that had previously been disturbed. A total of 93 rock samples and 129 soil samples were collected or retrieved. The results of the rock-sampling program are outlined in appendixes 1 and 2 and are plotted on Figures 8 and 9. The results of soil sampling are plotted (with pre-existing data) of figure 6.



140°



0 100 200
kilometres

EASTFIELD RESOURCES LTD. ALEXIS RESOURCES LTD.		
CANADIAN CREEK PROJECT Whitehorse M.D., Yukon		
General Location Map		
Scale	as shown	NTS
Date	October '96	Drawn by
J.E.L. Lindinger P.Geo.		Figure 1

CLIMATE, TOPOGRAPHY AND VEGETATION

The Property lies within a semiarid subarctic climatic regime. Precipitation is about 20 cm per year, and is about equally divided between rain and snow. Annual temperature range is from -30 to + 20 degrees centigrade with an annual average temperature of -5 degrees. The property lies within the Dawson Range physiographic province. This unglaciated terrain is characterised by tor peaked but otherwise rounded hills of up to 300 meters elevation. Elevation on the property range from 880 meters at the west edge where Coffee Creek leaves the property to over 1670 meters along the ridge near the northeast boundary. The axis of the Dawson Range passes through the Casino deposit and westward through the Canadian Creek property. Lower elevations are forested with coniferous species that give way to isolated stands of black spruce at about 1300 meters. Low buckbrush gives way to alpine grasses and mosses above 1525 meters.

PROPERTY STATUS

Claim Name	Grant No.s	No. Claims	of Expiry Month Day	Expiry Year
Ana 1 -10	YA86735- YA86744	10	17-Feb	2001
Ana 15 -26	YA86749- YA86760	12	17-Feb	2001
Ana 29 -40	YA86763- YA86774	12	17-Feb	2001
Ana 43 -54	YA86777- YA86788	12	17-Feb	2001
Koffee 1 -58	YB37482- YB37539	58	21-Sep	2000
Aztec 1 -12	YB37540- YB37551	12	21-Sep	2000
Aztec 17 -28	YB37556 YB37567	12	21-Sep	2000
Aztec 33 -52	YB37572 YB37591	12	21-Sep	2000

Claim Name	Grant No.s	No. Claims	of Expiry Month Day	Expiry Year
Maya 1 -40	YB37592 Yb37631	40	21-Sep	2000
Ice 1 -74	YB37801- YB37874	74	27-Jan	2001
TOTAL		262	Claims	
GEOLOGY				

Canadian Creek is a gold-copper (molybdenum) porphyry target similar to the adjoining Casino deposit 3 km east of the east property boundary (531 million tonnes grading 0.29 g/t gold, 0.26 % copper, and 0.026% MoS₂). An additional potential exists for the discovery of porphyry related distal gold mineralization which may occur as high grade veins or as lower grade high tonnage disseminated zones.

Upper Cretaceous quartz dioritic to quartz monzonitic intrusives and related breccias named the Casino Complex occur throughout the property. These rocks were formerly interpreted to significantly post date the mid Cretaceous Dawson Range batholithic rocks (quartz diorite to granodiorite). The Dawson Range intrusive rocks in turn intrude Paleozoic aged Yukon Metamorphic Complex. rocks. Recent work completed in 1997 by the Department of Earth and Atmospheric Sciences, University of Alberta (*1) has determined that the age of the Casino Plutonic Suite is indistinguishable from the Dawson Range Batholith – namely 104 million years (mid Cretaceous). Rare earth element content indicates that magmas of the Casino Plutonic suite are late stage fractionated magma derived from the Dawson Range Batholith. The batholith itself is interpreted to be the result of crustal melting that resulted from crustal thickening. A subsequent 70 million-year-old (subduction related) event then intruded the Casino Plutonic Suite. A diagnostic porphyritic unit locally named Patton Porphyry manifests this event. The University of Alberta work genetically relates porphyry mineralization at Casino to the fractionation of the Casino Plutonic Suite. This hypothesis is somewhat at odds with earlier work completed by several groups including Pacific Sentinel Gold Corporation in 1993 and 1994 which concluded that the porphyry mineralization is intimately associated with the younger porphyritic bodies (Patton Porphyry).

(*1) D. Selby, R. Creser and E. Nesbitt, Major and trace element compositions and Sr-Nd-Pb systematics of crystalline rocks from the Dawson Range, Yukon, Canada., Canadian Journal of Earth Sciences, September 1999.

Due to the unglaciated terrain of the central Yukon surface weathering has been ongoing

for at least 20 million years and has resulted in deep oxidation of sulphide bearing rocks, and in particular copper bearing sulphides. The generally recessive mineralized and altered areas additionally challenge exploration.

RECOMMENDATIONS

A program of diamond drilling targets originating from the work completed between 1993 and 1999 is recommended. This program should consist of 10 to 15 holes totaling a minimum of 2000 metres

COST STATEMENT

Item	Supplier	Cost
Professional Fees 3 days @ \$450 day Sept 19-21, 1999	J. Page P.Geo	\$1,350
Field Assistance 6 days @ \$275 day Sept 21-26, 1999	R Horton	\$1,650
Professional Fees 3 days @ \$450 day General Supervision Sept – Dec 1999	J.W. Morton P.Geo	\$1,350
Bulldozer Contract Charges (52 hours @ \$100 hr), Sept 19 – 26, 1999	Tintina Trenching Inc.	\$5,200
6x6 & Equipment Rental 6 days @ \$60 day), Sept 19 – 26, 1999	Tintina Trenching Inc.	\$360
Pickup Truck Rental, Sept 19 – 26, 1999	Tintina Trenching Inc.	\$370
Expenses	Tintina Trenching Inc.	\$707.81
Expenses	J. Page	\$320.98
Groceries		\$620.50
Airfare	J. Page Vancouver-Whitehorse	\$1229.82
Helicopter Charter 4 hrs	Trans North Air	\$3,765.54
Analytical Costs 218 samples @ \$13.54	Acme Analytical Laboratories Ltd.	\$2,950.75
Total		\$19,875.40

AUTHOR QUALIFICATIONS


I, J.W. Morton am a graduate of Carleton University Ottawa with a B.Sc. (1972) in Geology and a graduate of the University of British Columbia with a M. Sc. (1976) in Graduate Studies (Soil Science).

I, J.W Morton have been a member of the Association of Professional Engineers and Geoscientists of the Province of BC (P.Geo.) since 1991.

I, J.W. Morton have practiced my profession since graduation throughout Western Canada, the Western USA and Mexico.

I supervised the work outlined in this report.

Signed this 15th day of May 2000.

A handwritten signature in black ink, appearing to read 'JW Morton', is written over a horizontal line.

J.W Morton P.Geo.

SAMPLE NUMBER	LOCATION			TYPE/DEPTH	LITHOLOGICAL DESCRIPTION	SAMPLE ANALYSIS			
	CLAIM GROUP/AREA	NORTH	WEST			Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)
97-PR-1	ANA - south of Canadian Creek	119+49N	66+05W	Grab sample from outcrop.	Silicious, altered rock with a weak foliation defined by thin layers of limonite and muscovite (after biotite?). Composed largely of quartz with minor pyrite and tourmaline. A phyllic altered YMC quartz-biotite gneiss?	32	80	7	<.3
97-PR-2	ANA - Canadian Creek Road	124+33N	74+13W	Grab sample from road-cut.	Bleached, medium-grained rock, shot through with quartz-tourmaline veins, which vary from 1 mm to 3 cm in thickness. Tourmaline (80%) is dominant over quartz (20%) in the veins. No sulphides noted.	35	4	2	0.3
97-PR-3	KOFFEE - north branch of upper drill access road.	117+44N	102+32W	Grab sample from trench debris. Re-sample of '96 sample C103194.	Rusty-weathering, angular fragments of silicious, clay altered rock. Small, 1-2 mm quartz veins identify 2 fracture sets intersecting at 30 degrees. No sulphides visible, but very limonitic.	46	40	3	<.3
97-PR-4	KOFFEE - Main Koffee road north of saddle.	107+68N	103+65W	Grab sample of sub-crop exposed in ditch.	Rusty-weathering angular fragments of a fine-grained, silicious, intermediate volcanic rock. Small pin-heads of limonite suggest that about 1 percent pyrite existed in this rock prior to leaching.	4	10	5	<.3
97-PR-5	KOFFEE - Main Koffee road north of saddle.	107+50N	103+50W	Grab sample of sub-crop exposed in ditch about 25 metres from 97-PR-4.	Rusty-weathering angular fragments of a fine-grained, silicious, intermediate volcanic rock. Small, open "cusp" shaped fractures, 1-2 cm wide are partly filled with fine-grained pyrite and quartz.	7	11	2	<.3
97-PS-6	KOFFEE - creek zone	117+50N	116+30W	Grab sample of gossen soil.	Rusty-brown, limonitic soil with small angular bits.	35	29	4	0.8
97-PR-7	KOFFEE - creek zone	117+35N	118+05W	Grab sample from stream cut	Silicious breccia, quartz flooding with fine-grain pyrite. Open cavities common.	144	11	3	<.3
97-PR-8	KOFFEE - creek zone	117+35N	118+15W	Grab sample from stream cut. Previous sample at this site was 93-K-9.	Slightly rusty-weathering Hb-Bi-Granodiorite with 1-2 % pyrite stringers containing a trace of bornite? Hornblende appears partly altered to fine-grained biotite +/- chlorite.	7	105	3	<.3
97-PR-9	KOFFEE - creek zone	117+15N	118+90W	Grab sample from stream cut.	Potassic altered granodiorite containing 0.5 -1 % pyrite as fracture fillings, possible trace of bornite.	3	26	2	<.3
97-PR-10	KOFFEE - Ag/As anomaly area	124+13N	93+00W	Grab sample from sub-crop.	Rusty-weathering, quartz-rich intrusive with minor disseminated pyrite.	7	69	58	0.3
97-PR-11	KOFFEE - Ag/As anomaly area	124+13N	93+13W	Grab sample from sub-crop.	YMC dark-colored, fine-grained, equigranular, quartz/biotite gneiss. Weak foliation partly defined by thin, discontinuous limonite layers.	3	45	3	<.3
97-PR-12	KOFFEE - Ag/As anomaly area	124+13N	93+30W	Grab sample from sub-crop.	YMC equigranular fine-medium grained quartzite with limonitic partings giving a shistose appearance..	9	20	6	<.3
97-PR-13	KOFFEE - Ag/As anomaly area	122+32N	94+00W	Grab sample from outcrop.	YMC meta-sediment, rusty-weathering, hard, compact, grey-colored rock. Difficult to break. Weakly defined foliation visible on weathered surfaces. Trace of white-colored pyrite (marcasite?) noted.	8	108	2	<.3
97-PR-14	KOFFEE - Ag/As anomaly area	121+25N	94+00W	Grab sample from outcrop.	YMC dark-colored, rusty-weathering gneiss. Foliation defined by 1 cm layers of quartz alternating with layers of chlorite/muscovite/limonite.	1	13	1	<.3

SAMPLE NUMBER	LOCATION			TYPE/DEPTH	LITHOLOGICAL DESCRIPTION	SAMPLE ANALYSIS			
	CLAIM GROUP/AREA	NORTH	WEST			Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)
SAMPLE NUMBER	CLAIM GROUP/AREA	NORTH	WEST	TYPE/DEPTH	LITHOLOGICAL DESCRIPTION	Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)
97-PR-15	KOFFEE - Ag/As anomaly area	115+25N	94+30W	Grab sample from outcrop (97-MK-1)	YMC dark coloured thinly layered gneiss, rusty weathering with limonitic partings and slightly rusty 1-2 mm quartz veins parallel to cross cutting foliation.	13	40	2	0.3
97-PR-16	KOFFEE - creek zone	117+15N	118+90W	Grab sample from sub-crop by creek, previous samples at this site are 93-K-8, 97-PR-9.	A potassic altered Hb-Bi-granodiorite with Hb altered to chlorite, and biotite/k-spar veins; biotite largely unaltered, feldspars have slight green color. K-spar veins cut through the rock carrying associated minor amounts of pyrite and possibly bornite.	3	29	1	<.3
97-PR-17	KOFFEE - creek zone	117+35N	118+15W	Grab sample from outcrop by creek after excavation. Previous samples are 93-K-9, 97-PR-8.	Rusty weathering, Hb-Bi-Granodiorite, weak sericitic alteration of feldspars, possible secondary biotite alteration, 1mm pyrite filled fractures with 3-5 mm wide bleached envelopes. Trace of bornite.	4	36	2	<.3
97-PR-18	KOFFEE - creek zone	117+35N	118+00W	Grab sample from outcrop after excavation.	Bleached, clay-altered feldspars in coarse-grained Granodiorite(?). Alteration not texture destructive. Biotite altered to mauve-colored muscovite? Hair-line fractures contain pyrite.	117	16	1	<.3
97-PR-19	KOFFEE - creek zone	117+35N	117+85W	Grab sample from outcrop after excavation.	Propylitic altered Hb-Bi-Granodiorite, Hornblende altered to chlorite, biotite unaltered, Feldspars have slight green color, but still hard. Pyrite in hair-line fractures and after hornblende.	6	62	5	<.4
97-PR-20	KOFFEE - upper east road	116+90N	102+00W	Grab sample from ditch, not clear if s/c or float.	Rusty-weathering, angular pieces of fine-grained intrusive, silicious, minor amount of pyrite filling discontinuous fractures.	153	17	2	<.5
97-PR-21	KOFFEE - Koffee saddle	99+75N	98+60W	Grab sample of s/c from ditch.	Rusty-weathering, silicious breccia, silica flooding of breccia fragments has destroyed texture, disseminated pyrite in matrix, some angular spaces are open.	6	4	4	<.6
97-PR-22	KOFFEE - Koffee saddle	99+70N	98+35W	Grab sample of s/c from ditch.	Rusty-weathering, limonitic "rind" on boulders break open to expose a pervasive propylitically-altered coarse-grained rock (Granodiorite?) with a trace of pyrite.	2	5	1	<.7
97-PR-23	KOFFEE - Koffee saddle	98+85N	95+60W	Grab sample of float in ditch, many similar pieces at this location.	A crowded k-spar porphyry breccia with prominent k-spar and epidote veining. Matrix is medium to dark green and andesitic in composition. Breccia fragments display some chlorite alteration of biotite which define moderate foliation.	1	4	1	<.8
97-PR-24	KOFFEE - Koffee saddle	98+55N	94+30W	Grab sample of float in ditch.	A quartz-feldspar porphyry breccia with coarse-grained, chlorite altered biotite laths. Matrix is medium green, fine-grained, medium-soft (andesitic composition) with minor disseminated pyrite.	3	46	2	<.9
97-PR-25	KOFFEE - Koffee saddle	99+60N	98+00W	Grab sample of road material, large amount at this location suggests that it is subcrop.	Rusty-weathering silicious breccia with white-colored, hard fragments in dark grey fine-grained rhyodacitic matrix. Matrix carries fine-grained disseminated pyrite. Some fragments are of "matrix" with fragments, suggesting multiple stage brecciation.	3	33	6	1.2

SAMPLE NUMBER	LOCATION			TYPE/DEPTH	LITHOLOGICAL DESCRIPTION	SAMPLE ANALYSIS			
	CLAIM GROUP/AREA	NORTH	WEST			Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)
SAMPLE NUMBER	LOCATION			TYPE/DEPTH	LITHOLOGICAL DESCRIPTION	SAMPLE ANALYSIS			
	CLAIM GROUP/AREA	NORTH	WEST			Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)
97-PR-26	KOFFEE - Koffee saddle	100+15N	100+00W	Grab sample of float from ditch.	Yellowish limonitic weathering clay altered rock shot through with irregular quartz veining and including grey-coloured siliceous breccia fragments.	41	3	2	<.3
97-PR-27	KOFFEE - Koffee saddle	99+90N	98+90W	Grab sample of float from ditch.	Dark brown hematite weathering clay altered rock. Texture is largely destroyed but a vague impression of breccia persists.	2	12	11	0.3
97-PR-28	KOFFEE - Koffee saddle	99+80N	98+65W	Grab sample of float from ditch.	Bleached, argillic-altered breccia with prominent clay-altered feldspars in fragments and a bleached chlorite matrix. Grey silicious fractures cut through matrix and fragments.	1	3	1	<.3
97-PR-29	KOFFEE - Koffee saddle	100+10N	100+55W	Grab sample of float from ditch.	Medium-grained, propylitically altered hb-bi-granodiorite shot through with epidote veining and containing several short pyrite stringers.	1	5	1	<.3
97-PR-30	KOFFEE - Koffee saddle	100+00N	100+70W	Grab sample of float from ditch.	Rusty-weathering, bleached intrusive rock with grey silicified patches. Fine-grained, disseminated pyrite is associated with the silicification.	2	5	<1	<.3
97-PR-31	KOFFEE - Upper east Koffee road	114+50N	102+00W	Grab sample of subcrop from ditch.	Limonitic, bleached, clay-altered coarse-grained intrusive with silicious zones and weathered-out, fine-grained sulphide box-work. Rusty coated fractures.	26	44	2	<.3
97-PR-32	KOFFEE - Upper east Koffee road.	114+90N	102+00W	Grab sample of float from roadbed.	Limonitic, bleached, clay-altered coarse-grained intrusive with grey silicious patches.	210	68	5	0.4
97-PR-33	KOFFEE - Upper east Koffee road.	119+20N	105+50W	Grab sample of road material believed to subcrop from ditch.	Limonitic, bleached, clay-altered coarse-grained intrusive with grey silicious patches.	33	69	11	<.3
97-PR-34	KOFFEE - Upper east Koffee road.	119+10N	105+40W	Grab sample of road material.	Breccia with bleached, clay-altered, medium-grained intrusive fragments in a slightly more silicious, finer-grained, grey-colored matrix containing minor blebs of pyrite.	26	143	9	0.3
97-PR-35	KOFFEE - creek zone	116+25N	114+00W	Grab sample of float/subcrop from excavation in stream bank	Rusty-weathering, angular pieces of bleached, coarse-grained hb-bi-granodiorite. Hornblende is altered to biotite and chlorite. Fracture controlled pyrite is common. Trace of chalcopyrite. Pyrite has minor iridescence.	14	218	2	<.3
97-PR-36	KOFFEE - creek zone	117+20N	119+00W	Grab sample of float/subcrop from west side of excavation in stream bank.	Rusty-weathering, clay-altered, silicious, grey-colored intrusive shot through with pyrite veining and limonitic zones. small pockets and layers of wad.	722	3213	3	17.1
97-PR-37	KOFFEE - creek zone	117+10N	111+90W	Grab sample of float/subcrop from excavation in stream bank.	Fine-grained aplite (potassic flooding?) with 0.5% disseminated and fracture controlled pyrite.	8	18	2	<.3
97-PR-38	KOFFEE - creek zone	117+10N	111+90W	Grab sample of float/subcrop from excavation in stream bank.	Rusty-weathering, clay-altered porphyry with silicious grey-colored patches. Pyrite replaces mafics.	16	64	3	<.3
97-PR-39	KOFFEE - creek zone	117+00N	115+60W	Grab sample of outcrop from excavation in stream bank.	Propylitically-altered hb-bi-granodiorite, some biotite alteration of hornblende before chlorite, minor sericitic-alteration of feldspars. Pyrite-filled 1mm fractures may have k-spar envelopes. Pyrite has iridescent tarnish. Trace of chalcopyrite?	17	46	1	<.3

SAMPLE NUMBER	LOCATION			TYPE/DEPTH	LITHOLOGICAL DESCRIPTION	SAMPLE ANALYSIS			
	CLAIM GROUP/AREA	NORTH	WEST			Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)
SAMPLE NUMBER	LOCATION			TYPE/DEPTH	LITHOLOGICAL DESCRIPTION	SAMPLE ANALYSIS			
	CLAIM GROUP/AREA	NORTH	WEST			Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)
97-PR-40	KOFFEE - Koffee saddle	100+10N	100+06W	Grab sample of float from road bed.	Rusty weathering clay altered breccia with a limonitic and slightly siliceous matrix.	9	4	1	<.3
97-PR-41	KOFFEE - north Koffee road	119+40N	106+00W	Grab sample of float from road bed.	Yellowish-brown weathering, weakly clay-altered, medium-grained intrusive with 1cm wide, grey-colored silicification envelopes around hairline fractures with a rusty boxwork.	26	162	31	<.3
97-PR-42	KOFFEE - north Koffee road	119+50N	106+10N	Grab sample from ditch.	Rusty weathering moderately bleached and clay altered medium grained intrusive with 1cm wide silicification envelopes around 0.5 mm fractures containing pyrite and a open boxwork. Silicification more pervasive than in 97-PR-41.	9	178	10	<.3
97-PR-43	KOFFEE - north Koffee road.	120+00N	107+10W	Grab sample of float from road bed.	Slightly rusty-weathering, potassic-altered, medium-grained intrusive with disseminated pyrite and small fractures filled with pyrite.	17	519	23	<.3
97-PR-44	KOFFEE - north Koffee road.	120+40N	108+00W	Grab sample from float in road bed / subcrop in ditch.	Limonitic, bleached and weakly clay-altered medium-grained intrusive with 0.5 - 1mm wide grey-colored veins filled with black fine-grained mineral with an earthy texture (wad?).	20	105	8	<.3
97-PR-45	ANA - Access road to PD-3	113+30N	73+70W	Grab sample from road cut.	Sericitic-altered fine-grained intrusive with fine-grained disseminated pyrite.	33	78	<1	<.3
97-PR-46	ANA - Access road to PD-1	~115+50N	~70+50W	Grab sample from road cut.	Rusty-weathering, sericitic-altered, fine-grained intrusive with quartz veining.	58	64	11	<.3
A	KOFFEE - creek zone				Field notes lost	4	16	5	<.3
B	KOFFEE - creek zone				Field notes lost	52	19	3	<.4
C	KOFFEE - creek zone				Field notes lost	11	119	1	<.5
D	KOFFEE - creek zone				Field notes lost	23	601	1	.4
E	KOFFEE - creek zone				Field notes lost	34	49	14	<.3
F	KOFFEE - creek zone				Field notes lost	13	81	3	<.3
G	KOFFEE - creek zone				Field notes lost	941	84	8	<.3
H	KOFFEE - creek zone				Field notes lost	794	92	2	0.4
1	KOFFEE - creek zone				Field notes lost	41	90	3	<.3

APPENDIX 2 CANADIAN CREEK PROJECT 1999
TRENCH SAMPLES

TRENCH NUMBER	LOCATION			SIZE W x L x D	SAMPLE NUMBER	SAMPLE COMMENTS TYPE/DEPTH/LITHOLOGY	SAMPLE ANALYSIS							
	AREA	NORTH	WEST				Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)	As (ppm)	Zn (ppm)	Sb (ppm)	Pb (ppm)
97-T1	KOFFE E	117+50 N	116+00 W	3 x 40 x 3	97-T1-1	Grab sample of subcrop? from 3 metres depth from south end of trench. Pyrite-bornite stockwork is hosted by bleached and weakly clay-alteration overprint on a propylitically-altered, coarse-grained hb-bi-granodiorite.	57	84	2	<.3	2	20	<.3	8
					97-T1-2	Grab sample of float/subcrop? from 2 metres depth in center of trench. Pyrite rind on angular cobble-sized, propylitically-altered granodiorite. Trace of chalcoppyrite.	25	150	1	0.4	2	71	<.3	18
97-T2	KOFFE E	116+80 N	101+63 W	3 x 5 x 2	97-T2-1	Bleached, clay-altered and phyllic-altered angular float in a limonitic soil. Sample from 2 metres depth.	29	22	2	<.3	30	29	<.3	8
					97-T2-2	Silicious float with braided rusty fractures contain a trace of chalcoppyrite. Sample from 2 metres depth.	38	29	2	<.3	37	44	<.3	10
97-T3	KOFFE E	116+85 N	101+45 W	3 x 5 x 2	97-T3-S1	Soil sample of bright orangish-yellow limonitic soil from 2 metres depth.	43	54	3	<.3	225	48	<.3	10
					97-T3-2	Phyllic-altered, foliated float (YMC?) with 5% pyrite. From excavation pile, thought to be from about 2 metres depth.	33	20	1	<.3	59	34	<.3	7
97-T4	KOFFE E	117+50 N	102+15 W	3 x 5 x 2	97-T4-1	Rusty-weathering, silicious (veining and flooding) float with 1-5% pyrite and traces of bornite and chalcoppyrite.	59	40	8	<.3	26	26	<.3	11
97-T5	KOFFE E	116+65 W	101+90 W	3 x 10 x 6	97-T5-1	Grab sample of brown limonitic clay from 2 metres depth.	126	33	2	<.3	405	95	5	10
					97-T5-2	Pervasive phyllic/supergene clay alteration leaving clay, chlorite streaks and up to 10% pyrite. From 3 metres depth.	185	35	4	1.5	290	561	<.3	75
					97-T5-3	Dark-brown clay alteration from 4 metres depth	690	108	4	0.3	842	145	15	14
					97-T5-4	Silicious, phyllic-altered angular float from 6 metres depth.	388	39	1	<.3	97	21	<.3	9
					97-T5-5	Bleached, grey-colored clay with minor pyrite.	354	81	2	<.3	57	12	<.3	22

APPENDIX 2 CANADIAN CREEK PROJECT 1999
TRENCH SAMPLES

TRENCH	LOCATION			SIZE	SAMPLE	SAMPLE COMMENTS	SAMPLE ANALYSIS							
NUMBER	AREA	NORTH	WEST	W x L x D	NUMBER	TYPE/DEPTH/LITHOLOGY	Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)	As (ppm)	Zn (ppm)	Sb (ppm)	Pb (ppm)
TRENCH	LOCATION			SIZE	SAMPLE	SAMPLE COMMENTS	SAMPLE ANALYSIS							
NUMBER	AREA	NORTH	WEST	W x L x D	NUMBER	TYPE/DEPTH/LITHOLOGY	Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)	As (ppm)	Zn (ppm)	Sb (ppm)	Pb (ppm)
97-T6	KOFFE E	117+00 N	116+25 W	3 x 40 x 6	97-T6-1	Grab sample of layer of dark brown rusty, angular bits of propylitically-altered granodiorite from approximately 3 metres depth toward the southwest end of the trench.	21	170	3	<.3	4	29	<3	13
					97-T6-2	Grab sample of layer of dark brown rusty ferrocrite with both rounded and angular clasts of granodiorite from approximately 3 metres depth near the centre of the trench.	29	89	14	0.8	126	30	<3	10
					97-T6-3	Grab sample of "blue clay". A chlorite rich kaolinite with up to 5% pyrite from 6 metres depth in centre part of trench.	13	114	22	<.3	5	9	0.3	8
					97-T6-4	Grab sample of propylitically altered granodiorite float with a pyrite stockwork from 4 metres depth at the northeast end of the trench. This is approximately 10 metres from the south end of trench 97-T1, where stockwork mineralization was found in s/c.	125	112	2	<.3	4	29	<3	13
97-T7	KOFFE E	121+85 N	107+10 W	2 x 6 x 2	97-T7-1	Angular, silicious, phyllic-altered outcrop from 2 metres depth. approximately 2-4% pyrite.	39	9	8	0.5	13	3	<3	68
97-T8	KOFFE E	117+20 N	116+10 W	2 x 10 x 2	No Sample									
97-T9	KOFFE E	123+50 N	106+00 W	6 x 25 x 1	97-T9-1	Grab sample of phyllic-altered porphyry float and rusty soil. Subcrop/outcrop not reached because of permafrost.	45	22	2	0.5	224	122	3	102
97-T10	KOFFE E	125+50 N	110+00 W	3 x 10 x 1	97-T10-1	Grab sample of aplite/felsite float, of which there was an unusual amount in this trench. Subcrop/outcrop not reached because of permafrost.	3	12	4	<.3	67	25	4	15
97-T11	KOFFE E	118+00 N	122+00 W	4 x 5 x 0.3	97-T11-1	Aplite float in limonitic soil.	15	22	4	<.3	8	29	<3	11
97-T12	KOFFE E	118+35 N	121+30 W	4 x 5 x 0.15	97-T12-1	Aplite float.	12	23	3	<.3	8	31	<3	5
97-T13	KOFFE E	118+85 N	120+40 W	4 x 5 x 0.3	97-T13-1	Angular aplite float.	19	30	3	<.3	12	31	<3	9
97-T14	KOFFE E	119+35 N	119+50 W	4 x 5 x 0.15	97-T14-1	Angular aplite float and clay altered granodiorite float.	18	45	4	1.7	13	383	14	102

APPENDIX 2 CANADIAN CREEK PROJECT 1999
TRENCH SAMPLES

TRENCH NUMBER	LOCATION			SIZE W x L x D	SAMPLE NUMBER	SAMPLE COMMENTS TYPE/DEPTH/LITHOLOGY	SAMPLE ANALYSIS							
	AREA	NORTH	WEST				Au (ppb)	Cu (ppm)	Mo (ppm)	Ag (ppm)	As (ppm)	Zn (ppm)	Sb (ppm)	Pb (ppm)
TRENCH	LOCATION			SIZE	SAMPLE	SAMPLE COMMENTS	SAMPLE ANALYSIS							
NUMBER	AREA	NORTH	WEST	W x L x D	NUMBER	TYPE/DEPTH/LITHOLOGY	Au	Cu	Mo	Ag	As	Zn	Sb	Pb
97-T15	KOFFE E	119+75 N	118+65 W	4 x 5 x 2	97-T15-1	Aplite float in creek bed.	12	24	3	<.3	49	15	<3	8
					97-T15-2	Granodiorite with disseminated pyrite in creek bed.	8	57	4	.3	10	30	<3	12
97-T16	KOFFE E	121+00 N	118+00 W	4 x 5 x 0.5	97-T16-1	Clay-altered granodiorite float in yellowish-brown soil.	24	34	16	<.3	20	19	<3	10
					97-T16-2	Clay-altered granodiorite float with rusty pin heads (after pyrite?) in yellowish-brown soil.	72	6	1	<.3	12	2	<3	3
97-T17	KOFFE E	120+50 N	118+00 W	4 x 5 x 0.5	97-T17-1	Bleached, rusty granodiorite float.	12	21	2	<.3	13	28	<3	5
97-T18	KOFFE E	120+30 N	117+60 W	4 x 5 x 2	97-T18-1	Clay-altered granodiorite float.	9	31	5	<.3	14	31	<3	12
					97-T18-2	Rusty quartz float with vughs and disseminated crystalline pyrite.	394	19	3	6.9	29	71	26	140
97-T19	KOFFE E	120+00 N	118+00 W	4 x 5 x 0.5	97-T19-1	Clay-altered, fine-grained, rusty intrusive float.	12	25	3	<.3	33	51	<3	15
97-T20	ANA	112+70 N	82+25W	2 x 6 x 3	97-T20-1	Fine-grained intrusive float with a trace of pyrite. Located approximately 50 metres from trench 96-28.	130	381	7	0.7	27	25	<3	4
97-T21	ANA	113+25 N	81+95W	2 x 6 x 3	97-T21-1	Fine-grained intrusive float with quartz vein containing moly rossette. Located approximately 10 metres from trench 96-28.	44	40	21	<.3	2	11	<3	7
					97-T21-2	Micro-breccia float. Located approximately 10 metres from trench 96-28.	98	252	48	0.8	25	56	<3	14
97-T22	ANA	113+60 N	81+75W	2 x 6 x 3	No sample	No Micro-breccia float found. Located approximately 50 metres from trench 96-28.								



GEOCHEMICAL ANALYSIS CERTIFICATE

Mincord Exploration Consultants Ltd. PROJECT CELIN CREEK File # 9904829 Page 1

110 - 325 Howe St., Vancouver BC V6C 1Z7 Submitted by: Bill Morton



SAMPLE#	Mo ppm	Cu ppm	Pb ppm	Zn ppm	Ag ppm	Ni ppm	Co ppm	Mn ppm	Fe %	As ppm	U ppm	Au ppm	Th ppm	Sr ppm	Cd ppm	Sb ppm	Bi ppm	V ppm	Ca %	P %	La ppm	Cr ppm	Mg %	Ba ppm	Ti %	B ppm	Al %	Na %	K %	W ppm	Au* ppb
97-PR-1	7	80	13	29	<.3	15	2	44	.93	9	<8	<2	4	7	<.2	<3	<3	22	.02	.009	14	26	.26	355	<.01	<3	.51	.01	.23	6	31.5
97-PR-2	2	4	10	9	.3	2	1	37	.65	23	<8	<2	3	2	<.2	<3	<3	1	.02	.002	7	14	.02	66	<.01	18	.26	.01	.24	3	35.4
97-PR-3	2	40	5	14	<.3	2	2	102	2.22	38	<8	<2	15	24	<.2	<3	<3	7	.06	.017	12	7	.11	634	<.01	<3	.82	.01	.32	2	45.5
97-PR-4	5	10	19	107	<.3	2	1	135	.60	98	<8	<2	14	5	.7	<3	<3	1	.05	.006	22	10	.03	162	<.01	<3	.33	.05	.16	4	4.1
97-PR-5	2	11	15	50	<.3	3	1	205	.51	40	<8	<2	14	8	2.2	<3	<3	1	.03	.003	12	14	.01	501	<.01	<3	.24	.04	.13	4	7.1
97-PS-6	4	29	20	25	.8	3	1	44	3.03	13	<8	<2	3	9	<.2	<3	<3	27	.02	.026	14	11	.16	186	<.01	<3	.86	.01	.21	2	34.5
97-PR-7	3	11	8	6	<.3	4	5	35	1.72	109	<8	<2	6	10	<.2	<3	<3	9	.01	.010	17	12	.01	135	<.01	7	.37	.01	.27	3	144.1
97-PR-8	3	105	<3	27	<.3	5	11	339	3.48	3	<8	<2	11	25	<.2	<3	<3	99	.84	.054	19	25	1.09	267	.19	<3	1.79	.10	.81	2	7.0
97-PR-9	2	26	7	10	<.3	4	2	83	.67	3	<8	<2	31	10	<.2	<3	<3	6	.23	.003	8	14	.10	72	<.01	<3	.24	.04	.11	5	2.9
97-PR-10	58	69	9	63	.3	57	8	230	2.25	11	<8	<2	9	44	.5	5	<3	29	.23	.071	17	58	.26	222	.04	<3	.40	.03	.22	11	6.6
97-PR-11	3	45	7	44	<.3	18	5	103	1.70	9	<8	<2	4	8	.2	<3	<3	92	.01	.010	11	65	.69	1870	.11	<3	.83	.03	.58	8	2.8
97-PR-12	6	20	8	12	<.3	5	2	132	1.55	2	<8	<2	4	27	<.2	23	<3	108	.01	.021	14	70	.51	2438	.04	<3	.82	.04	.45	4	9.2
97-PR-13	2	108	7	43	<.3	73	25	138	1.95	13	<8	<2	2	175	.2	<3	<3	49	3.77	.294	20	63	.32	247	.27	<3	4.67	.55	.17	42	8.4
97-PR-14	1	13	11	88	<.3	35	16	492	3.99	29	<8	<2	13	46	<.2	<3	<3	40	.51	.020	37	50	1.02	179	.02	3	2.22	.03	.35	2	1.0
97-PR-15	2	40	5	82	<.3	36	15	367	4.38	7	<8	<2	7	9	<.2	<3	<3	88	.15	.045	19	82	1.23	88	.36	<3	2.41	.05	1.30	6	12.7
RE 97-PR-15	2	41	4	83	<.3	36	15	368	4.42	5	<8	<2	8	9	<.2	<3	<3	89	.15	.045	20	80	1.24	87	.36	<3	2.43	.05	1.30	6	12.5
97-PR-16	1	29	<3	15	<.3	4	6	202	1.88	2	<8	<2	15	21	<.2	<3	<3	45	.67	.028	14	16	.55	174	.06	<3	.91	.08	.32	2	2.6
97-PR-17	2	36	4	14	<.3	5	12	255	3.39	3	<8	<2	10	29	<.2	<3	<3	51	1.50	.049	20	15	.85	80	.07	3	1.19	.07	.49	3	3.8
97-PR-18	1	16	4	2	<.3	3	7	29	1.50	116	<8	<2	2	3	<.2	4	<3	10	.02	.004	23	8	.02	61	<.01	10	.46	<.01	.17	2	117.3
97-PR-19	5	62	<3	23	<.3	5	11	309	3.32	2	<8	<2	11	25	<.2	<3	<3	98	.73	.051	18	22	1.18	279	.22	<3	1.79	.11	.87	2	5.8
97-PR-20	2	17	26	14	<.3	4	1	150	1.01	57	<8	<2	6	10	<.2	5	<3	5	.02	.011	14	12	.05	262	<.01	<3	.41	.01	.28	3	152.8
97-PR-21	4	4	12	57	<.3	5	9	114	1.84	33	12	<2	7	29	5.1	6	<3	14	.04	.020	14	14	.02	121	<.01	<3	.55	<.01	.14	3	5.6
97-PR-22	1	5	9	52	<.3	4	10	723	2.97	4	<8	<2	9	64	<.2	<3	<3	34	4.16	.038	13	13	.41	97	<.01	<3	.54	<.01	.22	2	2.2
97-PR-23	1	4	9	57	<.3	6	11	702	2.77	3	<8	<2	23	68	<.2	<3	3	63	.88	.050	49	21	1.17	106	.15	<3	1.86	.06	.12	2	1.2
97-PR-24	2	46	7	70	<.3	11	20	855	4.86	6	<8	<2	14	49	<.2	<3	<3	117	.94	.099	23	26	1.57	458	.24	<3	2.26	.08	.09	3	3.2
97-PR-25	6	33	298	605	1.2	10	9	86	1.61	289	19	<2	<2	9	12.1	38	<3	10	.03	.008	2	28	.02	41	<.01	3	.24	<.01	.06	5	2.7
97-PR-26	2	3	67	58	<.3	2	1	29	.48	79	<8	<2	2	15	.4	8	<3	5	.04	.003	6	8	.02	1287	<.01	3	.24	<.01	.15	4	40.9
97-PR-27	1	12	10	57	<.3	5	12	1058	3.53	11	<8	<2	17	10	<.2	<3	4	52	2.21	.041	29	17	.07	529	.01	<3	.79	.01	.27	2	2.1
97-PR-28	1	3	16	54	<.3	5	13	138	1.07	14	<8	<2	11	6	<.2	5	<3	27	.12	.043	29	8	.01	346	<.01	3	.66	<.01	.11	2	.8
97-PR-29	1	5	4	57	<.3	6	12	642	2.99	4	<8	<2	10	36	<.2	<3	<3	69	1.14	.036	17	18	1.23	164	.08	<3	2.00	.09	.10	3	.5
97-PR-30	<1	2	10	48	<.3	4	12	835	2.61	69	<8	<2	9	23	<.2	6	<3	20	1.49	.035	13	8	.40	211	<.01	3	.61	.01	.21	2	1.5
97-PR-31	2	44	9	12	<.3	3	3	107	1.70	7	<8	<2	18	41	<.2	<3	<3	4	.06	.024	25	14	.14	609	<.01	4	.71	.01	.42	2	26.1
97-PR-32	5	68	22	18	.4	4	4	95	3.36	122	<8	<2	11	17	<.2	<3	7	6	.04	.018	11	9	.06	110	<.01	<3	.55	.01	.26	2	210.2
97-PR-33	11	69	13	12	<.3	2	1	44	1.12	7	<8	<2	20	13	<.2	<3	<3	5	.06	.017	25	11	.07	399	<.01	<3	.57	.04	.29	2	32.9
97-PR-34	9	143	9	10	.3	3	3	43	1.25	8	<8	<2	12	10	<.2	<3	<3	3	.03	.005	16	10	.04	237	<.01	<3	.41	.01	.34	3	26.4
STANDARD DS2	14	128	30	168	<.3	39	13	837	3.35	61	21	<2	4	30	11.6	7	10	85	.56	.086	18	172	.64	152	.12	3	1.84	.04	.17	9	205.7

GROUP 1D - 0.50 GM SAMPLE LEACHED WITH 3 ML 2-2-2 HCL-HNO₃-H₂O AT 95 DEG. C FOR ONE HOUR, DILUTED TO 10 ML, ANALYSED BY ICP-ES.
UPPER LIMITS - AG, AU, HG, W = 100 PPM; MO, CO, CD, SB, BI, TH, U & B = 2,000 PPM; CU, PB, ZN, NI, MN, AS, V, LA, CR = 10,000 PPM.
ASSAY RECOMMENDED FOR ROCK AND CORE SAMPLES IF CU PB ZN AS > 1%, AG > 30 PPM & AU > 1000 PPB
- SAMPLE TYPE: ROCK AU* BY ACID LEACHED, ANALYZE BY ICP-MS. (10 gm)
Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.

DATE RECEIVED: DEC 16 1999 DATE REPORT MAILED: Jan 5/2000 SIGNED BY: C. Leong D. TOYE, C. LEONG, J. WANG; CERTIFIED B.C. ASSAYERS

All results are considered the confidential property of the client. Acme assumes the liabilities for actual cost of the analysis only.

Data FA



SAMPLE#	Mo ppm	Cu ppm	Pb ppm	Zn ppm	Ag ppm	Ni ppm	Co ppm	Mn ppm	Fe %	As ppm	U ppm	Au ppm	Th ppm	Sr ppm	Cd ppm	Sb ppm	Bi ppm	V ppm	Ca %	P %	La ppm	Cr ppm	Mg %	Ba ppm	Ti %	B ppm	Al %	Na %	K %	W ppm	Au* ppb
97-PR-35	2	218	6	27	<.3	6	15	262	3.52	3	<8	<2	9	24	.4	<3	<3	124	.38	.060	11	16	1.56	234	.17	<3	2.12	.11	.92	3	13.5
97-PR-36	3	3213	235	136	17.1	6	17	540	10.18	23	<8	<2	11	10	1.4	<3	54	29	.34	.047	13	10	.98	42	<.01	7	1.29	.01	.30	5	721.5
97-PR-37	2	18	6	4	<.3	3	1	43	.63	2	<8	<2	9	4	<.2	<3	<3	2	.02	.004	8	17	.03	65	<.01	<3	.23	.05	.16	4	8.1
97-PR-38	3	64	6	18	<.3	5	3	139	3.14	4	25	<2	15	18	.5	<3	<3	83	.08	.026	14	19	1.45	141	.20	<3	1.68	.06	.14	3	16.4
97-PR-39	1	46	3	30	<.3	6	14	352	4.25	<2	<8	<2	7	41	.8	<3	<3	92	1.31	.058	9	16	1.19	119	.12	<3	1.84	.12	.64	3	17.0
97-PR-40	1	4	<3	26	<.3	3	7	1666	3.78	29	<8	<2	4	111	.7	<3	<3	10	6.80	.021	5	4	1.57	174	<.01	3	.45	.01	.26	3	8.8
97-PR-41	31	162	19	15	<.3	3	2	58	1.11	5	<8	<2	22	23	<.2	<3	<3	5	.07	.015	23	10	.08	881	<.01	<3	.56	.01	.37	3	25.9
97-PR-42	10	178	87	28	<.3	3	1	84	1.30	11	<8	<2	23	15	<.2	17	<3	4	.07	.018	23	7	.08	238	<.01	3	.53	.01	.41	3	9.1
97-PR-43	23	519	12	16	<.3	4	6	62	1.24	4	<8	<2	23	10	<.2	<3	<3	6	.06	.013	24	13	.06	232	<.01	<3	.52	.03	.24	3	17.2
97-PR-44	8	105	9	15	<.3	3	1	42	1.58	2	<8	<2	18	29	<.2	<3	<3	8	.08	.010	21	6	.06	1176	<.01	<3	.56	.03	.20	2	20.4
97-PR-45	<1	78	14	17	<.3	2	3	54	1.59	16	<8	<2	10	61	<.2	<3	<3	4	.50	.033	9	5	.19	175	<.01	<3	1.83	.18	.15	<2	32.5
97-PR-46	11	64	<3	9	<.3	8	1	40	1.63	11	<8	<2	9	12	<.2	<3	<3	19	.02	.015	19	23	.15	182	.02	<3	.57	.02	.46	4	57.5
97-T1-1	2	84	8	20	<.3	4	13	282	4.37	2	9	<2	13	21	.7	<3	<3	22	2.00	.049	18	9	.56	64	<.01	<3	.89	.01	.26	2	57.3
97-T1-2	1	150	18	71	.4	5	11	398	3.52	2	<8	<2	12	24	.6	<3	<3	79	.31	.053	19	13	1.10	442	.11	<3	1.60	.08	.56	3	24.8
97-T2-1	2	22	8	29	<.3	4	3	120	1.76	30	<8	<2	13	17	<.2	<3	<3	3	.07	.010	26	13	.06	357	<.01	<3	.48	.01	.28	3	28.8
97-T2-2	2	29	7	44	<.3	5	3	145	1.82	37	<8	<2	9	21	<.2	<3	<3	2	.08	.011	22	7	.05	770	<.01	<3	.42	.01	.23	4	37.6
97-T3-1	3	54	10	48	<.3	4	1	90	2.66	225	<8	<2	31	25	.3	<3	<3	4	.18	.045	33	14	.09	175	<.01	<3	.64	.01	.24	3	43.4
97-T3-2	1	20	7	34	<.3	4	5	45	2.05	59	<8	<2	13	20	<.2	<3	3	1	.07	.012	8	10	.04	215	<.01	<3	.35	.01	.22	4	33.0
97-T4-1	8	40	11	26	<.3	4	6	224	3.32	26	<8	<2	12	33	.5	<3	<3	6	.04	.011	14	10	.13	279	<.01	3	.82	.01	.35	<2	58.6
97-T5-1	2	33	10	95	<.3	8	4	224	2.10	405	<8	<2	11	14	<.2	5	3	8	.18	.022	11	9	.11	197	<.01	<3	.70	.01	.16	4	126.1
97-T5-2	4	35	75	561	1.5	13	7	115	8.49	290	<8	<2	7	7	5.0	<3	10	5	.11	.023	7	10	.07	35	<.01	4	.56	.01	.29	<2	184.5
97-T5-3	4	108	14	145	.3	15	16	1585	3.23	842	<8	<2	19	11	.3	15	9	3	.10	.050	33	9	.05	208	<.01	3	.46	.01	.22	5	690.3
97-T5-4	1	39	9	21	<.3	6	7	69	2.88	97	<8	<2	4	6	<.2	<3	6	2	.11	.030	10	6	.03	88	<.01	<3	.50	.01	.27	2	388.1
97-T5-5	2	81	22	12	<.3	5	3	46	1.29	57	<8	<2	8	8	<.2	<3	3	2	.06	.005	17	7	.03	130	<.01	<3	.32	.01	.21	4	354.0
TR 97-T5-5	3	84	28	12	<.3	5	3	47	1.35	59	<8	<2	9	9	<.2	4	<3	1	.06	.005	17	9	.03	132	<.01	<3	.33	.01	.22	3	214.1
97-T6-1	3	170	3	14	<.3	5	12	68	3.93	14	<8	<2	14	18	.3	<3	3	42	.29	.062	18	9	.56	60	.01	4	1.55	.02	.42	2	21.1
97-T6-2	14	89	10	30	.8	6	6	217	6.64	126	<8	<2	26	24	<.2	<3	<3	60	.19	.084	12	15	.42	242	.04	<3	1.25	.04	.22	3	29.3
97-T6-3	22	114	8	9	<.3	5	15	198	3.86	5	<8	<2	8	27	.3	<3	<3	37	2.52	.057	13	8	.88	58	<.01	4	1.68	.01	.36	<2	12.6
97-T6-4	2	112	13	29	<.3	6	12	476	7.77	4	<8	<2	9	39	<.2	<3	<3	37	3.42	.058	13	8	.78	17	.01	5	1.20	.01	.30	2	125.1
97-T7-1	8	9	68	3	.5	2	1	28	1.28	13	<8	<2	12	9	<.2	<3	<3	3	.02	.010	16	7	.03	228	<.01	<3	.36	.02	.42	5	39.1
97-T9-1	2	22	102	122	.5	6	4	356	1.69	224	<8	<2	18	17	1.0	3	<3	19	.11	.032	17	15	.17	136	.01	<3	.65	.03	.25	5	44.8
97-T10-1	4	12	15	25	<.3	4	3	152	1.22	67	<8	<2	20	6	<.2	4	<3	8	.04	.012	22	14	.06	122	.01	<3	.39	.03	.22	2	3.0
97-T11-1	4	22	11	29	<.3	9	6	226	2.30	8	<8	<2	7	34	<.2	<3	<3	54	.27	.037	12	27	.42	232	.07	<3	1.10	.06	.19	4	14.9
97-T12-1	3	23	5	31	<.3	10	5	191	2.45	8	<8	<2	9	28	<.2	<3	<3	59	.26	.037	14	27	.51	263	.09	4	1.32	.05	.23	4	11.7
97-T13-1	3	30	9	31	<.3	10	4	167	2.26	12	<8	<2	6	29	<.2	<3	<3	49	.27	.031	15	26	.45	303	.06	4	1.40	.03	.19	4	19.0
STANDARD DS2	15	131	28	170	<.3	41	14	817	3.49	62	21	<2	4	32	12.3	8	12	91	.61	.086	17	187	.67	156	.10	4	1.99	.04	.18	10	204.9

Sample type: ROCK.



Mincord Exploration Consultants Ltd. PROJECT CELIN CREEK FILE # 9904829 Page 3



SAMPLE#	Mo ppm	Cu ppm	Pb ppm	Zn ppm	Ag ppm	Ni ppm	Co ppm	Mn ppm	Fe %	As ppm	U ppm	Au ppm	Th ppm	Sr ppm	Cd ppm	Sb ppm	Bi ppm	V ppm	Ca %	P %	La ppm	Cr ppm	Mg %	Ba ppm	Ti %	B ppm	Al %	Na %	K %	W ppm	Au* ppb
97-T14-1	4	45	102	383	1.7	9	4	157	1.65	13	<8	<2	7	24	1.1	14	<3	38	.22	.030	15	23	.37	384	.07	3	1.16	.04	.20	3	18.3
97-T15-1	3	24	8	15	<.3	4	3	83	1.72	49	<8	<2	16	25	<.2	<3	<3	14	.05	.016	14	10	.12	1590	.01	4	.59	.02	.26	6	11.5
97-T15-2	4	57	12	30	<.3	7	6	181	2.32	10	<8	<2	15	23	<.2	<3	<3	56	.14	.039	16	20	.65	182	.10	3	1.25	.07	.41	6	7.7
97-T16-1	16	34	10	19	<.3	5	3	136	2.64	20	<8	<2	13	34	<.2	<3	<3	38	.11	.038	19	12	.39	266	.03	3	1.12	.05	.29	2	23.6
97-T16-2	1	6	3	2	<.3	1	1	25	.83	12	<8	<2	5	13	<.2	<3	<3	6	.01	.005	20	6	.03	621	<.01	4	.48	.01	.29	2	71.9
97-T17-1	2	21	5	28	<.3	7	4	181	1.98	13	<8	<2	9	16	<.2	<3	<3	35	.16	.029	18	16	.34	200	.07	3	1.10	.03	.27	6	12.3
97-T18-1	5	31	12	31	<.3	5	7	249	2.64	14	<8	<2	13	31	.2	<3	<3	59	.22	.036	14	18	.71	401	.14	5	1.44	.07	.42	2	8.6
97-T18-2	3	19	140	71	6.9	6	2	83	1.90	29	<8	<2	3	10	.6	26	3	14	.05	.011	3	16	.09	282	.03	4	.29	.01	.06	9	393.6
97-T19-1	3	25	15	51	<.3	15	8	288	2.58	33	<8	<2	9	25	<.2	<3	<3	62	.33	.045	17	36	.44	240	.11	4	1.58	.04	.12	4	11.9
97-T20-1	7	381	4	25	.7	11	31	867	4.05	27	<8	<2	8	12	<.2	<3	<3	37	.19	.072	20	19	.22	273	.01	8	.66	.01	.27	4	129.6
RE 97-T20-1	7	395	8	25	.6	12	32	894	4.21	27	<8	<2	8	13	<.2	<3	3	38	.20	.073	20	18	.22	303	.01	7	.68	.01	.28	4	140.5
97-T21-1	21	40	7	11	<.3	3	3	84	.73	2	<8	<2	22	5	<.2	<3	<3	5	.06	.003	7	15	.08	52	<.01	<3	.37	.05	.14	4	43.5
97-T21-2	48	252	14	56	.8	7	11	252	4.50	25	<8	<2	8	14	<.2	<3	3	41	.24	.061	36	24	.42	216	.01	15	1.21	.02	.23	4	98.3
99-A	5	16	11	48	<.3	10	10	452	1.80	36	<8	<2	11	13	<.2	<3	<3	28	.16	.030	18	26	.36	120	.09	3	.91	.05	.38	6	4.2
99-B	3	19	7	4	<.3	2	1	37	1.14	341	<8	<2	3	13	<.2	10	<3	10	.02	.007	16	11	.06	243	<.01	4	.45	.01	.28	3	51.7
99-C	1	119	5	22	<.3	4	8	141	4.93	4	<8	<2	10	23	<.2	<3	<3	67	.21	.067	13	12	.76	249	.07	4	2.09	.02	.43	2	11.2
99-D	1	601	9	29	.4	7	14	369	4.76	10	<8	<2	7	18	<.2	<3	<3	61	1.26	.062	14	15	.78	54	.15	5	1.08	.05	.58	5	23.4
99-E	14	49	15	54	<.3	5	7	239	2.84	7	<8	<2	15	24	<.2	<3	<3	40	.12	.025	18	10	.39	639	.04	3	1.04	.03	.34	3	34.2
99-F	3	81	10	21	<.3	5	10	311	6.14	15	<8	<2	17	18	<.2	<3	<3	41	1.24	.068	18	6	.65	72	.03	6	1.83	.01	.36	2	13.3
99-G	8	84	7	12	<.3	5	18	267	8.42	<2	<8	<2	7	25	<.2	<3	<3	13	2.75	.042	11	7	.35	50	<.01	4	.82	.01	.29	3	941.3
99-H	2	92	5	15	.4	5	14	240	5.89	3	<8	<2	6	19	<.2	<3	<3	15	1.42	.042	8	7	.42	48	<.01	<3	.80	.01	.28	2	795.9
99-I	3	90	15	38	.3	6	17	443	5.31	13	<8	<2	8	22	<.2	<3	<3	37	1.86	.060	12	15	.44	33	<.01	4	.61	.01	.23	3	41.8
STANDARD DS2	14	132	34	170	<.3	39	13	804	3.43	62	21	<2	4	30	11.7	8	11	86	.58	.089	18	181	.65	153	.12	6	1.87	.04	.17	10	205.9

Sample type: ROCK. Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.



GEOCHEMICAL ANALYSIS CERTIFICATE



Mincord Exploration Consultants Ltd. PROJECT CELIN CREEK File # 9904828 Page 1

110 - 325 Howe St., Vancouver BC V6C 1Z7 Submitted by: Bill Morton

SAMPLE#	Mo ppm	Cu ppm	Pb ppm	Zn ppm	Ag ppm	Ni ppm	Co ppm	Mn ppm	Fe %	As ppm	U ppm	Au ppm	Th ppm	Sr ppm	Cd ppm	Sb ppm	Bi ppm	V ppm	Ca %	P %	La ppm	Cr ppm	Mg %	Ba ppm	Ti %	B ppm	Al %	Na %	K %	W ppm	Au* ppb
KSS-1	1	34	48	120	<.3	23	16	535	3.56	38	<8	<2	7	28	1.6	<3	<3	89	.52	.092	23	41	.74	228	.13	4	1.82	.02	.08	2	3.9
KSS-2	1	49	26	104	<.3	19	21	1081	3.97	35	<8	<2	13	32	1.5	<3	<3	112	.61	.106	31	36	.84	269	.19	<3	2.38	.02	.24	2	2.8
KSS-3	1	77	26	112	<.3	22	17	627	3.95	45	<8	<2	10	30	1.5	<3	<3	115	.52	.087	25	42	.85	230	.19	<3	2.22	.02	.24	4	2.7
KSS-4	1	117	32	163	.4	43	25	821	4.98	53	<8	<2	7	39	1.7	<3	<3	141	.66	.088	22	83	1.21	263	.23	3	2.75	.02	.31	4	4.4
KSS-5	2	74	12	75	<.3	29	15	438	3.33	30	<8	<2	4	32	.8	<3	<3	94	.62	.080	13	55	.77	141	.13	3	1.89	.03	.07	3	2.9
KSS-6	3	249	49	302	.6	41	40	2077	4.90	99	<8	<2	8	36	2.3	5	<3	112	.69	.074	24	56	.92	274	.09	3	2.42	.02	.08	<2	22.6
KSS-7	3	23	82	80	.4	5	4	456	5.89	97	<8	<2	18	169	1.6	9	<3	51	.23	.083	46	14	.06	106	<.01	4	.50	.02	1.00	3	3.0
KSS-8	3	13	27	127	.3	8	20	1182	4.70	12	<8	<2	15	13	1.1	3	3	63	.35	.083	31	16	.17	223	.01	3	.74	.01	.11	3	10.4
KSS-9	<1	16	14	75	<.3	6	14	605	4.04	184	<8	<2	12	21	.8	21	<3	35	.20	.043	37	11	.12	202	<.01	5	.48	.01	.20	2	4.8
KSS-10	1	8	133	724	<.3	4	7	1002	3.84	483	<8	<2	16	31	6.7	20	<3	45	.28	.046	24	10	.09	393	<.01	4	.54	.01	.17	<2	11.2
KSS-11	2	14	313	787	1.8	10	12	1010	5.65	587	<8	<2	11	14	3.8	41	3	33	.36	.043	14	17	.22	254	.02	3	.85	.01	.12	<2	93.5
KSS-12	1	13	13	86	<.3	6	15	839	3.87	208	<8	<2	19	16	.7	11	<3	96	.72	.105	40	15	.34	315	.01	<3	1.45	.01	.17	<2	7.5
KSS-13	1	16	9	73	<.3	9	15	962	4.42	49	<8	<2	21	26	.9	13	<3	68	1.04	.099	32	18	.80	229	.02	<3	2.38	.01	.21	2	5.3
KSS-14	<1	27	15	84	<.3	19	13	781	3.61	26	<8	<2	10	26	.7	4	<3	86	.64	.077	24	30	.75	221	.07	<3	1.91	.02	.14	2	9.7
KSS-15	<1	31	13	82	<.3	22	15	1189	3.80	21	<8	<2	12	32	.8	<3	<3	95	.70	.087	33	32	.94	254	.09	<3	2.24	.02	.15	3	15.8
KSS-16	4	27	9	127	<.3	23	16	1301	3.51	18	12	<2	6	29	1.0	<3	<3	82	.59	.096	21	42	.76	374	.08	3	2.17	.02	.07	3	4.5
KSS-17	4	12	14	68	<.3	10	8	634	2.60	6	<8	<2	19	20	.6	<3	<3	56	.48	.055	35	20	.64	201	.04	<3	1.70	.01	.08	2	1.9
RE KSS-17	4	12	15	69	<.3	10	8	658	2.69	8	<8	<2	20	21	.5	<3	<3	58	.50	.058	36	21	.66	206	.04	<3	1.73	.01	.08	2	.5
KSS-18	17	11	20	327	<.3	9	11	971	3.37	47	<8	<2	29	20	1.7	<3	<3	54	.43	.058	43	14	.36	217	.01	<3	1.31	.01	.13	<2	1.0
KSS-19	11	10	5	196	<.3	6	16	1892	3.99	35	<8	<2	33	20	1.6	<3	<3	100	.43	.066	44	16	1.11	545	.25	<3	2.46	.02	.94	8	1.5
KSS-20	27	14	34	141	<.3	6	6	635	2.72	126	11	<2	31	13	.6	<3	<3	20	.24	.030	38	7	.13	125	<.01	<3	.64	.01	.12	4	7.7
KSS-21	1	9	25	142	<.3	6	9	701	3.01	82	<8	<2	35	21	.5	<3	<3	66	.42	.065	50	14	.68	277	.13	<3	1.93	.02	.53	5	2.0
KSS-22	3	18	21	130	.3	6	9	1105	2.47	19	<8	<2	33	22	.2	<3	<3	66	.47	.063	44	13	.70	364	.12	<3	1.78	.02	.51	4	7.2
KSS-23	2	114	72	609	2.3	9	10	445	3.32	39	<8	<2	7	29	.7	<3	<3	101	.33	.053	19	23	.88	437	.20	<3	2.11	.03	.42	2	61.6
KSS-24	11	44	104	302	.4	20	12	1562	3.49	119	<8	<2	11	24	3.1	<3	<3	89	.46	.083	21	31	.72	363	.14	<3	1.56	.02	.28	3	23.0
KSS-25	5	31	42	222	<.3	15	14	1026	3.41	36	<8	<2	11	22	1.2	<3	<3	110	.55	.103	20	31	1.05	328	.22	<3	1.97	.02	.46	8	5.2
KSS-26	3	29	39	143	<.3	20	13	483	3.07	20	<8	<2	7	28	.5	<3	<3	81	.48	.083	21	37	.74	253	.13	<3	1.88	.02	.10	6	8.3
KSS-27	5	83	82	397	.8	15	12	755	4.44	183	<8	<2	8	26	1.1	3	3	93	.42	.089	20	25	.87	328	.17	3	2.10	.02	.33	2	13.8
KSS-28	1	45	9	27	<.3	6	4	343	2.80	5	<8	<2	16	27	<.2	<3	<3	67	.40	.062	33	6	.53	338	.03	<3	1.88	.01	.27	3	11.9
KSS-29	2	44	19	76	<.3	13	14	523	3.39	11	<8	<2	12	44	<.2	<3	3	81	.41	.075	26	26	.82	310	.12	<3	1.81	.02	.21	3	78.5
KSS-30	1	39	22	90	<.3	10	12	648	3.68	15	<8	<2	12	31	<.2	<3	<3	89	.43	.090	22	18	.76	257	.12	<3	1.58	.02	.19	3	172.4
KSS-31	1	52	25	101	<.3	12	11	583	3.93	19	<8	<2	12	26	<.2	<3	<3	93	.36	.090	20	22	.86	282	.15	<3	1.72	.02	.27	5	40.7
KSS-32	<1	29	6	71	<.3	12	11	576	3.17	16	<8	<2	8	26	<.2	<3	3	82	.42	.093	14	26	.69	198	.14	<3	1.36	.02	.14	5	104.6
KSS-33	1	35	16	105	.3	13	13	666	3.84	19	<8	<2	10	27	<.2	<3	3	91	.36	.076	17	25	.86	295	.13	<3	1.96	.02	.16	4	85.8
STANDARD DS2	15	136	32	166	<.3	38	13	833	3.32	62	23	<2	4	30	11.9	8	9	84	.56	.085	17	178	.63	150	.12	3	1.83	.04	.16	11	210.3

GROUP 1D - 0.50 GM SAMPLE LEACHED WITH 3 ML 2-2-2 HCL-HNO3-H2O AT 95 DEG. C FOR ONE HOUR, DILUTED TO 10 ML, ANALYSED BY ICP-ES.

UPPER LIMITS - AG, AU, HG, W = 100 PPM; MO, CO, CD, SB, BI, TH, U & B = 2,000 PPM; CU, PB, ZN, NI, MN, AS, V, LA, CR = 10,000 PPM.

- SAMPLE TYPE: SOIL AU* BY ACID LEACHED, ANALYZE BY ICP-MS. (10 gm)

Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.

DATE RECEIVED: DEC 16 1999

DATE REPORT MAILED: Jan 5/2000

SIGNED BY: C. LEONG, J. WANG; CERTIFIED B.C. ASSAYERS



SAMPLE#

Mo	Cu	Pb	Zn	Ag	Ni	Co	Mn	Fe	As	U	Au	Th	Sr	Cd	Sb	Bi	V	Ca	P	La	Cr	Mg	Ba	Ti	B	Al	Na	K	W	Au*
ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	%	ppm	ppm	%	ppm	%	%	%	%	%	ppm	ppb
1	29	8	75	<.3	10	13	873	3.41	16	<8	<2	9	28	<.2	3	<3	80	.37	.070	17	23	.75	297	.13	<3	1.56	.02	.14	4	26.6
1	30	13	64	<.3	8	11	554	3.50	21	<8	<2	10	23	<.2	<3	3	77	.28	.062	18	15	.66	248	.11	<3	1.39	.02	.14	6	30.4
2	23	13	65	<.3	9	10	701	3.24	21	<8	<2	16	22	<.2	<3	<3	75	.31	.073	24	18	.60	213	.10	<3	1.21	.02	.14	3	6.7
5	157	32	37	<.3	6	6	128	4.04	31	13	<2	33	49	<.2	<3	<3	6	.43	.062	33	2	.14	270	<.01	<3	.82	.01	.09	2	37.3
4	164	16	74	.3	10	10	100	4.70	98	14	<2	55	29	<.2	<3	3	9	.33	.101	26	5	.14	192	<.01	<3	1.09	.01	.06	3	162.2
5	50	29	197	.4	37	15	2478	4.44	384	<8	<2	10	32	.9	4	<3	76	.41	.100	23	38	.66	213	.12	3	1.79	.01	.15	5	46.0
7	78	29	526	.3	33	11	1093	6.17	3576	9	<2	6	22	1.7	16	<3	79	.39	.096	28	39	.63	240	.10	<3	1.78	.01	.12	<2	12.3
4	26	40	235	.5	21	13	949	4.48	724	<8	<2	6	24	.2	11	4	78	.44	.085	26	38	.61	171	.11	<3	1.77	.01	.11	2	12.2
3	16	60	40	.3	6	1	458	2.25	59	<8	<2	33	90	<.2	5	<3	5	.32	.028	49	4	.08	271	<.01	<3	.49	.01	.45	3	52.3
5	55	35	124	.3	16	10	750	4.19	459	<8	<2	15	33	<.2	11	<3	52	.35	.071	34	22	.44	224	.08	<3	1.19	.01	.17	5	410.6
3	39	38	245	.4	18	13	1225	4.12	702	<8	<2	16	35	.7	9	5	58	.38	.089	31	31	.52	225	.09	<3	1.45	.01	.12	<2	183.5
5	135	43	173	.5	19	10	889	2.95	284	<8	<2	22	29	.5	15	<3	45	.42	.066	33	22	.39	127	.04	<3	1.11	.01	.12	4	45.7
<1	23	40	218	.3	21	11	433	2.64	69	<8	<2	10	24	.2	<3	<3	62	.44	.075	32	40	.73	157	.15	<3	1.86	.02	.14	5	7.5
1	32	39	207	<.3	23	9	566	2.93	132	<8	<2	10	28	.2	4	<3	69	.50	.083	32	40	.70	163	.16	<3	1.68	.02	.15	3	16.0
1	32	42	208	<.3	23	10	573	2.90	130	<8	<2	10	28	.3	<3	<3	68	.51	.084	32	39	.71	163	.16	<3	1.68	.02	.15	<2	8.5
<1	58	101	336	1.6	26	13	1003	3.36	333	<8	<2	10	29	1.1	4	4	78	.49	.075	50	44	.64	242	.10	<3	2.32	.01	.11	<2	37.3
1	24	60	242	.7	19	13	651	3.08	310	<8	<2	7	22	.3	7	3	56	.38	.073	23	32	.56	169	.10	<3	1.63	.01	.09	<2	14.7
12	53	73	162	<.3	13	7	380	3.09	180	<8	<2	17	28	.2	10	<3	51	.37	.072	24	23	.46	188	.11	<3	1.08	.03	.19	4	74.7
34	98	40	72	.5	5	2	164	1.67	67	<8	<2	36	34	.2	23	<3	16	.20	.041	43	8	.15	217	.01	<3	.60	.01	.21	2	19.1
2	30	103	248	.8	19	13	562	3.01	185	<8	<2	11	25	.5	8	<3	62	.42	.075	21	33	.66	177	.12	<3	1.70	.01	.11	<2	15.8
49	132	46	143	<.3	11	6	334	2.70	103	<8	<2	20	28	.5	6	4	46	.31	.065	26	23	.37	181	.08	<3	1.00	.02	.13	4	19.4
13	55	72	233	.9	16	10	588	3.04	135	<8	<2	12	29	.5	5	<3	61	.37	.061	26	32	.56	246	.08	<3	1.82	.01	.11	<2	45.8
7	42	57	189	.5	17	12	645	2.81	109	<8	<2	9	25	.3	3	<3	57	.32	.059	20	30	.54	203	.09	<3	1.62	.01	.09	4	27.6
27	71	67	64	1.2	7	4	192	3.23	51	<8	<2	24	46	<.2	<3	<3	30	.23	.044	28	13	.28	260	.04	<3	.92	.04	.33	<2	70.5
81	77	26	15	.7	2	2	74	6.07	6	<8	<2	28	40	<.2	<3	5	5	.12	.027	16	<1	.09	151	<.01	<3	.62	.08	.96	<2	74.3
44	71	34	53	<.3	11	5	198	3.41	25	<8	<2	19	42	<.2	3	<3	48	.21	.045	19	24	.39	213	.08	<3	1.10	.03	.21	2	26.2
12	31	47	118	.4	15	7	237	3.03	62	<8	<2	6	30	<.2	5	4	65	.27	.063	20	27	.52	250	.07	3	1.71	.01	.10	2	42.4
7	27	46	100	.4	12	6	198	2.87	69	<8	<2	12	26	<.2	<3	4	56	.27	.061	20	25	.53	249	.09	<3	1.57	.01	.11	<2	97.4
8	44	55	100	.4	8	5	257	4.01	69	<8	<2	23	51	<.2	5	<3	45	.30	.064	25	17	.44	313	.06	<3	1.27	.02	.26	2	119.7
2	56	74	158	.6	17	11	538	3.21	150	<8	<2	10	32	<.2	6	3	72	.48	.085	29	34	.66	185	.13	<3	1.83	.02	.12	3	94.3
2	38	74	168	.3	21	14	510	2.69	69	<8	<2	10	33	.9	6	3	62	.46	.065	25	37	.70	228	.13	<3	1.82	.02	.15	3	17.5
<1	21	71	143	.3	25	12	876	2.97	170	<8	<2	5	19	.5	7	<3	76	.29	.075	14	34	.57	109	.10	4	1.76	.01	.08	3	9.3
<1	18	108	157	.8	23	9	538	2.50	225	<8	<2	6	17	.2	7	<3	57	.28	.062	15	35	.60	94	.10	3	1.93	.01	.07	3	5.9
1	24	523	661	2.0	35	14	1355	4.14	1165	<8	<2	11	26	3.3	29	4	64	.30	.063	28	44	.72	183	.05	<3	2.21	.01	.10	<2	27.9
14	132	29	163	.3	38	13	842	3.31	61	19	<2	4	31	10.7	9	9	84	.57	.084	18	178	.62	150	.12	4	1.85	.04	.17	10	210.1

Sample type: SOIL. Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.



SAMPLE#	Mo	Cu	Pb	Zn	Ag	Ni	Co	Mn	Fe	As	U	Au	Th	Sr	Cd	Sb	Bi	V	Ca	P	La	Cr	Mg	Ba	Ti	B	Al	Na	K	W	Au*
	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	%	ppm	ppm	%	ppm	%	%	%	%	%	ppm	ppb
CC 100+00W 124+50N	1	24	75	216	.6	25	11	619	3.31	668	<8	<2	6	21	.8	4	3	64	.33	.071	16	42	.66	104	.10	<3	2.13	.01	.09	2	26.2
CC 100+00W 124+00N	2	19	135	249	1.1	19	8	428	3.74	305	<8	<2	5	18	.4	23	<3	78	.23	.054	16	40	.56	96	.10	<3	2.15	.01	.08	<2	4.5
CC 100+00W 123+50N	1	24	177	192	2.0	19	7	435	2.65	729	<8	<2	4	23	.6	19	5	63	.33	.054	19	35	.54	114	.09	<3	1.53	.01	.09	4	35.2
CC 100+00W 123+00N	<1	38	476	596	3.0	16	9	733	4.72	1542	<8	<2	19	23	3.3	15	13	58	.24	.063	26	32	.48	164	.09	<3	1.57	.02	.23	<2	50.1
CC 100+00W 122+50N	1	24	128	381	1.0	21	10	614	3.04	628	<8	<2	6	27	1.3	18	<3	67	.37	.053	35	38	.65	148	.10	<3	1.80	.01	.11	<2	18.5
CC 100+00W 122+00N	<1	16	144	486	.9	16	9	771	2.65	328	<8	<2	5	26	2.3	8	<3	65	.43	.068	18	31	.51	107	.10	3	1.36	.01	.10	<2	6.2
CC 100+00W 121+50N	2	19	120	265	.7	15	7	620	2.60	329	<8	<2	5	24	1.5	5	3	66	.37	.051	22	33	.47	112	.10	<3	1.39	.02	.10	<2	11.1
CC 100+00W 121+00N	1	28	130	366	1.2	23	11	748	3.49	648	<8	<2	5	25	.8	12	4	70	.39	.079	24	39	.70	144	.07	<3	2.09	.01	.10	<2	35.6
CC 100+00W 120+50N	1	28	149	403	.8	22	11	767	3.40	716	<8	<2	7	30	1.1	8	4	67	.41	.058	34	39	.66	160	.08	<3	1.86	.01	.11	<2	14.2
CC 100+00W 120+00N	<1	29	85	387	1.0	22	10	744	3.08	361	<8	<2	6	30	.9	6	4	66	.43	.066	32	42	.75	168	.11	<3	2.08	.02	.11	<2	30.7
CC 100+00W 119+50N	1	25	71	247	.4	24	14	1012	3.46	214	<8	<2	7	30	1.1	<3	4	80	.47	.069	28	38	.70	169	.13	<3	2.08	.02	.12	<2	8.2
CC 100+00W 119+00N	1	31	44	214	.7	24	11	767	3.12	163	<8	<2	7	43	.6	4	<3	68	.78	.074	74	36	.71	229	.12	<3	2.08	.02	.13	3	26.1
CC 100+00W 118+50N	2	26	100	357	.8	22	11	755	3.30	478	<8	<2	7	30	.5	18	5	68	.50	.072	39	37	.75	180	.11	<3	1.99	.02	.18	<2	14.8
CC 100+00W 118+00N	3	43	84	417	1.1	32	16	900	3.77	264	<8	<2	7	39	1.2	9	3	90	.50	.080	50	49	.81	375	.13	3	2.63	.02	.14	<2	14.9
RE CC 100+00W 118+00N	2	43	80	419	1.1	32	16	907	3.79	264	<8	<2	7	40	1.1	9	4	90	.50	.081	50	49	.81	380	.13	<3	2.67	.02	.14	<2	14.7
CC 100+00W 117+50N	<1	26	60	279	.3	23	12	797	3.29	343	<8	<2	7	33	.3	11	<3	77	.45	.059	26	40	.72	212	.13	<3	2.11	.02	.14	2	12.4
CC 100+00W 117+00N	1	24	44	349	.3	24	12	1008	2.95	224	<8	<2	3	39	1.6	10	<3	79	.59	.058	17	41	.72	223	.13	<3	2.01	.02	.13	<2	5.8
CC 100+00W 116+50N	2	27	47	263	.3	26	12	752	3.19	148	<8	<2	7	28	.3	7	<3	84	.43	.063	17	44	.79	203	.15	3	2.03	.02	.14	2	7.5
CC 100+00W 116+00N	2	38	69	165	.7	26	11	417	2.90	80	<8	<2	6	44	<.2	6	3	83	.65	.066	31	43	.77	321	.11	<3	2.34	.02	.11	2	8.3
CC 100+00W 115+50N	2	26	57	133	1.1	21	12	606	3.65	146	<8	<2	8	23	<.2	<3	3	100	.27	.065	22	43	.65	198	.12	3	2.46	.02	.11	3	4.0
CC 100+00W 115+00N	2	31	68	196	.8	24	14	911	3.89	124	<8	<2	6	37	.2	4	5	97	.50	.082	31	40	.73	394	.11	3	2.53	.02	.11	3	6.5
CC 98+50W 108+00W	<1	29	18	72	<.3	14	9	255	4.05	25	<8	<2	12	18	<.2	<3	4	74	.31	.074	14	25	.67	146	.10	<3	1.89	.01	.09	2	30.1
CC 98+50W 107+50W	1	56	24	80	<.3	17	11	304	2.87	19	<8	<2	12	21	<.2	<3	4	77	.35	.057	26	34	.78	229	.11	<3	2.31	.02	.08	2	40.2
CC 98+50W 107+00W	1	55	18	147	<.3	18	16	475	2.87	18	<8	<2	13	23	.5	6	<3	77	.44	.079	31	31	.76	254	.15	<3	1.97	.01	.10	4	19.5
CC 98+50W 106+50W	<1	49	18	128	.3	23	22	2814	3.39	16	<8	<2	6	32	.5	7	<3	79	.47	.136	38	36	.61	382	.07	<3	2.35	.02	.08	3	15.7
CC 98+50W 106+00W	<1	36	20	162	<.3	22	11	431	3.24	37	<8	<2	11	28	<.2	<3	3	85	.49	.085	24	37	.80	285	.14	<3	2.10	.02	.11	4	22.9
CC 98+00W 108+00N	1	43	20	92	<.3	21	10	420	3.41	23	<8	<2	10	24	<.2	<3	3	84	.46	.095	23	33	.81	235	.14	<3	2.09	.02	.12	2	26.2
CC 98+00W 107+50N	<1	56	24	106	<.3	24	14	636	3.86	35	<8	<2	9	27	<.2	<3	5	89	.42	.078	19	40	.83	275	.11	<3	2.43	.01	.11	2	42.2
CC 98+00W 107+00N	<1	37	25	86	<.3	19	11	335	2.78	12	<8	<2	11	22	<.2	<3	3	72	.39	.061	20	36	.78	222	.12	<3	2.22	.01	.08	<2	42.4
CC 98+00W 106+50N	<1	61	18	93	<.3	21	11	568	3.22	25	<8	<2	11	22	.2	<3	5	78	.40	.087	31	39	.76	229	.12	<3	2.23	.01	.09	2	39.6
CC 98+00W 106+00N	2	42	23	137	<.3	25	14	608	4.01	39	<8	<2	13	25	.2	<3	<3	94	.45	.086	29	43	.88	317	.16	<3	2.38	.02	.14	3	85.5
CC 97+50W 108+00N	2	42	32	114	.3	22	15	478	2.64	22	<8	<2	10	27	.2	<3	<3	99	.47	.073	27	37	.88	328	.11	<3	2.29	.01	.09	2	27.8
CC 97+50W 107+50N	1	33	27	105	<.3	20	13	382	3.04	43	<8	<2	9	22	<.2	<3	5	83	.40	.074	20	33	.74	240	.12	<3	2.02	.01	.08	2	20.8
CC 97+50W 107+00N	1	33	23	88	<.3	19	10	470	3.05	37	<8	<2	7	24	<.2	<3	4	78	.41	.066	17	36	.75	212	.09	<3	2.15	.01	.07	4	39.7
STANDARD DS2	15	135	33	167	<.3	38	13	816	3.34	63	21	<2	4	29	11.0	7	11	83	.56	.086	17	175	.63	150	.11	4	1.85	.04	.17	10	225.3

Sample type: SOIL. Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.



SAMPLE#	Mo	Cu	Pb	Zn	Ag	Ni	Co	Mn	Fe	As	U	Au	Th	Sr	Cd	Sb	Bi	V	Ca	P	La	Cr	Mg	Ba	Ti	B	Al	Na	K	W	Au*
	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	%	%	ppm	ppm	%	ppm	%	ppm	%	%	%	ppm	ppb
CC 97+50W 106+50N	1	35	33	111	<.3	19	10	451	2.88	34	<8	<2	8	22	<.2	<3	3	78	.37	.076	18	37	.72	191	.08	4	2.01	.01	.09	2	29.8
CC 97+50W 106+00N	1	40	23	111	<.3	26	11	866	4.08	38	<8	<2	12	28	<.2	<3	<3	97	.50	.090	20	42	.87	325	.10	3	2.59	.01	.15	3	24.9
CC 96+00W 126+50N	1	37	82	214	1.0	27	13	2070	3.25	318	<8	<2	6	48	2.0	35	<3	61	.91	.101	32	36	.58	215	.04	4	1.78	.02	.10	3	22.6
CC 96+00W 126+00N	1	21	49	133	<.3	22	10	822	2.63	152	<8	<2	9	21	.4	16	<3	62	.32	.060	20	30	.51	127	.06	<3	1.37	.01	.08	4	18.3
CC 96+00W 125+50N	1	26	34	101	.4	24	10	575	2.67	124	<8	<2	8	28	.2	14	3	73	.48	.073	22	36	.56	136	.08	<3	1.35	.02	.08	2	19.0
CC 96+00W 125+00N	<1	25	47	126	.5	18	8	489	2.49	150	<8	<2	10	27	.5	12	<3	60	.45	.056	24	31	.51	115	.08	3	1.41	.02	.10	3	24.4
CC 96+00W 124+50N	2	34	105	206	1.0	20	11	935	3.38	373	<8	<2	13	28	.6	23	<3	57	.30	.059	37	35	.51	135	.05	3	1.76	.01	.12	5	33.5
CC 96+00W 124+00N	2	37	264	296	1.6	15	10	1141	3.72	1295	<8	<2	13	30	1.5	23	<3	46	.25	.064	25	23	.32	131	.04	3	1.16	.01	.19	2	59.7
CC 96+00W 123+50N	1	31	53	102	.3	24	9	356	2.70	81	<8	<2	12	19	<.2	16	<3	66	.27	.059	30	35	.62	142	.08	<3	2.00	.01	.08	3	16.3
CC 96+00W 123+00N	2	37	128	187	.7	29	15	929	3.84	242	<8	<2	7	24	.9	37	5	72	.34	.093	33	38	.67	179	.06	3	2.00	.01	.10	6	15.5
CC 96+00W 122+50N	2	25	47	132	<.3	25	12	686	3.20	361	<8	<2	7	31	<.2	10	<3	72	.36	.078	17	33	.69	138	.09	<3	1.91	.01	.12	4	14.2
CC 96+00W 122+00N	1	56	21	83	<.3	26	12	411	3.26	13	<8	<2	7	25	<.2	<3	<3	77	.39	.078	27	41	.86	137	.11	<3	2.41	.01	.13	<2	2.9
CC 96+00W 121+50N	<1	29	13	69	<.3	27	11	441	2.99	7	<8	<2	7	24	<.2	3	<3	77	.37	.075	20	39	.77	130	.12	<3	2.09	.02	.12	<2	2.3
CC 96+00W 121+00N	1	26	17	69	<.3	25	11	469	3.19	13	<8	<2	7	20	<.2	5	<3	75	.27	.052	21	37	.79	107	.11	<3	2.12	.01	.12	<2	3.3
RE CC 96+00W 121+00N	1	24	14	67	<.3	25	11	453	3.10	14	<8	<2	7	20	<.2	5	<3	73	.27	.051	20	36	.77	105	.11	4	2.10	.01	.12	2	1.8
CC 96+00W 120+50N	2	31	112	305	1.7	23	13	1123	3.65	662	<8	<2	9	28	.2	65	5	74	.28	.052	38	40	.81	160	.10	3	2.40	.01	.17	<2	26.0
CC 96+00W 120+00N	3	35	195	517	2.2	29	9	282	5.70	430	8	<2	10	24	.2	17	<3	82	.31	.089	19	47	.71	264	.07	3	2.61	.01	.17	2	25.2
CC 96+00W 119+50N	2	38	199	252	2.3	29	9	325	2.77	131	<8	<2	7	26	.8	46	<3	79	.38	.079	22	52	.92	246	.11	<3	2.41	.01	.15	<2	22.3
CC 92+00W 127+00N	2	30	47	138	.4	33	12	701	3.19	169	<8	<2	7	25	.2	13	<3	79	.38	.087	21	41	.68	174	.08	<3	1.87	.01	.10	<2	9.6
CC 92+00W 126+50N	3	32	84	253	.8	36	13	878	3.69	247	<8	<2	7	26	.5	31	<3	82	.31	.068	21	43	.70	202	.06	<3	2.06	.01	.11	<2	10.7
CC 92+00W 126+00N	3	36	81	261	.7	34	15	1069	3.75	288	<8	<2	5	27	.3	18	3	91	.33	.080	26	47	.75	255	.05	<3	2.32	.01	.10	<2	25.6
CC 92+00W 125+50N	3	30	93	284	2.7	31	11	721	2.92	328	<8	<2	12	18	.3	21	<3	66	.27	.082	25	42	.66	162	.06	<3	2.03	.01	.16	<2	33.2
CC 92+00W 125+00N	4	83	85	267	5.1	35	10	2934	3.08	750	15	<2	4	42	3.3	21	4	56	.53	.210	55	38	.42	355	.02	<3	2.00	.01	.09	<2	58.6
CC 92+00W 124+50N	4	56	94	367	3.2	32	10	1350	3.75	1112	9	<2	8	50	3.2	25	6	49	.68	.126	56	30	.42	357	.02	<3	1.83	.01	.14	2	81.0
CC 92+00W 124+00N	3	42	115	274	3.2	62	11	579	3.91	1196	<8	<2	12	31	<.2	17	14	79	.46	.095	27	105	1.06	247	.08	<3	2.26	.01	.21	<2	131.3
CC 92+00W 123+50N	4	36	76	232	2.1	33	9	370	3.59	743	<8	<2	7	24	<.2	13	13	81	.31	.058	20	44	.77	186	.07	<3	2.00	.01	.11	6	51.8
CC 92+00W 123+00N	4	36	107	397	1.0	38	18	1380	3.69	215	<8	<2	4	38	1.1	23	<3	93	.48	.082	20	43	.73	359	.06	<3	1.96	.01	.10	<2	8.9
CC 92+00W 122+50N	3	37	206	471	3.9	38	17	1810	3.97	404	<8	<2	11	32	1.4	36	<3	92	.39	.083	28	56	.84	315	.07	3	2.40	.01	.10	<2	14.3
CC 92+00W 122+00N	3	30	160	309	2.3	29	8	511	2.93	270	<8	<2	7	26	.3	29	<3	72	.35	.078	15	32	.63	186	.05	<3	1.80	.01	.11	<2	8.9
CC 92+00W 121+50N	4	41	131	243	2.0	42	14	817	3.52	320	<8	<2	5	30	.3	24	<3	97	.39	.071	15	56	.80	161	.07	<3	1.96	.01	.12	<2	13.5
CC 92+00W 121+00N	5	57	515	324	7.0	41	13	616	4.05	1247	<8	<2	6	35	1.5	24	<3	108	.43	.136	16	47	.69	231	.08	3	1.60	.02	.18	<2	18.4
STANDARD DS2	14	138	31	168	.3	39	14	829	3.42	62	18	<2	4	30	11.5	7	11	85	.57	.088	18	182	.64	153	.09	4	1.87	.04	.17	10	217.4



Sample type: SOIL. Samples beginning 'RE' are Reruns and 'RRE' are Reject Reruns.

Appendix 3

094206

97-PR-36

Altered, veined Intrusive (granodiorite?)



Photomicrographs 00R V 1 and 3

Scale 0.1 mm _____

Reflected light

Scale 0.1 mm. _____

Pictured: pyrite, chalcopyrite, and gold (fresh polish)

Summary Description

Altered intrusive, with abundant original quartz, strongly sericite- and quartz-altered feldspar, and chloritized, silicified mafics. Quartz and pseudomorphs after feldspar show an interlocking relationship. Original rock could have been a granodiorite (very little yellow staining is visible on the slab stained for K-spar, and quartz is abundant). Biotite pseudomorphs are recognizable among the mafics, with probable pseudomorphs after hornblende. At one end of the section is a banded (multistage) vein containing quartz with open spaces, pyrite, chalcopyrite, traces of galena, and native gold.

Microscopic Description

Transmitted Light

Quartz; 30-35%, anhedral (<0.01 to ~5 mm). Original quartz appears to have had an interlocking granitic texture with mafics and feldspar (now altered). This quartz shows strained extinction. In addition to original quartz there is some irregular veining in and silicification of altered plagioclase and mafics. There is also a drusy quartz and sulphide vein at the edge of the section containing native gold. Introduced quartz is relatively weakly strained to unstrained.

Sericite; 10-20%, anhedral (<0.01 to 0.05 mm). Masses of fine sericite with lesser intermixed quartz form pseudomorphs, presumably after feldspar (plagioclase). Typically has a few pyrite and chalcopyrite grains and patchy iron staining. Original feldspar would have had an interlocking relation with quartz.

Clay minerals; $\leq 10\%$, microcrystalline. Not positively identified, but appears to be present in patches among the sericite. This material has birefringence lower than is typical of sericite. Commonly iron stained, as for the sericite.

Chlorite; 7-10%, anhedral (<0.01 to 0.3 mm). Fine bladed chlorite occurs in pseudomorphs after mafic minerals, where it is intermixed with quartz (and probably albite) and lesser pyrite. The mafic pseudomorphs are commonly subhedral and in some cases euhedral, and have forms suggesting both biotite and hornblende.

Plagioclase; traces+, anhedral to subhedral (<0.01 to 0.5 mm). A few remnants survive, strongly sericite-altered or surrounded by sericite.

Zircon; traces, anhedral (<0.01 to 0.1 mm). Very sparse grains with very high positive relief and high birefringence. Found in pseudomorphs after mafics and feldspar.


Apatite; traces, anhedral to subhedral (0.01 to 0.5 mm). A few prismatic apatite crystals are found in chloritized mafics.

Fe oxide staining; traces, amorphous. Found in fractures and spreading through sericite, apparently associated with pyrite.

Veins:

Multistage quartz vein at the edge of the section is ≤ 1 cm in width and consists of drusy, euhedrally-terminated quartz with open spaces, some of which have been filled with pyrite, lesser chalcopyrite, traces of galena, tetrahedrite, and native gold. There appear to be two 'bands' or stages of quartz and sulphide with a thin strip of wallrock in between (however the vein is at the edge of the section). Chalcopyrite and the gold are found near the wallrock.

094206



Photomicrograph 00R V 18

Reflected light and plane polarized light

Scale 0.1 mm ____

Pictured: quartz and sulphides in vein

Reflected Light

Pyrite; 7-12%, anhedral to euhedral (<0.01 to ~ 1 mm). Concentrated in a quartz vein at the edge of the section where it commonly fills drusy cavities. Also disseminated in the altered wallrock, in the pseudomorphs (not in original quartz), and with some preferential concentration in mafic pseudomorphs.

Chalcopyrite; 1-2%, anhedral (<0.01 to 0.5 mm). Associated with pyrite, both in the vein and disseminated in the wallrock. Contains a few euhedral pyrite inclusions -- probably later than much of the pyrite.


Leucoxene; traces+, anhedral (<0.01 to 0.1 mm). Commonly found with pyrite in pseudomorphs after mafics, particularly biotite.

Covellite ($\pm$ chalcocite); traces, anhedral (<0.01 to 0.05 mm). Minor local alteration of chalcopyrite around grain edges.

Galena; traces, anhedral (<0.01 to 0.05 mm). Small inclusions in pyrite, in some cases with chalcopyrite. Some is also found at edges of coarse pyrite, with chalcopyrite.

Tetrahedrite; traces, anhedral (<0.01 to 0.05 mm). Inclusions / blebs in pyrite. Occurrence similar to the galena.

Native Au; trace, anhedral (<0.01 to 0.05 mm). A few small grains are found in the quartz-sulphide vein, at the contact between chalcopyrite and pyrite (see photomicrograph).



LEGEND

Trails
Trench
Lake
River
Stream
Tress
Contours
Index
Intermediate

Sample Sites

Rock X Pb, Zn, Ag, As (ppm)

Silt / Pb, Zn, Ag, As (ppm)

Soil Pb, Zn, Ag, As (ppm)

Road cut / Cu, Mo (ppm), Au (ppb)

○ Diamond Drill Hole

1999 additions from double lined lines

○ 100 ppm Arsenic contour

○ 50 ppm Lead contour

Alexis Resources Ltd.

CANADIAN CREEK PROJECT

Whitehorse M.D., Yukon

ROCK, SILT and SOIL

GEOCHEMISTRY


Pb, Zn, Ag, As (ppm)

Scale 1 : 5000

Date April 2000

By J.C. / gel/qpb

Figure 7



LEGEND

- Trails
- Trench
- Lake
- River
- Stream
- Trees
- Contours
- Index
- Intermediate

- 4** CASINO COMPLEX - Undivided
 - 4e** Hornblende Intrusive Breccia
 - 4d** Metatuffaceous Intrusive Breccia
 - 4c** Patton Porphyry
 - 4b** Quartz Monzonite Breccia
 - 4a** Quartz Monzonite
- Kqd** KOFFEE CREEK QUARTZ DIORITE STOCK
- 3** DAWSON RANGE BATHOLITH - Undivided
 - 3b** Hornblende Biotite Granodiorite
 - 3a** Biotite Granodiorite
- 2** PRE DAWSON RANGE INTRUSIVES
 - Polished Quartz Diorite

- Outcrop
- Faults interpreted from airphoto
- Geological contact
- K-5 Rock sample site and sample number
- K-4 Silt " " " " "
- TR-3 Trench

- Proposed drill site
- Trench location and sample number
- Start of soil line
- Claim L.C.P.

- cp - chalcopyrite
- sil - silicification
- py - pyrite
- mg - magnetite
- bx - breccia
- b - bleached
- QM - quartz monzonite
- GDR - granodiorite
- QD - quartz diorite

0 50 100 200 300 400 500 METRES


CANADIAN CREEK PROJECT
Whitehorse, M.D., Yukon



064206
PROPERTY GEOLOGY
8 SAMPLE LOCATIONS

MINCORD
EXPLORATION
CONSULTANTS
LIMITED


Scale 1 : 5000
Date April 2000
By

NTS 115-J/10
Map 5





LEGEND



- 4 CASINO COMPLEX - Undivided**
 - 4d** Metrolithic Breccia
 - 4c** Patton Porphyry
 - 4b** Quartz Monzonite Breccia
 - 4a** Quartz Monzonite
- 3 DAWSON RANGE BATHOLITH - Undivided**
 - 3b** Hornblende Biotite Granodiorite
 - 3a** Biotite Granodiorite
- 2 PRE DAWSON RANGE INTRUSIVES - Undivided**
 - 2a** Foliated Quartz Diorite
- 1 YUKON METAMORPHIC COMPLEX**
Gneiss and Quartzite

- Trench location (1997)
- Claim L.C.P.
- Diamond Drill Hole
- Rock Sample Site and Sample Number
- Trench (1993)
- Outcrop
- Foliation
- Py - Pyrite
- To - Tourmaline
- Bx - Breccia
- Faults visible in outcrop
- Faults interpreted from airphoto
- Geological Contact (defined, assumed)
- Alteration Boundary

0 50 100 200 300 400 500 METRES

CANADIAN CREEK PROJECT
Whitehorse M.D., Yukon

GEOLOGY⁹⁴²⁰⁶
& SAMPLE LOCATIONS

MINCORD
EXPLORATION
CONSULTANTS
LIMITED

Scale 1 : 5000
Date April 2000
By JC/JP
MTS 115-JV10
Figure 4

Sample #	Pb	Zn	Ag	As	Sample #	Cu	Mo	Au
97-PR-1	13	23	<3	9	97-PR-1	80	7	32
97-PR-2	10	9	<3	23	97-PR-2	4	2	35
97-PR-3	16	17	<3	16	97-PR-3	78	<1	33
97-PR-4	<3	9	<3	11	97-PR-4	64	11	58
97-T21-1	4	25	0.7	37	97-T21-1	381	7	130
97-T21-2	7	11	<3	2	97-T21-2	40	21	44
	14	56	0.8	25	97-T21-2	232	48	66