

Assessment Report on the

2015 DRILLING PROGRAM

on the

BOULEVARD PROJECT, YUKON

Grant Number	Claim Name	Grant Number	Claim Name
YC46726 - YC46733	Tiger 1 - Tiger 8	YD51301 - YD51328	XT 301 - XT 328
YC64776 - YC64871	BLVD 1 - BLVD 96	YD51331 - YD51358	XT 331 - XT 358
YC65316 - YC65319	AVE 1 - AVE 4	YD51361 - YD51378	XT 361 - XT 378
YC81284 - YC81315	DRIVE 1 - DRIVE 32	YD51595 - YD51640	XY 95 - XY 140
YC82926 - YC83031	BLVD 97 - BLVD 202	YD51642 - YD51701	XY 142 - XY 201
YC98845 - YC98852	HAN 1 - HAN 8	YD51801 - YD51810	BDW 1 - BDW 10
YD08611 - YD08694	SOL 1 - SOL 84	YD51901 - YD51932	XZ 1 - XZ 32
YD12511 - YD12516	VIP 501 - VIP 506	YD61666 - YD61667	VIP 326 - VIP 327
YD15231 - YD15248	GRT 1 - GRT 18	YD62088 - YD62097	VIP 388 - VIP 397
YD15269 - YD15270	GRT F 19 - GRT F 20	YD62182 - YD62195	VIP 342 - VIP 355
YD15271 - YD15286	GRT 21 - GRT 36	YD62198 - YD62200	VIP 358 - VIP 360
YD49101 - YD49194	BDW 11 - BDW 104	YD62233 - YD62244	VIP 433 - VIP 444
YD50701 - YD50754	XY 1 - XY 54	YD62257 - YD62268	VIP 457 - VIP 468
YD50756 - YD50780	XY 56 - XY 80	YD62287 - YD62292	VIP 487 - VIP 492
YD51236 - YD51238	XT 236 - XT 238	YD62299 - YD62300	VIP 499 - VIP 500
YD51240	XT 240	YD62301 - YD62309	VIP 361 - VIP 369
YD51242	XT 242	YD62384 - YD62387	VIP 384 - VIP 387
YD51244	XT 244	YE27193 - YE27206	XY 81 - XY 94
YD51269 - YD51286	XT 269 - XT 286	YE27207	XY 141

WHITEHORSE MINING DISTRICT
Dates Worked: June 15 - September 11, 2015

NTS 50k Mapsheet: 115J13, 115J14
 UTM 566,500E; 6,969,000N (NAD 83, Zone 7)

Prepared by:
 David Gale PGeo, MSc
 Kendra A. Johnston, PGeo, BSc
 Independence Gold Corp.

Prepared for:
 Independence Gold Corp.
 P.O. Box 11584
 Suite 1410 – 650 West Georgia Street
 Vancouver, British Columbia, V6B4N8

December 16, 2015

Table of Contents

SUMMARY	4
INTRODUCTION	4
CLAIM DATA AND OWNERSHIP	6
PROPERTY DESCRIPTION	7
LOCATION	7
CLIMATE AND GEOMORPHOLOGY	7
INFRASTRUCTURE	8
HISTORY	8
PREVIOUS WORK	8
RECENT HISTORY	10
GEOLOGICAL SETTING	11
REGIONAL GEOLOGY	11
PROPERTY GEOLOGY	11
GEOPHYSICS	15
DRILLING	15
SUNSET-SUNRISE ZONE DRILLING	15
DENALI ZONE DRILLING	26
SAMPLING PROCEDURES	28
QUALITY ASSURANCE/QUALITY CONTROL	28
DISCUSSIONS AND CONCLUSIONS	30
SUNSET-SUNRISE ZONE	30
DENALI ZONE	32
RECOMMENDATIONS	33
SUNSET-SUNRISE ZONE	33
DENALI ZONE	34

REFERENCES	35
STATEMENT OF QUALIFICATIONS.....	36
STATEMENT OF QUALIFICATIONS.....	37
STATEMENT OF EXPENDITURES.....	38

List of Tables

Table 1 – Boulevard Project Claim Information.....	6
Table 2 – Drill Hole Summary Data.....	17
Table 3 - 2015 Significant Assay Intervals.....	18

List of Figures

Figure 1 – Location Map	5
Figure 2 – Boulevard Claim Map.....	in back pocket
Figure 3 – Regional Geology.....	12
Figure 4 – Total Field Ground Magnetic Survey Data for the Sunrise Zone.....	16
Figure 5 – Total Field Ground Magnetic Survey Data for the Denali Zone.....	17
Figure 6 – Drill Location Map for the Sunrise-Sunset....	20
Figure 7 – Drill Location map for the Denali Target.....	21
Figure 8 – Splitter used to obtain 2-5 kg representative sample.....	28
Figure 9 – Blank Gold Values from the 2015 drilling programs.....	29
Figure 10 – Gold standard values from the 2015 drilling programs.....	30

List of Appendices

Appendix 1 – Ground Geophysical Results

Appendix 2 – Drill Logs

Appendix 3 – Laboratory Assay Certificates, Certification and Analytical Method Summaries

SUMMARY

The Boulevard Project (“Boulevard”), consisting of 4 contiguous properties: Boulevard, Tiger, Solitude and YCS, is an early stage exploration project, operated by Independence Gold Corp. (“InGold” or “the Company”). Work completed in 2015 focused primarily on drill testing geologic, geophysical and soil targets. Activities over the past ten years have targeted structurally hosted, orogenic gold mineralization. During this time, InGold and its predecessors have conducted multiple rounds of soil geochemical surveys, collecting more than 19,000 soil and rock samples within the project’s boundary. This year, a total of 2,841 metres (m) were drilled on the Boulevard Project including, 1,912 m at the Sunset-Sunrise zone and 929 m at the Denali Zone. The 2015 exploration program also included 73 line kilometres (km) of ground geophysics at the Sunrise and Denali Zones. The work in 2015 successfully discovered the new mineralized Sunrise Zone, and confirmed the presence of the Denali Zone in bedrock by drilling.

The Boulevard Project, comprising 888 quartz claims, is located in west-central Yukon approximately 135 km south of Dawson City, Yukon, 35 km south of Kinross Gold Corp’s White Gold deposit, and contiguous to the western side of the Coffee Gold project owned by Kaminak Gold Corporation (Figure 1).

INTRODUCTION

Early in 2015 a review of all the Company’s Yukon-based exploration properties was conducted to determine their exploration potential. Drilling programs were approved for the Sunset and Denali zones on the Boulevard Project. Two reverse circulation drill programs were conducted between June and September, 2015. This report provides a discussion of results, interpretations and recommendations for the two programs. The authors participated in and managed the programs. The Statements of Qualification are contained within this report.

The objective of the drilling was to delineate continuous zones of gold mineralization on Boulevard by testing linear magnetic trends that are coincident with gold, arsenic and antimony geochemical soil anomalies.



CLAIM DATA AND OWNERSHIP

InGold's predecessor Silver Quest Resources Ltd. ("Silver Quest") acquired the BLVD claims from Rimfire Minerals and Northgate Minerals in July 2009. Through various staking campaigns, and the acquisition of the VIP claims from Goldspike Exploration Inc., the project was expanded in 2010 and 2011. The project was transferred to the newly formed Independence Gold in January 2012 after Silver Quest was taken over by New Gold Inc. Quartz claims are registered with the Whitehorse Mining Recorder. This report covers 888 contiguous quartz mining claims, an approximate area of 18,568 hectares (ha) which encompasses the entire Boulevard Project.

Boulevard centres on UTM 566,500E and 6,969,000N (NAD 83, Zone 7) on NTS map sheets 115J13, 115J14, 115O03 and 115O04 as shown on Figure 2 (in pocket). A five-year, Class 3 Permit (LQ00328) was granted to Silver Quest on September 1, 2011 and transferred to InGold in 2012. Claim data for the claims covered in this report are listed below.

Table 1 – Boulevard Project Claim Information

Grant Number	Claim Name	Registered Owner
YC46726 - YC46733	Tiger 1 - Tiger 8	Independence Gold Corp.
YC64776 - YC64871	BLVD 1 - BLVD 96	Rimfire Minerals 50%, Aurico Gold 50%
YC65316 - YC65319	AVE 1 - AVE 4	Rimfire Minerals 50%, Aurico Gold 50%
YC81284 - YC81315	DRIVE 1 - DRIVE 32	Rimfire Minerals 50%, Aurico Gold 50%
YC82926 - YC83031	BLVD 97 - BLVD 202	Rimfire Minerals 50%, Aurico Gold 50%
YC98845 - YC98852	HAN 1 - HAN 8	Independence Gold Corp.
YD08611 - YD08694	SOL 1 - SOL 84	Independence Gold Corp.
YD12511 - YD12516	VIP 501 - VIP 506	Independence Gold Corp.
YD15231 - YD15248	GRT 1 - GRT 18	Independence Gold Corp.
YD15269 - YD15270	GRT F 19 - GRT F 20	Independence Gold Corp.
YD15271 - YD15286	GRT 21 - GRT 36	Independence Gold Corp.
YD49101 - YD49194	BDW 11 - BDW 104	Independence Gold Corp.
YD50701 - YD50754	XY 1 - XY 54	Independence Gold Corp.
YD50756 - YD50780	XY 56 - XY 80	Independence Gold Corp.
YD51236 - YD51238	XT 236 - XT 238	Independence Gold Corp.
YD51240	XT 240	Independence Gold Corp.
YD51242	XT 242	Independence Gold Corp.
YD51244	XT 244	Independence Gold Corp.
YD51269 - YD51286	XT 269 - XT 286	Independence Gold Corp.
YD51301 - YD51328	XT 301 - XT 328	Independence Gold Corp.

YD51331 - YD51358	XT 331 - XT 358	Independence Gold Corp.
YD51361 - YD51378	XT 361 - XT 378	Independence Gold Corp.
YD51595 - YD51640	XY 95 - XY 140	Independence Gold Corp.
YD51642 - YD51701	XY 142 - XY 201	Independence Gold Corp.
YD51801 - YD51810	BDW 1 - BDW 10	Independence Gold Corp.
YD51901 - YD51932	XZ 1 - XZ 32	Independence Gold Corp.
YD61666 - YD61667	VIP 326 - VIP 327	Independence Gold Corp.
YD62088 - YD62097	VIP 388 - VIP 397	Independence Gold Corp.
YD62182 - YD62195	VIP 342 - VIP 355	Independence Gold Corp.
YD62198 - YD62200	VIP 358 - VIP 360	Independence Gold Corp.
YD62233 - YD62244	VIP 433 - VIP 444	Independence Gold Corp.
YD62257 - YD62268	VIP 457 - VIP 468	Independence Gold Corp.
YD62287 - YD62292	VIP 487 - VIP 492	Independence Gold Corp.
YD62299 - YD62300	VIP 499 - VIP 500	Independence Gold Corp.
YD62301 - YD62309	VIP 361 - VIP 369	Independence Gold Corp.
YD62384 - YD62387	VIP 384 - VIP 387	Independence Gold Corp.
YE27193 - YE27206	XY 81 - XY 94	Independence Gold Corp.
YE27207	XY 141	Independence Gold Corp.

PROPERTY DESCRIPTION

LOCATION

Boulevard is located in the Independence Creek area on the south side of the Yukon River in west-central Yukon, approximately 135 km south of Dawson City, Yukon (Figure 1).

CLIMATE AND GEOMORPHOLOGY

Boulevard lies within the Dawson Range, an area characterized by its rolling hills. Local elevations range from 400 m to 1,700 m above sea level. The higher elevation areas of the property are above tree line and covered with blocky felsenmeer and a thin layer of moss cover. Lower elevations support a mixture of stunted aspen, birch and spruce forest with thick willow and birch brush. Cold north facing slopes contain permafrost, and favour spruce tree growth, in contrast to the warm south facing slopes where permafrost is more intermittent, and aspen and birch trees along with ground brush predominate the vegetation cover of the hill sides.

In spite of the mild summer temperatures, permafrost can be found throughout the geographic region. Locally, permafrost is discontinuous, depending on slope direction, elevation and drainage patterns. Regionally, multiple freeze thaw cycles have resulted in an abundance of

felsenmeer covered slopes and plateaus. The Dawson Range remained unglaciated during the Pleistocene, making outcrops rare, and maintaining a soil profile that is relatively in-place. The few outcrops that are present are located along sparsely vegetated ridges and in main creek drainages.

INFRASTRUCTURE

Access to Boulevard in 2015 was via Piper Navajo, Cessna Caravan or deHavilland Beaver aircraft operated by Tintinia Air of Whitehorse Yukon. These planes were based in Whitehorse and flew to the Independence airstrip located on the Boulevard Project. Access to other parts of the project was via a Bell 407 helicopter operated by either Summit Helicopters of Yellowknife, Northwest Territories or Range Helicopters out of Drayton Valley, Alberta.

Alternatively, several barge operators are available for hire to barge supplies down the Yukon River from Whitehorse or Carmacks. A barge landing does not currently exist at Boulevard; however one could be built off claim at the confluence of Independence Creek and the Yukon River with the appropriate permits. Supplies would need to be mobilized by helicopter from there to the project area. There is a barge landing near by at the confluence of Coffee Creek and the Yukon River, this landing belongs to Kaminak and could be used with permission.

There are no roads that provide access to the Boulevard project.

A tent camp was located 2.5 km north of the Independence airstrip (Camp Coordinates: Lat: 62° 51' 41", Long: 139° 41' 42").

HISTORY

PREVIOUS WORK

A staking rush of historic proportions followed the discovery of the Casino Copper Gold project in 1969. At this time much of the land now included in Boulevard was staked and worked. The first known work in the area was completed by the Dawson Range Joint Venture and consisted of a regional silt sampling program during July and August 1969, targeting Casino-style copper-molybdenum mineralization. Positive results from the regional silt program led to more staking

and the completion of a soil sample grid on the eastern portion of the current Boulevard Project. The soil survey conducted by the Dawson Range Joint Venture identified an anomaly with greater than 10 grams per tonne (g/t) molybdenum over an area of 12 hectares. A follow-up trenching program consisting of 7 tractor trenches was conducted in 1970 resulting in the Toni-Tiger discovery, Minfile occurrence 115J 052, later re-staked by S. Ryan as the Tiger claims in June 2006.

In December of 1969, Fawn Bay Development Company Ltd. and Hanna Gold Mines Ltd. staked the FBH claims on the south-western portion of Boulevard and conducted reconnaissance grid soil sampling and limited geological mapping in 1970 (Minfile 115J 048). At this time two other companies were actively staking in the area. Gold Hawk Exploration Ltd. was staking the Monarch claims a few kilometres south of Toni Tiger, while Northern Empire Mines Ltd. was staking the Keg claims a few kilometres to the west of Toni Tiger. No exploration was carried out at the time on either of these claim blocks (Minfile 115J 050 and 115 051).

The majority of claims staked in 1969 lapsed within a couple of years, and the land lay untouched until 2006. In June 2006, Rimfire Minerals and Northgate Minerals formed a Joint Venture to carry out a regional silt sampling program in selected areas of the Dawson Range, targeting Pogo-style intrusion-related gold systems. Several elevated gold values including one silt sample which returned 323 parts per billion gold (ppb) were returned from a creek draining the land west of Toni Tiger. These samples were followed up on in August of 2006 by prospecting, geological mapping and conducting a reconnaissance style soil sampling grid consisting of approximately 400 samples at 100 m sample intervals (Roberts and Baker, 2007). These results led to the staking of the central core of the Boulevard property in 2007.

In 2007, the Joint Venture between Rimfire Minerals and Northgate Minerals conducted a detailed soil sampling program, collecting 1,707 soils and completed an additional round of staking in 2008 creating the current Boulevard property. During the spring and early summer of 2008, the Joint Venture carried out a significant exploration program consisting of excavator trenching, prospecting, soil sampling and minimal geological mapping. The group returned in the fall of 2008 to extend the soil sampling grid, to complete a ground magnetic and an induced polarization geophysical surveys and completed 7 diamond drill holes (Minfile 115J 050). In

July of 2009, the Joint Venture optioned the Boulevard property to Silver Quest, who conducted a short reconnaissance soil sample program that fall.

In 2009, Atac Resources Ltd. was also on the ground in the area, staking and working the Han claims (Minfile 115J 048). Atac Resources Ltd. completed a reconnaissance style soil sample survey consisting of 56 samples on behalf of Archer, Cathro & Associates (1981) Limited (Smith, 2010), who optioned these claims to Silver Quest in December 2009.

RECENT HISTORY

2010 was an exciting year for Yukon exploration enthusiasts and for the Dawson Range in particular. Kinross Gold Corp announced the takeover of Underworld Resources for \$139.2 million in March (Fleming, 2010) and Kaminak Gold Corp announced a discovery diamond drill hole grading 17.1 g/t gold over 15.5 m at the end of May (Carpenter, 2010a) followed by a second diamond drill hole 2 weeks later which graded 1.08 g/t over 84 m (Carpenter, 2010b). The news from Kaminak sparked another staking rush of historic proportions across the district and marks the beginning of a significant period of growth for the Boulevard project.

During the 2010 field season Silver Quest completed a large exploration program with the help of Equity Exploration Consultants Ltd. The program included a detailed sampling program collecting 4,362 soils, 62 rocks and 8 silt samples, 62.5 line km of ground geophysics, 20 diamond drill holes totalling 3,006 m and minor amounts of prospecting and mapping (Baker 2011). During July of that year, Silver Quest staked the VIP claims (YCS Property), extending the project to the north and truncating Kaminak's ground to the west. In August 2010, Silver Quest acquired the Tiger claims from S. Ryan.

Silver Quest continued to work the Boulevard project in 2011. A variety of contractors as well as Silver Quest staff were used to complete the work. The majority of exploration work focused on soil sampling to identify anomalous geochemical trends in soil, as well as to identify drill targets. A total of 5,473 soil samples were taken, 9 diamond drill holes were completed (1,434 m) and 3,410 line km of airborne geophysics including magnetics and radiometrics were flown. At the end of the season Silver Quest conducted a small staking program, staking 592 claims on the north end of the YCS property; these new claims comprised the Solo property and were allowed to lapse in March 2015.

In December of 2011, Silver Quest was taken over by New Gold Inc. As part of the transaction Silver Quest's Yukon properties were transferred to Independence Gold Corp. In 2012, continued exploration by the Company included the collection of 5,906 soil samples from a variety of detailed sampling grids and selected ridge and spur sample lines which successfully extended the Sunset and Vegas Zones discovered in 2010, by 100 m to reach a total strike length of 2.2 km. Samples were collected at 100 m intervals along ridges, or at 25 m intervals in grid formation. The 2012 season concluded with a 240 m trenching program on the YCS property, which confirmed mineralization in bedrock within the Denali Zone.

Today, Boulevard encompasses the Boulevard, Solitude, YCS and Tiger properties.


GEOLOGICAL SETTING

REGIONAL GEOLOGY

Boulevard is situated within the Yukon-Tanana Terrane approximately 130 km southwest of the Tintina Fault in west-central Yukon. This area is characterized by various pericratonic terranes that were accreted to the ancestral continental margin of North America in the early Jurassic. During the mid-Cretaceous the pericratonic terranes were intruded by a northwest-southeast trending plutonic suite known as the Dawson Range Plutonic Belt (Hart et al., 2004).

PROPERTY GEOLOGY

Boulevard has most recently been mapped and interpreted by G. McKenzie, M. Allan, and C. Hart of the Mineral Deposit Research Unit at the University of British Columbia. McKenzie (2014) completed a master's research thesis on the geology and mineralization of the Boulevard project (Figure 3). Ryan et al. (2014) recently mapped the Stevenson Ridge area which includes the Boulevard Project area. This section summarizes the current understanding of the geology in the project area.



Boulevard is underlain by 3 main rock units; the Dawson Range Batholith, the Klondike Schists and the Coffee Creek plutonic suite. The majority of the project is underlain by a west-northwest trending belt of volcanic, hypabyssal and sedimentary rocks, the Klondike Schist assemblage. This assemblage is composed of local variations of schist including variably pyritic chlorite-biotite-actinolite schist, amphibolite, variable micaceous quartzite, biotite feldspar-augen schist and quartz-augen metamorphic rocks. These rocks have been metamorphosed to a greenschist facies as identified by the presence of chlorite, biotite and actinolite assemblages within the package (McKenzie et al, 2012). This late Paleozoic aged assemblage has a number of age dates ranging from 249.6 to 267.2 Ma (McKenzie et al., 2012).

The southern portion of the project is underlain by the Dawson Range Batholith, a medium to coarse grained biotite-hornblende granodiorite of mid-Cretaceous age (dated at 100.2 to 102 Ma). This rock unit is characterized by its massive to lineated nature and locally aligned hornblende and elongated quartz crystals. The intrusive Dawson Range Batholith is separated from the Klondike schists to the north by a major crustal-scale thrust fault that is marked by minor lenses of ultramafic harzburgite along the fault's surface trace (McKenzie et al, 2012). The third main rock unit, the Coffee Creek plutonic suite, intrudes metamorphosed mafic schists in the central portion of the project area. The Coffee Creek plutonic suite is composed of medium to coarse grained biotite granite with minor pegmatitic phases. A minor garnet-bearing, quartz-pyritic porphyry phase has been identified at the western end of the Coffee Creek pluton (McKenzie et al, 2012). This intrusive has been age dated at 99.4 Ma +/- 0.9 (McKenzie et al, 2012).

McKenzie et al. (2012) identified five major vein types and developed a paragenetic sequence for the veins in the Sunset-Sunrise area based on cross-cutting relationships observed in core drilling. The first two stages (V_1 and V_2) contain quartz, iron carbonate, chalcopyrite, +/- pyrrhotite. The gold bearing event is V_3 , and these veins are comprised of quartz, iron carbonate, pyrite, arsenopyrite, gold and varying concentrations of sphalerite, galena, stibnite and tetrahedrite. V_3 veins are typically 2-20 mm wide and occur in sheeted sets that cross cut the metamorphic fabric at a very high angle. Geometric reconstructions of vein orientations using the metamorphic fabric yield a southeast strike and dips of 30° to the southwest. The final two vein stages (V_4 and V_5) are comprised of variable amounts of quartz and/or iron carbonate and V_4 contains pyrite.

A number of structural studies have been completed on the project (McKenzie et al., 2012; Sanchez and Ciolkiewicz, 2012, and Cooley, 2015a and 2015b). The following summary is a synthesis of these studies:

- Northeast verging thrust nappes interleave and juxtapose the metamorphic rock package. These thrust faults are sub-parallel with a penetrative foliation (striking east-southeast and dip steeply to the south). The foliation is interpreted as S_2 and is axial planar to isoclinal F_2 folds that are rarely observed. The F_2 fold axes plunge shallowly $10-20^\circ$ towards the ESE.
- NNW-SSE trending dextral fault zones offset the NNW trending thrust nappes. Seven major structures were identified on the basis of up to 800 m of offset of the ESE trending magnetic domains (i.e. stratigraphic horizons) but other, similar minor structures are also recognized. These structures are interpreted to postdate the NE directed thrusting.
- Rarely observed NW to NE-trending linear magnetic anomalies locally cross the WNW-ESE structures. Cooley (2015a) interpreted an increased concentration of these northerly trending structural features in the Sunrise area based on the 2015 ground magnetic data. Cooley attributed these linear magnetic highs to mafic dykes emplaced along northerly trending fault structures.

A limited amount of geological mapping was completed in the Denali area in 2015 (Cooley, 2015b). The observed rock types include quartzite, biotite schist, quartz biotite schist and chlorite schist. A quartz-eye feldspar porphyry dyke appears to intrude all of the metamorphic rocks, and an ultramafic unit is interpreted based on elevated Ni, Cr and Mg values, both in soils and in samples from drill hole YCS15-04. A well developed, penetrative foliation was observed in all outcrops (except the porphyry), and this foliation strikes 306° and dips 70° to the northeast. This fabric is weakly folded by a set of open folds that plunge 30° towards 323° and locally cause the foliation to dip to the southwest.

GEOPHYSICS

Two ground magnetic surveys were completed in August, 2015 to help with targeting for the fall drilling campaign. The Sunrise Zone survey was completed in three days (August 23-25) and included 48 line km (Figure 4). The Denali survey was completed in two days (Aug 26-27) and included 25 line km (Figure 5). A walking magnetic unit was utilized which collected continuous readings along lines spaced 50 m apart. Raw data from the survey is provided in Appendix 1.


DRILLING

During the 2015 programs, 21 reverse circulation drill holes were completed from 21 different drill sites. The summer drill program ran from June 15th to July 4th and included eight holes (819.92 m) at the Sunset-Sunrise target area and five holes (929.65 m) at the Denali target (Figure 3). The follow-up fall drill program extended from August 25th to September 11th and targeted the Sunrise area with an additional eight drill holes (1092.10 m). Collar locations and survey information are provided in Table 2 and drill logs are in Appendix 2. Intervals of geochemically significant mineralization are summarized in Table 3. The locations of all drill holes are shown on Figures 6 and 7, and are summarized below. A small representative sample of each drilled interval was retained in chip trays and these were transported to a storage facility in Vancouver, BC. There are no drill chips retained on site.

SUNSET-SUNRISE ZONE DRILLING

BV15-28 and BV15-29

These two holes are 250 m apart and tested the same northeast-trending magnetic lineament. Each drill hole tested significant multi-element gold (Au), arsenic (As) and antimony (Sb) soil geochemical anomalies. BV15-28 was drilled towards the southeast and intersected mostly quartz biotite schist with intercalated quartzite bands 1.5 to 4 m wide. BV15-29 only intersected quartz biotite schist. No significant gold values, alteration, veining or sulphide mineralization was intersected in these two drill holes.






Table 2 - Drill Hole Summary Data

Hole ID	Target/Objective	Easting NAD83-Z7	Northing NAD83-Z7	Elev (m)	Depth (m)	Azim	Dip
BV15-28	NE trending magnetic low/Soil anomaly	576132	6966800	1305	124.97	126	-55
BV15-29	NE trending magnetic low/Soil anomaly	576044	6966560	1331	102.11	310	-55
BV15-30	NE trending magnetic low/Soil anomaly	576544	6966510	1304	60.96	130	-55
BV15-31	NE trending magnetic low/Soil anomaly	576732	6965912	1240	70.1	90	-55
BV15-32	NE trending magnetic low/Soil anomaly	577247	6966447	1373	135.64	280	-55
BV15-33	NE trending magnetic low/Soil anomaly	577121	6966691	1294	102.11	110	-55
BV15-34	Magnetic low	577291	6967390	1260	124.97	115	-55
BV15-35	Scissor hole to BV15-31 (discovery hole)	576805	6965912	1247	169.16	272	-50
BV15-36	BV15-31 follow-up	576809	6965935	1252	170.69	270	-50
BV15-37	BV15-31 follow-up	576811	6965885	1243	140.21	270	-50
BV15-38	BV15-31 follow-up	576789	6965812	1230	132.59	270	-50
BV15-39	Discrete soil anomaly	576607	6965995	1250	156.97	110	-50
BV15-40	Soil geochem anomaly and magnetic low	576669	6965862	1227	105.16	119	-50
BV15-41	Soil geochem anomaly and magnetic low	576680	6965601	1179	50.29	120	-50
BV15-42	Soil geochem anomaly	576805	6965692	1211	167.03	300	-50
TG15-01	Depth potential below 2012 trench	576983	6966109	1287	99.06	90	-55
YCS15-01	Mag low, soil anomaly	566541	6975673	1223	169.17	242	-55
YCS15-02	NNW trending soil anomaly, ENE trending mag low	566497	6975699	1212	201.17	205	-55
YCS15-03	NNW trending soil anomaly	566659	6975604	1217	167.64	205	-55
YCS15-04	Perpendicular to easterly trending soil anomaly	566300	6975498	1218	199.64	335	-55
YCS15-05	Perpendicular to easterly trending soil anomaly	566102	6975500	1220	192.02	155	-55

Table 3 - 2015 Significant Assay Intervals

Hole ID	From (m)	To (m)	Length (m)	Au (g/t)
BV15-28			NSR	
BV15-29			NSR	
BV15-30	47.24	50.29	3.05	0.20
BV15-31	41.15	65.53	24.38	3.77
Incl	41.15	53.34	12.19	7.23
TG15-01	44.2	48.77	4.6	0.22
BV15-32	9.14	12.19	3.05	0.38
BV15-32	68.58	70.1	1.52	0.12
BV15-32	79.25	80.77	1.52	0.17
BV15-33	24.38	25.91	1.53	0.63
BV15-34			NSR	
BV15-35	45.72	68.58	22.86	0.94
Incl	53.34	54.86	1.53	2.78
BV15-36	57.91	67.06	9.15	0.40
Incl	57.91	59.44	1.53	1.04
BV15-37	73.15	97.54	24.39	0.26
BV15-38			NSR	
BV15-39	74.68	80.77	6.09	0.63
BV15-40	12.19	30.48	18.29	2.78
Incl	24.38	27.43	3.05	15.0
BV15-41			NSR	
BV15-42	3.05	4.57	1.52	2.10
BV15-42	91.44	96.01	4.57	1.63
YT15-01	18.29	21.34	3.05	3.11
YT15-02	71.63	74.68	3.05	0.5
	185.9	187.5	1.52	1.3
YT15-03	9.14	15.24	6.1	4.25
Incl	13.72	15.24	1.52	12.1
YT15-04			NSR	
YT15-05			NSR	





Figure 6: Drill Location Map for the Sunrise-Sunset Zones



LEGEND

- | | | | |
|--------------------------------|--------------------------|--|--|
| Reverse Circulation Drill hole | Gold 90th percentile | Metavolcanic/Amphibolite/
Feldspar Augen Schist | Ultramafic/Serpentinized
Hazburgite |
| Claim boundary | Arsenic 90th percentile | Undivided biotite-rich
metasediments | Feldspar Porphyry |
| 2012 Trench | Antimony 90th percentile | Quartz Biotite Schist | Metaplutonic/Leucogranite/Qtz-
Fspar Augen Schist |
| Dextral Fault | Thrust Fault (Inferred) | Thrust Fault | |

Figure 7: Drill Location Map for the Denali Zone

BV15-30

This drill hole is located on the eastern edge of a multi-element (Au-As-Sb) soil geochemical anomaly along a northeast-trending magnetic lineament. The geology is comprised of quartz biotite schist for the entire drill hole. A low grade intercept of 0.20 g/t over 3.1 m was intersected starting at 47.24 m down hole. This mineralization is associated with elevated As and Sb, combined with an increase in quartz veining (up to 65%) and pyrite (up to 5%).

BV15-31

This hole tested an area of northeast-trending magnetic lineaments that coincide with northeast-trending multi-element Au, As and Sb anomalies. The uppermost 30 m is comprised of discontinuous chlorite schist that changes to a mixed sequence of quartz biotite schist and quartzite. Starting at 41.15 m depth there is a mineralized zone that returned 3.8 g/t Au over 24.38 m (Table 3). Within this broad zone, a higher grade interval returned 7.23 g/t Au over 12.2 m. An interval of quartzite 16.77 m wide hosts the mineralization. The mineralized zone is recognizable in the rock chips by an increase in quartz veining (veins variably comprise from 0 to 30%), an increase in silicification (locally up to intense pervasive alteration) and an increase in pyrite (locally up to 12%). The drill hole ended in mineralization at 70.1 m depth.

TG15-01

This hole is located 320 m to the northeast from BV15-31, and tested a northeast-trending magnetic lineament which is coincident with an extensive As and Sb-in-soil anomaly and a discrete Au-in-soil anomaly. This drill hole has the identifier prefix “TG” that is different from the other holes in the Sunset-Sunrise area, because it was drilled on the Tiger (TG) property. TG15-01 intersected a sequence of quartzite and quartz biotite schist, alternating on 1.5 to 10 m scale. Low grade mineralization (<100 ppb Au) was returned discontinuously over the entire hole and increased to 216.8 ppb Au over 4.6 m, starting at 44.2 m depth. Quartz veins were variably developed in the upper 30 m of the drill hole, ranging up to 35% of the rock volume. Local, weak chlorite and biotite alteration occurs throughout the hole and pyrite varies from absent to 4%.

BV15-32

This drill hole is positioned 425 m northeast of TG15-01. It is situated on the southeastern margin of a robust As-in-soil anomaly and is coincident with a discrete 353 ppb Au-in-soil

anomaly. A mixed sequence of quartz biotite schist, interlayered with thin quartzite horizons (typically <3 m but locally up to 16 m wide) was intersected. Three low grade mineralized intervals are all hosted within quartzite; these have a maximum width of 3 m (Table 3). These three intervals are associated with subtle increases in both pyrite mineralization and quartz veining. Alteration is not significant.

BV15-33

This drill hole is situated 275 m north-northwest of BV15-32. It tested a northeast trending magnetic anomaly and is on the northern margin of an As-in-soil anomaly with weakly elevated gold. The hole intersected quartz biotite schist with thin (1-9 m) intervals of quartzite. A single interval returned 631 ppb Au over 1.53 m starting at 24.38 m down hole, and is associated with increased quartz and calcite veining. There is no associated alteration and only minor disseminated pyrite.

BV15-34

This drill hole is located 720 m north of BV15-33 and tested a break in the magnetic data near the southern margin of the Coffee Creek granite. The hole collared in massive, unfoliated granite and at 6.1 m depth intersected interbanded quartz biotite schist and quartzite. A continuous, 33.5 m wide, interval of chlorite schist extends down hole from 47.24 m depth. No significant mineralization was intersected.

BV15-35

This is the first drill hole of the fall drilling program, and was designed to follow-up the high grade intercept from discovery hole BV15-31. BV15-35 was collared 72 m due east of BV15-31 and drilled westwards, scissoring across BV15-31 in an attempt to determine the full width of the mineralized zone. The hole intersected alternating bands of quartz biotite schist and quartzite, with the quartzite ranging from 1.5 to 36.57 m in apparent thickness. The best gold grades were hosted within quartzite. A 22.9 m long interval, starting at 45.72 m depth, returned 0.94 g/t Au, with a high of 2.78 g/t Au over 1.52 m (Table 3). This mineralization is associated with increased quartz veining, locally up to 80% but more typically averaging 5-10% veins, weak to moderate pervasive silicification and 0.5-2% disseminated pyrite. Based on the mineralization intersected in this hole and in BV15-31, the mineralized zone is interpreted to be sub-vertically dipping and approximately 20 m wide. A second interval in the lowermost 32 m of the drill hole returned discontinuous mineralization ranging between 100-400 ppb Au associated with

reduced As and Sb, and a complete absence of an elevated base metal (Cu, Zn, Pb) signature. The contrasting pathfinder element signature suggests that the lower mineralized interval could be geochemically distinct from both the upper mineralized zone, and also distinct from the mineralization within hole BV15-31. This lower mineralized zone might be geochemically related to the mineralized zone in hole BV15-40 described below.

BV15-36

This hole is a 25 m step out to the north of BV15-31 and was drilled to the west to test the full extent of the interpreted northeast trending structure. The hole intersected quartz biotite schist interlayered with quartzite. The majority of the >100 ppb Au is hosted within the quartzite horizons which range in apparent thickness from 1.5 to 13.7 m. A 9.15 m wide interval that returned 0.40 g/t Au starts at 57.91 m depth. Lower grade gold intervals occur in the other quartzite layers. The mineralization is sometimes associated with a small increase in quartz veining, weak, discontinuous silicification and 1-2% disseminated pyrite. The discontinuous mineralization occurs within a sub-vertical zone that appears to be northeast trending, when correlated with the mineralized intervals in BV15-31 and BV15-35.

BV15-37

This hole is a 25 m step out to the south of BV15-31 and was inclined to the west, to test the extent of the interpreted northeast trending structure. The rocks in the upper half of the hole are dominated by quartz biotite schist with minor quartzite layers and this relationship is reversed in the lower half of the drill hole. The elevated gold grades in the lower half of the hole are associated with quartzite. One interval returned 0.24 g/t Au over 24.39 m, starting at 73.15 m depth. Another mineralized interval returned 0.50 g/t Au over 6.1 m, starting at 131.06 m depth. Both of these intervals are marked by an increase in quartz veining and weak to moderate, pervasive silicification. These two intercepts help to define a subvertical, northeast-trending, discontinuous structure that is up to 35 m wide, and open to the north. This mineralized structure is closed off 70 m to the south by hole BV15-38.

BV15-38

This hole is a 100 m step out to the south from BV15-31, and in combination with BV15-40, tests the southern extent of an interpreted northeast-trending mineralized zone. Approximately 15 m of quartzite occurs in both the top and bottom of the hole. These thick quartzite bands are

separated by quartz biotite schist with only thin interbands of quartzite. No significant mineralization was returned from this drill hole.

BV15-39

This hole is located 150 m northwest from BV15-31, and was positioned to test a northeast trending gold-in-soil anomaly (distinct from the anomaly tested by hole BV15-31). Similar to the other drill holes in the Sunrise area, this hole intersected a mixed sequence of quartzite and quartz biotite schist, with quartzite dominating the uppermost 33.5 m, and also the lowermost 18.3 m. A single quartzite horizon returned 6.1 m grading 0.63 g/t Au, starting at 74.68 m depth. Mineralization is associated with an increase in quartz veining (up to 5%), weak to moderate pervasive silicification and 1-2% disseminated pyrite. A sub-vertically dipping, northeast-trending structure is interpreted to cross-cut this quartzite.

BV15-40

This hole is located 80 m southwest of BV15-31 and tested an area with elevated gold, arsenic and antimony in soils, as well as interpreted northeast-trending structures. The rocks are comprised of a mixed sequence of quartz biotite schist with quartzite. A 3.1 m wide interval starting at 24.38 m depth returned 15.00 g/t Au. This intercept contains a high grade sample that returned 27.90 g/t Au over 1.52 m, and occurs within a broad, lower grade mineralized envelope that returned 2.78 g/t Au over 18.29 m. The high grade 3.1 m interval is in marked contrast to the other mineralized zones in the Sunrise Zone because it has no quartz veins, does not have any pervasive silicification nor associated sulphides. Quartz veins increase to 5% in the lower grade mineralized halo, above and to the west of the high grade interval. The pathfinder element geochemical signature also contrasts with that of the other mineralized intervals at Sunrise. There is comparatively lower arsenic, no elevated copper, only weakly elevated zinc, irregularly elevated lead and no sulphur reflecting the absence of sulphides with the gold in BV15-40. This is similar to the mineralization intersected at the western, deeper end of BV15-35 (and therefore interpreted to be part of the same northeast trending zone), but different from the mineralized zones intersected in BV15-31 and in the upper part of hole BV15-35 described above.

BV15-41

This hole is 315 m south of BV15-31 and tested a northeast-trending magnetic discontinuity, as well as a weakly elevated gold-in-soil anomaly. The rocks are dominated by quartzite with

intervals up to 12.2 m wide of quartz biotite schist. No significant mineralization was intersected. This hole was designed to test the area along trend to the south from the interpreted structures hosting mineralization in BV15-31. This drill hole only reached a final depth of 50.29 m (versus a target depth of 200 m) because of bad ground conditions, therefore this target has not been fully tested.

BV15-42

This hole is collared 230 m southeast from BV15-31 and tested northeast-trending magnetic discontinuities, multi-element soil anomalies, and the possible extension of mineralization in historical diamond drill hole BV08-07 which returned 1.69 g/t Au over 1.8 m. The rocks intersected are dominated by quartz biotite schist with interlayered horizons of quartzite up to 10 m thick. Two mineralized intervals were intersected: a single sample returned 2.10 g/t Au over 1.52 m starting at 3.05 m depth, and 1.63 g/t Au over 4.57 m starting at 91.44 m depth. The upper, thinner mineralized interval is hosted within quartz biotite schist while the wider interval is hosted within quartzite with an associated significant increase in quartz veining (up to 80%), strong pervasive silica alteration and disseminated pyrite up to 3%.

DENALI ZONE DRILLING

YCS15-01

This hole is collared 30 m northeast from the mineralization discovered in YCS-Trench 1 (4.56 g/t Au over 10 m; trenched in 2012) and was planned to test this mineralization at depth, as well as a robust multi-element soil anomaly (Figure 7). The hole intersected a sequence of quartz biotite schist, quartzite, chlorite schist and argillite. A single zone of mineralization returned 3.1 g/t Au over 3.05 m, starting at 18.29 m depth. The mineralization is hosted within quartz biotite schist that does not contain any quartz veins or silica alteration, and shows traces of fine grained disseminated pyrite. Rare arsenopyrite needles explain the elevated arsenic within the mineralized interval. This intercept is interpreted to be the down dip extension of the gold mineralization observed in the surface trench and therefore defines a shallow, northeast-dipping (~35°) mineralized zone. Mineralization is open, both at depth and along strike.

YCS15-02

This hole was collared 65 m northwest of the mineralization in YCS Trench-1 and was designed to test the north-central part of the Denali multi-element soil anomaly, the northwestern extension of the mineralization observed in the surface trench, as well as a northeast-trending

magnetic anomaly. The rock in the top 53 m of the drill hole is comprised of quartz biotite schist with minor quartz muscovite schist. The lower 148 m is dominated by quartzite with minor quartz biotite schist. Two mineralized zones were intersected; the first starts at 71.15 m depth and returned 0.5 g/t Au over 3.05 m, while the second interval is from a single 1.52 m sample that returned 1.30 g/t Au. Neither of these intervals have any associated quartz veins or alteration. The lower interval is marked by a subtle increase in disseminated pyrite, from absent to 1-2%. Both intervals are hosted within quartzite.

YCS15-03

This hole was collared 135 m southeast of the mineralization in YCS Trench-1 and was planned to test the southeastern extent of the Denali multi-element soil anomaly. The rocks are comprised of an interlayered sequence of quartz biotite schist and quartzite. A 6.1 metre wide zone grading 4.25 g/t Au was intercepted, starting at 9.14 m depth. With no quartz veining or silicification, the only visible evidence of mineralization is trace, disseminated fine grained pyrite and a subtle decrease in the oxidation intensity within the two sample intervals that contain the higher grade gold. This mineralized interval is interpreted to be part of the same shallowly northeast-dipping mineralized zone intersected in YCS15-01. Mineralization remains open at depth and along strike.

YCS15-04

This hole was collared 275 m southwest of the mineralization in YCS Trench-1 and was designed to test a northeast trending magnetic anomaly, coincident with a multi-element soil anomaly. The rocks are dominated by a sequence of quartz biotite schist with minor interlayered quartzite and chlorite schist. No significant mineralization was intersected.

YCS15-05

This hole was collared 450 m southwest of the mineralization in YCS Trench-1 and was planned to test a northeast trending magnetic anomaly, coincident with a multi-element soil anomaly. The rocks are dominated by broad intervals of quartz biotite schist within minor horizons of chlorite schist and quartzite. No significant mineralization was intersected.

SAMPLING PROCEDURES

For each 1.52 metre interval (5 feet), the reverse circulation drill produces approximately 15-25 kilograms of typically dry rock chips (chips measure up to 20 mm diameter) collected in a bucket. A representative sample for analysis was either collected using a splitter (Figure 8) or else homogenized directly within the bucket. A 2-5 kg representative sample for analysis was collected from each 1.52 m interval. Samples were submitted to SGS Canada Inc.'s laboratory facility in Vancouver, an ISO 17025 certified facility.

Upon arrival at the laboratory, rock chip samples were weighed and dried. Each sample was then crushed to 75% passing 2 mm, split into a 250

gram (g) sample that was pulverized to 85% passing 75 microns. A 0.25 g (Inductively Coupled

Plasma - ICP) and 30 g (fire assay- FA) subsample was obtained from the resulting material for analysis. Samples were analysed by aqua regia digestion and inductively coupled plasma with optical emission spectroscopy (ICP-OES) analysis for 34-elements. Gold was analysed by fire assay and atomic absorption spectroscopy (FAA313). If greater than 10,000 ppb Au was returned, a second analysis was completed using lead fusion fire assay and gravimetric finish. For those samples containing greater than 1 g/t Au, a metallic screen analysis was completed to determine the percentage of coarse gold in each sample. Assay certificates of analysis, laboratory certification and analytical method summaries are presented in Appendix 3.



Figure 8 Splitter used to obtain 2-5 kg representative sample.

QUALITY ASSURANCE/QUALITY CONTROL

For Quality Assurance-Quality Control (QA-QC) purposes, blanks (CDN-BL-10, purchased from CDN Resource Laboratories) were inserted on every sample identification number ending in 20, 50 and 80; while a gold standard (CDN-GS-5R, purchased from CDN Resource Laboratories) was inserted on every sample identification number that ended in 30, 60 and 90. The values

from these blanks were used to check for contamination between samples, and gold standards were used to check the consistency of the analytical procedures used by SGS Canada Inc. Erroneous QA-QC results were investigated and appropriate re-analysis undertaken when necessary. Internal SGS Canada Inc. blanks, duplicates, standards and spikes were also used to confirm the accuracy of the analytical methods and instruments.

Two charts below summarize the results from the QA-QC sampling. QA-QC samples passed without any significant concerns. Only seven blank samples returned a result above detection and between 5-12 ppb which is less than 2.5 times the detection limit (Figure 9). All the gold standards returned gold values less than or greater than 2 standard deviations from the expected value (Figure 10). As a result no QA-QC failures were reported.




Figure 9 - Blank Gold Values from the 2015 drilling programs.




Figure 10 - Gold standard values from the 2015 drilling programs. The expected gold standard value, along with the 1st and 2nd standard deviation values are indicated.

DISCUSSIONS AND CONCLUSIONS

A total of 1,912.02 m were drilled at the Sunset and Sunrise Zones in sixteen RC drill holes. A total of 929.65 m were drilled at the Denali Zone in five RC drill holes. A new discovery, the Sunrise Zone, was made in the Sunset target area, and gold mineralization was confirmed in bedrock below Trench 1 on the Denali Zone. Both Zones are summarized below.

SUNSET-SUNRISE ZONE

The 2015 exploration program was designed to test coincident northeast-trending magnetic anomalies and multi-element soil anomalies in the Sunset-Sunrise Zone area. Drill holes were designed to cross the target anomalies orthogonally. Lower grade intersections were returned from a number of these drill holes, but BV15-31 was the discovery hole at the Sunrise Zone. The fundamental control on the gold mineralization is the presence of quartzite. Quartzite is the host of, or directly proximal to (1's of metres away), the majority of the significant (>100 ppb Au) mineralization at the Sunrise Zone. This quartzite is part of a metamorphosed package of rocks that strike east-southeast (100°-110°) and dip steeply 75°-90° towards the southwest or northeast. Seven distinct quartzite horizons have been interpreted from drilling results; these

horizons are comprised of quartzite with minor intercalated quartz biotite schist (Figure 6). The Sunrise Zone area is distinct; it has the highest percentage of quartzite within the metamorphic rock package, when compared to the Sunset and Vegas trend areas located further to the west. However, there are numerous intervals of quartzite in the Sunrise Zone area that are not mineralized with gold. The gold mineralization is interpreted to be associated with later, cross-cutting structures, and gold mineralization is focused where these structures cross-cut the favourable quartzite. These structures are interpreted to strike north-northeast based on the following reasons:

1. McKenzie et al.'s work (2012) on the paragenetic sequence of quartz veins using drill holes from the 2008 and 2010 diamond drilling programs in the Sunrise-Sunset areas concluded that there is only one set of gold-bearing veins (V_3) and these veins cross cut, and therefore post-date S_2 , the main metamorphic fabric. There are only two recognized structural events that post-date this primary fabric: the NNW trending dextral faults and the NNE trending faults.
2. Sanchez and Ciolkiewicz's (2012) structural report recognized a main set of quartz veins that strike towards 210° and dip 75° to the northwest.
3. The linear magnetic anomalies identified and tested by drilling around the Sunrise Zone consistently strike north and northeast. These anomalies may mark (mineralized) fault structures.
4. The mineralized intercepts in drill holes BV15-31, -35, -36 and -37 align to define discontinuous, north northeast-trending mineralized zones that range from 3 to 38 m wide.

Four north-northeast striking mineralized zones have been identified using drill sections (Figure 6). These mineralized zones are also variably associated with an increase in quartz vein content, an increase in silica alteration that ranges from weak to intense, and an increase in disseminated sulphides, most commonly pyrite. The widest of the four zones measures up to 38 m in true thickness, is discontinuous across strike and only carries significant mineralization where the structure crosses the favourable quartzite host rock.

At least two geochemically distinct episodes of significant gold mineralization have been identified at Sunrise based on variations in pathfinder element concentrations. Gold generally is associated with highly elevated values of arsenic together with an elevated suite of base metal

values. However, the high grade gold mineralization in BV15-40 between 12.19 m and 30.48 m depth has much less arsenic, in combination with decreased base metal values than are seen with the gold mineralization in hole BV15-31.

The two highest grade gold intercepts were returned from drill holes that were drilled from west to east. The main set of quartz veins measured on surface all dip steeply to the WNW so holes with azimuths directed towards the east appear to be better oriented to evaluate the true thickness of these mineralized systems. More drilling is required to confirm this relationship.

DENALI ZONE

The main focus of the 2015 RC drill program was to test below the 2012 discovery trench that returned 4.56 g/t Au over 10 m. This trench is centrally located within a 1,300 m long, arcuate, northwest trending multi-element soil anomaly. Three of the 2015 drill holes tested this anomaly. The mineralized zone encountered in the discovery trench is interpreted to occur in two of these drill holes, with YCS15-03 returning the best grades. The mineralized zone is difficult to recognize in the RC rock chips due to the complete absence of quartz veins or any recognizable alteration. The only visible evidence for the gold mineralization is the presence of disseminated, fine grained pyrite within the quartz biotite schist and a subtle change (minimal oxidation within the zone) in the normally gradational decrease in oxidation extending from surface to 18 m depth. A significant (up to 0.98%) concentration of arsenic, and increase in antimony with moderate increases in zinc and silver are associated with gold at Denali. YCS15-02 probably missed the mineralized zone because it was collared too far to the west, in the footwall of the zone (Figure 7). The zone likely dips shallowly to the northeast, and the surface trace of the mineralization projects to the east of the collar of YCS15-02.

The other two Denali holes, YCS15-04 and -05, drilled during the summer 2015 program tested the same northeast trending, subtle magnetic anomaly and two separate, robust multi-element soil anomalies, without intersecting significant mineralization. With the knowledge gained from holes YCS15-01 and YCS15-03, the known mineralized zone at Denali is interpreted to be northwest striking and shallowly northeast dipping. The multi-element soil anomalies that were tested by holes YCS15-04 and -05 may well be due to mineralized structures that parallel the known mineralized zone at Denali. These soil anomalies should be tested by holes inclined to the southwest, from beyond the northeastern side of the anomaly area. Airborne magnetic survey results show a northwesterly trending magnetic low along the eastern side of the

geochemical soil anomaly tested by hole YCS15-05. This magnetic feature could be the expression of a mineralized fault structure that would be parallel to the known mineralized zone at Denali, which is located 400 m northeast of the multi-element soil anomaly and the magnetic low feature.

RECOMMENDATIONS

The following recommendations can be made for the Sunrise-Sunset and Denali Zone areas.

SUNSET-SUNRISE ZONE

1. The majority of the 2015 drilling occurs within a magnetically subdued region that is underlain by a mixed package of quartzite and quartz biotite schist, with minor chlorite schist. Similar magnetic lows should be drill-tested because of their potential to be underlain by mineralized quartzite.
2. The 2015 drilling confirmed that a gold-in-soil anomaly is a robust indicator of bedrock gold mineralization. The majority of the gold mineralization intersected by 2015 drill holes occurs below a >90th percentile gold-in-soil anomaly. Future drilling should focus on untested gold-in-soil anomalies.
3. Geological mapping in the area identified a main set of quartz veins that strike 210° and dip 75° to the northwest. If this is the orientation of the mineralized veins, then drill holes inclined towards the east and southeast would best test the area.
4. The possible extensions of the known zones, both to the northeast and to the southwest, should be drill tested at Sunrise. The main northeast trending mineralized zone identified in holes BV15-31, BV15-35 and BV15-36 trends into an area with highly anomalous gold-in-soil values, 100 to 220 m north of BV15-36. This area should be drill-tested. The areas northeast and southwest of BV15-40 should also be drill tested. BV15-39 is the only drill hole within a separate northeast-trending structure which is open to the northeast and southwest. BV15-42 intersected mineralization that should be targeted along trend to the northeast, where the mineralized zone is interpreted to cross other quartzite horizons.
5. The Hollywood area is located to the northwest of the Sunset Zone and is host to a robust gold-in-soil anomaly. This anomaly was drill tested in 2010 with 4 drill holes, two of which (BV10-22 and BV10-23) intersected 19.2 metres of 0.36 g/t Au and 37.62 metres of 0.64 g/t Au, respectively. While no quartzite was described within the two

historical drill holes, this area is a potential target because of the robust gold-in-soil anomaly and the previous drill results.

6. The Vegas area is located north of the Sunrise Zone and has been tested by only two drill holes, both of which were drilled towards the northeast. This target area is defined by some of the most robust soil anomalies on the property, up to 1370 ppb Au. Mapping by Cooley (2015a) concluded that this area is underlain by a coarse grained feldspar augen gneiss and felsic gneisses and schists. These rock units are favourable hosts for gold mineralization, so the Vegas area also warrants additional drilling.

DENALI ZONE

1. The most robust, coincident and continuous soil geochemical anomalies are located along trend to the southeast from hole YCS15-03 at the new Denali Zone. YCS15-03 is 680 m from the eastern claim boundary; 3 to 5 drill holes are recommended to evaluate this area.
2. The area along strike to the north of hole YCS15-01 at Denali should also be evaluated, with 1 or 2 holes collared east and north of YCS15-02 and inclined to the west and southwest.
3. The down-dip potential of the Denali Zone should be tested by drilling east of YCS15-01 and YCS15-03, towards the property boundary. These proposed holes should be inclined to the southwest.
4. Two to four drill holes should be drilled in the vicinity of YCS15-04 and YCS15-05. These holes should be inclined to the southwest to test for mineralized zones that parallel the known Denali Zone. These proposed holes should be collared well to the east or northeast of the eastern margin of the target soil anomaly, to ensure that the anomaly is properly tested.

REFERENCES

- Baker, D. (2011), Silver Quest Resources Ltd. 2010 Geological, Geochemical, Geophysical and Diamond Drilling Report on the Boulevard Property, Whitehorse Mining District, Yukon, Assessment Report
- Carpenter, R., (2010a) Kaminak Drills New Yukon Gold Discovery: First Hole Intersects 17.1 g/t Au over 15/5 m, Kaminak Gold Corporation, Vancouver, New Release
- Carpenter, R., (2010b) Kaminak Makes Second Yukon Gold Discovery: Latte Zone Yields 1.08 g/t Au over 84 m, Kaminak Gold Corporation, Vancouver, New Release
- Cooley, M., (2015a). Structural Geology Interpretation of the Sunrise/Sunset/Toni Tiger area, Boulevard property, Yukon. Internal Company Report. 13p.
- Cooley, M., (2015b). Structural Interpretation of the Denali Area. Internal Company Report. 7p.
- Fleming, A., (2010) Kinross Gold Corporation Announces Proposed Acquisition of Underworld Resources Inc., Underworld Resources Inc, Vancouver, News Release
- Hart, J. R., Goldfarb, R., Lewis, L. L., and Mair, J. L. (2004) The northern Cordilleran mid-Cretaceous plutonic province: Ilmenite/magnetite-series granitoids and intrusion-related mineralization: *Resource Geology*, v. 54, p. 253-280.
- Minfile 115J 048 (2012) HANNA; Yukon Geological Survey, Mineral Occurrence
- Minfile 115J 050 (2012) BOULEVARD; Yukon Geological Survey, Mineral Occurrence
- Minfile 115J 051 (2012) GOLD HAWK; Yukon Geological Survey, Mineral Occurrence
- Minfile 115J 052 (2012) TONI TIGER; Yukon Geological Survey, Mineral Occurrence
- McKenzie G.G., Allen, M.M., Hart, C.J.R, and Mortensen, J.K., (2012) Mid-Cretaceous Orogenic mineralization: an example from the Independence Creek area, Yukon Gold Project Final Technical Report, Mineral Deposit Research Unit, University of British Columbia, p. 101.
- McKenzie G.G. (2014) Geology and mineralization at Independence Creek: Dawson Range, West-central Yukon Territory; M.Sc. thesis presented to the University of British Columbia.
- Roberts, M., and Baker, D. (2007), 2006 Geological and Geochemical Report on the Rimfire-Northgate Alliance; Stewart River area, Yukon, unpublished Company report, p. 53.
- Ryan, J.J., Zagorevski, A., Roots, C.F., and Joyce, N., 2014. Paleozoic Tectonostratigraphy of the Northern Stevenson Ridge area, Yukon; Geological Survey of Canada, Current Research 2014-4, 13p.
- Sanchez, M. and Ciolkiewicz, W., 2012. Field and Magnetometric Interpretations for Independence Gold's Boulevard Property, Yukon. Internal Company Memo.
- Smith, H. (2010), Assessment Report describing Soil Geochemical Sampling at the HAN Property, Archer, Cathro & Associates (1981) Limited.

STATEMENT OF QUALIFICATIONS

I, David Gale, PGeo, MSc, of 3358 Passaglia Place, Coquitlam, British Columbia, hereby certify that:

I am a graduate of the Memorial University of Newfoundland having obtained the degree of Bachelor of Science in Geology in 1994 and am a graduate of Queen's University in Kingston having obtained a Master's of Science degree in Geology in 1997..

I am a registered member of the Association of Professional Engineers and Geoscientists of British Columbia (#27366).


I have been continuously employed in the mineral exploration industry in Canada since 1994.

I am currently employed as a Senior Geologist, by Independence Gold Corp. Suite 1410-650 West Georgia Street, Vancouver, British Columbia, Canada, V6B 4N8.

I am the author of the report entitled "2015 Drilling Program on the Boulevard Project, Yukon"

I helped to manage, log rock chips and reviewed the geological work on site reported herein.

Dated this 16th day of December, 2015.



David Gale, PGeo, MSc

STATEMENT OF QUALIFICATIONS

I, Kendra A. Johnston, PGeo, BSc, of Suite 206-1550 Barclay Street, Vancouver, British Columbia, hereby certify that:

I am a graduate of the University of Victoria, British Columbia having obtained the degree of Bachelor of Science in Earth and Ocean Science and Geography, 2005.

I am a registered member of the Association of Professional Engineers and Geoscientists of British Columbia (#37719).

I have been continuously employed in the mineral exploration industry in Canada since 2005.

I am currently employed as a Project Geologist, by Independence Gold Corp. Suite 1410-650 West Georgia Street, Vancouver, British Columbia, Canada, V6B 4N8.


I am an author of the report entitled "2015 Drilling Program on the Boulevard Project, Yukon" dated Dec 16, 2015.

I helped to manage, log rock chips and reviewed the geological work on site reported herein.

Dated this 16th day of December, 2015.



Kendra A. Johnston, PGeo, BSc



STATEMENT OF EXPENDITURES

Boulevard Statement of Expenditure

	<u>Quantity</u>		<u>Rate</u>		<u>Cost</u>	
Rock Sample Analysis	1,941		\$65.00		\$126,165.00	
Geologist day(s)	76		\$500.00		\$38,000.00	
Camp hand and Cook day(s)	76		\$500.00		\$38,000.00	
Planning and reporting day(s)	20		\$500.00		\$10,000.00	
Driller day(s)	112		\$0.00		\$0.00	
Pilot and Engineer day(s)	76		\$0.00		\$0.00	
Camp Costs (per man day)	340		\$500.00		\$170,000.00	
Helicopter Hour(s)	120		\$1,735.00		\$208,200.00	
Helicopter Fuel (drums)	105		\$900.00		\$94,500.00	
Geophysical Contract	5		\$2,700.00		\$13,500.00	
Drilling Contract	28		\$9,800.00		\$274,400.00	
			Subtotal:		\$972,765.00	
			Supervision:	12%	\$116,731.80	
			Total:		\$1,089,496.80	
			Claims Worked:	23	\$47,369.43	per claim worked
Dates Worked: June 15 - July 4 and Aug 23 - Sept 11, 2015						
Work Completed by: Independence Gold Corp. and Ground Truth						

APPENDIX 1
GROUND GEOPHYSICAL
RESULTS
(See Attached CD)

APPENDIX 2

DRILL LOGS

Legend

CAS	Casing	ppm	parts per million
QBS	Quartz Biotite Schist	Ox	Oxidation 1-5 with 1: very weak, 2:Weak, 3: Moderate, 4: Strong and 5: Intense
QMS	Quartz Muscovite Schist	Qtz	Quartz
CS	Chlorite Schist	sx	Sulphides
QTZT	Quartzite	cm	Centimeter
GR	Granite	mm	Millimeter
ARG	Argillite	py	Pyrite
VN	Vein	po	Pyrrhotite
From, To	measured in metres	cpy	Chalcopyrite
ppb	parts per billion	aspy	Arsenopyrite
-1	Below Detection Limit	LB	Logged By
Qtz Vn	Quartz Vein (pct)		

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-28	0	1.52		CAS					No Sample		
BV15-28	1.52	3.05	Grey	QBS	DG	2		Mostly fine dust - very minimal chips. The ones that can be seen are qtz and biotite schist. ~20-30% biotite.	E862001	70	19
BV15-28	3.05	4.57	Grey	QBS	DG	2		Oxidation is minimal. Qtz is non descript, no sx or oxidation. Rock is almost black--very dark grey reflecting biotite on foliation planes. Strong foliation within the rock. Note: First drill rod inserted into hole at the end of this run.	E862002	-1	9
BV15-28	4.57	6.1	Grey	QBS	DG	2		Very minimal qtz grains, looking like vein material. 'Wk' chlorite alteration observed within qtz within schist.	E862003	-1	4
BV15-28	6.1	7.62	Grey	QBS	DG	2		Qtz grains are now absent. ~30-40% biotite within the schist. Oxidation of biotite visible.	E862004	11	5
BV15-28	7.62	9.14	Grey	QBS	DG	2		Biotite decreases ~15-20%. Select pieces are moderately oxidized, some show minimal to absent oxidation.	E862005	26	7
BV15-28	9.14	10.67	Grey	QBS	DG	2		Slightly more quartz material, chips larger in size and more blocky.	E862006	-1	12
BV15-28	10.67	12.19	Grey	QBS	DG	2		Slight increase in qtz vein material. No sx. Minimal oxidation. Strong foliation.	E862007	24	6
BV15-28	12.19	13.72	Grey	QBS	DG	2		Qtz vein minimal to absent. Oxidation is minimal to absent. Biotite ~20%.	E862008	-1	8
BV15-28	13.72	15.24	Grey	QBS	DG	2		"Salt + pepper texture" defined by mix of biotite and quartz. All grains are the same size ~2-4mm	E862009	-1	4
BV15-28	15.24	16.76	Grey	QBS	DG	2		Biotite content is increasing to ~35-40% of rock. Grey. Minimal oxidation, no qtz.	E862010	35	6
BV15-28	16.76	18.29	Grey	QBS	DG	2		Biotite ~35-40% of rock. Strongly foliated. No sx. No alteration. Minimal oxidation.	E862011	26	5
BV15-28	18.29	19.81	Grey	QBS	DG	2		Biotite ~15-15% of rock. Forms setps, sub mm size bands between qtz bands. Strongly foliated. No qtz vein material - all host rock. Minimal oxidation. Grey. 85% quartz.	E862012	12	6
BV15-28	19.81	21.34	Grey	QBS	DG	2		Biotite ~10-15% of rock. Comprises the septa that defines the foliation. Oxidation does not appear homogeneous; mainly observed within qtz bands in rock. Oxidation imparts a brownish orange to the qtz bands. Minimal oxidation, 85% quartz.	E862013	-1	-1
BV15-28	21.34	22.86	Grey	QBS	DG	2		Biotite 10-15%. 85% quartz. Strongly foliated rock. Minimal oxidation.	E862014	-1	7
BV15-28	22.86	24.38	Grey	QBS	DG	2		Biotite ~ 10-15%. Rare chips of qtz exhibiting an oxidized orange brown colour. Minimal oxidation. 85% quartz.	E862015	7	4
BV15-28	24.38	25.91	Grey	QBS	DG	2		Biotite increasing to 15-20%. Rare chips of qtz exhibiting an oxidized orange brown colour. Minimal oxidation. 85% quartz.	E862016	9	5
BV15-28	25.91	27.43	Grey	QBS	DG	2		Biotite ~10-15%. Single 1cm fragment of qtz vein with sugary texture. Oxidation appears to be reduced - still minimal. 85% quartz.	E862017	-1	5
BV15-28	27.43	28.96	Grey	QBS	DG	2		Biotite ~15-20%. Oxidation staying constant; only discolouring the qtz layers. Very subtle green discolouration in rare qtz foliation - could be developing chlorite alteration. 80% qtz.	E862018	-1	6
BV15-28	28.96	30.48	Grey	QBS	DG	2		Mimimal oxidation. 80% qtz.	E862019	-1	5
BV15-28	30.48	32	Grey	QBS	DG	2		Biotite 10-15%. Minimal oxidation. 85% qtz.	E862021	-1	5
BV15-28	32	33.53	Grey	QBS	DG	3		Biotite ~10%. Oxidation of qtz increases (moderate). 90% quartz. This run things started to speed up.	E862022	5	15
BV15-28	33.53	35.05	Grey	QBS	DG	3		Biotite ~10-15%. Increased percentage of yellow brown oxidized qtz grains. Does not appear to be vein material. Just qtz bands. Moderate oxidation. 85% qtz.	E862023	11	12
BV15-28	35.05	36.58	Grey	QBS	DG	2		Minimal oxidation	E862024	-1	7
BV15-28	36.58	38.1	Grey	QBS	DG	2		Minimal oxidation	E862025	-1	7
BV15-28	38.1	39.62	Grey	QBS	DG	3		See the slight increase in level of oxidation in the qtz grains. Biotite ~10-15%. 85% qtz	E862026	-1	9
BV15-28	39.62	41.15	Grey	QBS	DG	2		Biotite 10-15%, Qtz 85%. Oxidation decreases from previous run. Rock is maintaining strong foliation.	E862027	-1	9
BV15-28	41.15	42.67	Grey	QBS	DG	2		Biotite 10-15%, 85% qtz. Oxidation is still minimal.	E862028	-1	16
BV15-28	42.67	44.2	Grey	QBS	DG	3		Biotite 10-15%, qtz 90%. Colour is a lot lighter than previous reflecting decrease biotite and increase in qtz. Oxidation of qtz increases to moderate.	E862029	11	14
BV15-28	44.2	45.72	Grey	QBS	DG	2		Biotite 5-10%, 90% qtz. Oxidation has decreased since last interval (minimal).	E862031	-1	6
BV15-28	45.72	47.24	Grey	QBS	DG	2		Biotite is 15-20%, qtz 80%. Oxidation is still minimal.	E862032	-1	8
BV15-28	47.24	48.77	Grey	QBS	DG	2		Limonite in fine material, biotite 20%, 5% qtz. Minimal oxidation.	E862033	16	11
BV15-28	48.77	50.29	Grey	QBS	DG	1		5% muscovite, biotite 20/25%, 5% quartz. Oxidation = 1 (fresh rock).	E862034	-1	11
BV15-28	50.29	51.82	Grey	QBS	KJ	2		Biotite 20%, 10% qtz. Minimal oxidation.	E862035	-1	15
BV15-28	51.82	53.34	Grey	QBS	KJ	2		Biotite 25%, 15% qtz. Minimal oxidation.	E862036	50	14
BV15-28	53.34	54.86	Grey	QBS	KJ	2		Biotite 25%, 10% qtz. Minimal oxidation.	E862037	6	8
BV15-28	54.86	56.39	Grey	QBS	KJ	3		Oxidized quartz. Moderate oxidation. 20% quartz.	E862038	8	9
BV15-28	56.39	57.91	Grey	QBS	KJ	3		Limonite on quartz. Moderate oxidation. 20% quartz.	E862039	31	8
BV15-28	57.91	59.44	Grey	QBS	KJ	1		Qtz veinlets, pinkification. 45% quartz.	E862040	7	7
BV15-28	59.44	60.96	Grey	QBS	KJ	1		Biotite 25-30%, 15% qtz.	E862041	12	8
BV15-28	60.96	62.48	Grey	QBS	KJ	1		1cm qtz chips, muscovite. 20% qtz.	E862042	13	8
BV15-28	62.48	64.01	Grey	QBS	KJ	4		40% limonite, muscovite. 60% quartz. Moderate to intense oxidation.	E862043	-1	13
BV15-28	64.01	65.53	Grey	QBS	KJ	4		35% limonite, muscovite. Moderate to intense oxidation. 20% quartz.	E862044	-1	12
BV15-28	65.53	67.06	Grey	QBS	KJ	3		20% limonite, muscovite. 35% quartz. Moderate oxidation.	E862045	-1	9
BV15-28	67.06	68.58	Grey	QZT	KJ	4		85% quartz, biotite 15%. 85% oxidation. Moderate to intense oxidation.	E862046	-1	12
BV15-28	68.58	70.1	Grey	QZT	KJ	2		1.5cm quartz chips, 10% biotite. Minimal oxidation. 90% quartz.	E862047	-1	3
BV15-28	70.1	71.63	Grey	QZT	KJ	1		1cm quartz and smokey qtz, biotite 40%. 60% quartz.	E862048	-1	-1
BV15-28	71.63	73.15	Grey	QBS	KJ	1		Pinkification, 60% biotite, 40% quartz.	E862049	-1	4
BV15-28	73.15	74.68	Grey	QBS	KJ	1		70% biotite, 30% quartz.	E862051	-1	6
BV15-28	74.68	76.2	Grey	QZT	KJ	1		Hit water prior to this run. Veyr weak chlorite, 15% biotite, 80% quartz.	E862052	-1	4

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-28	76.2	77.72	Grey	QBS	KJ	0		Chlorite alteration, 60% biotite, 40% quartz.	E862053	-1	6
BV15-28	77.72	79.25	Grey	QBS	KJ	1		Weak chlorite, pinkification. 30% quartz.	E862054	-1	8
BV15-28	79.25	80.77	Orange	QBS	KJ	5		Fines turned from grey to orange. Intense oxidation. 30% quartz.	E862055	-1	4
BV15-28	80.77	82.3	Grey	QBS	KJ	2		Silicified, muscovite, limonite. Minimal oxidation. 30% quartz.	E862056	7	3
BV15-28	82.3	83.82	Grey	QBS	KJ	2		Minimal oxidation. 30% quartz.	E862057	-1	3
BV15-28	83.82	85.34	Grey	QBS	KJ	1		Chlorite and pinkification evident. 30% quartz.	E862058	-1	5
BV15-28	85.34	86.87	Grey	QBS	KJ	1		Chlorite and pinkification. 30% quartz.	E862059	-1	3
BV15-28	86.87	88.39	Grey	QBS	KJ	1		Pretty clean. 10% quartz.	E862061	-1	18
BV15-28	88.39	89.92	Grey	QBS	KJ	4		More powder, less chips produced. Limonite, qtz chips heavily oxidized. Moderate to intense oxidation, 40% quartz.	E862062	6	8
BV15-28	89.92	91.44	Grey	QBS	KJ	2		Sample more sandy than powder. Minimal oxidation, 20% quartz.	E862063	-1	7
BV15-28	91.44	92.96	Grey	QBS	KJ	1		Moist ground, some 1cm qtz, pitted qtz. 30% quartz.	E862064	-1	8
BV15-28	92.96	94.49	Grey	QBS	KJ	0		40% quartz.	E862065	-1	6
BV15-28	94.49	96.01	Grey	QBS	KJ	1		Oxidized biotite. 60% quartz.	E862066	-1	6
BV15-28	96.01	97.54	Grey	QBS	KJ	1		Almost a quartzite. 70% quartz, 30% biotite.	E862067	6	8
BV15-28	97.54	99.06	Grey	QBS	KJ	0		Very dark, 90% biotite, 5% quartz.	E862068	-1	6
BV15-28	99.06	100.58	Grey	QBS	KJ	0		Very dark, 80% biotite, 10% quartz.	E862069	-1	6
BV15-28	100.58	102.11	Grey	QBS	KJ	0		20% quartz, minimal potassic alteration. Possible Fe staining rather than potassic alteration.	E862070	-1	4
BV15-28	102.11	103.63	Grey	QBS	KJ	0		35% quartz. No alteration. Hit a soft spot before this sample, small cave in.	E862071	-1	6
BV15-28	103.63	105.16	Grey	QBS	KJ	0		20% quartz. Mainly small qtz chips must have gone through a qtz vein cm's thick, no alteration.	E862072	-1	7
BV15-28	105.16	106.68	Black	QBS	KJ	0		5% quartz, No oxidation, no alteration, black biotite	E862073	-1	5
BV15-28	106.68	108.2	Grey	QBS	KJ	0		30% quartz, no oxidation. Small pure qtz chips, likely from vein material.	E862074	-1	4
BV15-28	108.2	109.73	Grey	QBS	KJ	0		10% quartz, no oxidation. Qtz chips from mm-cm wide vein.	E862075	-1	6
BV15-28	109.73	111.25	Grey	QBS	KJ	0		40% quartz, no oxidation. No alteration.	E862076	-1	5
BV15-28	111.25	112.78	Grey	QBS	KJ	0		20% quartz. No alteration, no oxidation.	E862077	-1	7
BV15-28	112.78	114.3	Grey	QBS	KJ	0		30% quartz. No oxidation. Drilled through cm scale veins. No alteration.	E862078	-1	4
BV15-28	114.3	115.82	Grey	QBS	KJ	0		15% quartz. No oxidation.	E862079	-1	-1
BV15-28	115.82	117.35	Grey	QBS	KJ	0		5% quartz, no oxidation. Mostly biotite.	E862081	-1	6
BV15-28	117.35	118.87	Grey	QBS	KJ	0		45% quartz, no oxidation. Minimal chlorite alteration.	E862082	-1	-1
BV15-28	118.87	120.4	Grey	QBS	KJ	0		30% quartz, no oxidation. Water in hole, no alteration visible.	E862083	-1	3
BV15-28	120.4	121.92	Grey	QBS	KJ	0		40% quartz, no oxidation. Lots of water, sample really gucky, possibly some silicification.	E862084	-1	-1
BV15-28	121.92	123.44	Grey	QBS	KJ	0		70% quartz, no oxidation. Lots of water in hole, broken bit pulling out after this sample. This sample is 400-404', bull quartz chips, likely from vein.	E862085	-1	4
BV15-29	0	1.52		CAS					No Sample		
BV15-29	1.52	3.05	Grey	QBS	DG	2	5	~40% biotite. 5% quartz veins grains.	E862099	7	11
BV15-29	3.05	4.57	Black	QBS	DG	2	0	~50-60% biotite	E862100	11	23
BV15-29	4.57	6.1		QBS	DG	0	15	50% biotite, no oxidation. Qtz vein material smokey grey to white, ~15% of the interval. More clay in interval - getting banged up in the splitter	E862101	6	14
BV15-29	6.1	7.62		QBS	DG	0	40	50% biotite, 50% quartz. 40% of interval is qtz veins, white and massive.	E862102	-1	8
BV15-29	7.62	9.14	Grey	QBS	DG	2	15	60% quartz in rock and veins. Biotite ~40%. Qtz vein ~15%. Sugary medium to coarse grained qtz. No sx.	E862103	6	11
BV15-29	9.14	10.67		QBS	DG	0	5	Medium Grey. Biotite has dropped off dramatically ~15%. White qtz vein material still present. Weak potassic alteration possibly - massive as subtle pink colour of qtz. One large piece of rock can see a qtz vein 1mm thick perpendicular to the foliation in the rock.	E862104	-1	7
BV15-29	10.67	12.19	Grey	QBS	DG	0	10	30% biotite, 70% qtz. 10% qtz vein material. 1% py in wall rock, not veins. Dark grey. No oxidation.	E862105	9	7
BV15-29	12.19	13.72	Grey	QBS	DG	0	10	30% biotite, 70% qtz. Wall rock strongly foliated. 10% qtz vein frags. No oxidation.	E862106	18	7
BV15-29	13.72	15.24	Grey	QBS	DG	0	7	35% biotite, 65% qtz. Medium grey. Qtz vein 7% of rock. Strong foliation in host rock. No oxidation.	E862107	9	5
BV15-29	15.24	16.76	Grey	QBS	DG	0	15	25-30% biotite, 70% qtz. Qtz vein material increases ~15%. Medium grey. No other sign alteration. No oxidation. Sx in wall rock, not in veins.	E862108	5	5
BV15-29	16.76	18.29	Grey	QBS	DG	0	12	Medium grey. Biotite ~25%, Qtz ~75%. Qtz vein ~12%. No oxidatin - only very rare grains are weakly oxidized. Py in qtz vein and host rock.	E862109	-1	5
BV15-29	18.29	19.81	Grey	QBS	DG	0	15	Biotite 20%, Qtz 80% (in host rock). Qtz vein ~15% of sample. **0.5% Cpy observed along vein margin. Medium grey. No oxidation.	E862110	-1	7
BV15-29	19.81	21.34	Grey	QBS	DG	0	30	30% biotite, 70% qtz. Grey. Qtz vein comprises 30% of rock. White to light grey in colour. Essentially no oxidation. Single piece shows discolouring.	E862111	8	4
BV15-29	21.34	22.86	Grey	QBS	DG	0	7	Biotite 25%, Qtz 75%. Qtz vein decreases to 7%. No oxidation. Medium grey. Potassic alteration imparts a subtle brownish pink colour to the rock	E862112	7	8
BV15-29	22.86	24.38	Grey	QBS	DG	0	7	20% biotite, 80% qtz (in host rock). Qtz vein 7% of rock; typically white, medium to coarse grained. Medium grey, no oxidation.	E862113	-1	6
BV15-29	24.38	25.91	Grey	QBS	DG	0	25	20% biotite, 80% qtz. Medium grey. 25% qtz vein. No oxidation	E862114	6	9

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-29	25.91	27.43	Grey	QBS	DG	0	15	25% biotite, 75% qtz. 15% white and grey qtz vein medium to coarse grained. No oxidation. No potassic alteration evident.	E862115	-1	6
BV15-29	27.43	28.96	Grey	QBS	DG	0	3	20% biotite, 75% qtz. Medium grey. No oxidation. Qtz vein decreases to ~3%. No other alteration.	E862116	-1	3
BV15-29	28.96	30.48	Black	QBS	DG	0	2	50% biotite, 50% quartz. Black. No oxidation. 2% qtz veining. Looks like py is disseminated through rock and coating fractures.	E862117	-1	5
BV15-29	30.48	32	Grey	QBS	DG	0	5	Biotite 20%, Qtz 80%. Qtz veining 5%. No oxidation. Medium grey. Po appears to occur along margins of qtz vein.	E862118	-1	3
BV15-29	32	33.53	Grey	QBS	DG	0	12	Medium grey, no oxidation. 20% biotite, 80% Qtz. 12% of sample is qtz vein. Minimal potassic alteration--patches.	E862119	-1	-1
BV15-29	33.53	35.05	Grey	QBS	DG	0	3	85% qtz, 15% biotite. No oxidation. Medium grey. 3% quartz veining. Strong foliation best observed on dry surface	E862121	-1	5
BV15-29	35.05	36.58	Grey	QBS	DG	0	5	15% biotite, 85% qtz. 5% qtz veining - white, medium coarse grained. Strong foliation; pervasive.	E862122	-1	3
BV15-29	36.58	38.1	Grey	QBS	DG	0	3	15% biotite, 85% quartz, 3% qtz veining. No oxidation.	E862123	-1	4
BV15-29	38.1	39.62	Grey	QBS	DG	0	5	~20% biotite, 80% qtz. 5% quartz veining. No sx. No oxidation. Medium grey.	E862124	-1	20
BV15-29	39.62	41.15	Grey	QBS	DG	0	13	15% biotite, 80% qtz. 14% qtz calcite veins. Milky white calcite mixed with white grey qtz veins. No oxidation. Medium grey.	E862125	-1	8
BV15-29	41.15	42.67	Grey	QBS	DG	0	0	15% biotite. No oxidation. Medium grey. No veining. Trace pyrite, coating fractures.	E862126	-1	7
BV15-29	42.67	44.2	Grey	QBS	DG	0	3	15% biotite, 85% qtz. White coarse grained qtz with minor calcite veinlets. Medium grey, no oxidation.	E862127	11	8
BV15-29	44.2	45.72	Grey	QBS	DG	0	2.5	15% biotite, 85% qtz. 5% qtz-calcite veins. No oxidation. Dark grey. 0.5% py as fracture coatings, perpendicular to foliation and disseminated wisps within foliation.	E862128	6	6
BV15-29	45.72	47.24	Grey	QBS	DG	0	1.5	15% biotite, 85% qtz. 3% quartz calcite veinlets. No sx, no oxidation. Medium grey. Possibly very weak potassic alteration developing	E862129	-1	4
BV15-29	47.24	48.77	Grey	QBS	DG	0	0	20% biotite, 80% qtz. Dark grey. No oxidation. Wisp disseminated py +/- po within wall rock.	E862131	-1	11
BV15-29	48.77	50.29	Grey	QBS	DG	0	0	25% biotite, 75% qtz. No oxidation, no sx. Trace calcite vein parallel to foliation.	E862132	-1	4
BV15-29	50.29	51.82	Grey	QBS	DG	0	0	20% biotite, 80% qtz. No veining. Medium grey. No sx, no oxidation.	E862133	-1	6
BV15-29	51.82	53.34	Grey	QBS	DG	0	1.5	20% biotite, 80% qtz. 3% qtz and calcite veinlets. Medium grey. No oxidation. 2% py on margin of rare qtz vein.	E862134	34	5
BV15-29	53.34	54.86	Grey	QBS	DG	0	12	20% biotite, 80% qtz. 12% qtz vein with 1% calcite. 0.5%py disseminated as wisp in foliation.	E862135	-1	7
BV15-29	54.86	56.39	Grey	QBS	DG	0	12	25% biotite, 75% qtz. 12% qtz +/- calcite. No oxidation. Medium grey. 0.5% py in host rock.	E862136	-1	5
BV15-29	56.39	57.91	Grey	QBS	DG	0	7	25% biotite, 75% qtz. 7% qtz veining +/- calcite veinlets. No oxidation. No py.	E862137	-1	7
BV15-29	57.91	59.44	Grey	QBS	DG	0	0	15% biotite, 85% qtz. Medium grey. No vein. No sx. No oxidation.	E862151	-1	6
BV15-29	59.44	60.96	Grey	QBS	DG	0	0	15% biotite, 85% qtz. No vein. No oxidation. Medium grey. 1% py dissminated as wisps aligned in foliation.	E862152	-1	9
BV15-29	60.96	62.48	Grey	QBS	DG	0	0	15% biotite, 85% qtz. Medium grey. No vein. No sx. No oxidation.	E862153	-1	6
BV15-29	62.48	64.01	Grey	QBS	DG	0	1	15% biotite, 85% qtz. 1% qtz veining. No oxidation. Medium grey. 0.5% py within schist.	E862154	-1	7
BV15-29	64.01	65.53	Grey	QBS	DG	0	1	15% biotite, 85% qtz. 1% qtz veining. No sx. No oxidation. Very subtle brownish pink colouration of qtz--possibly representing potassic alteration. Only observed in a single piece.	E862155	-1	-1
BV15-29	65.53	67.06	Grey	QBS	DG	0	5	15% biotite, 85% qtz. No sx. No alteration. No oxidation. 5% qtz veining.	E862156	12	5
BV15-29	67.06	68.58	Grey	QBS	DG	0	20	15% biotite, 85% qtz. 20% qtz vein +/- calcite. 0.5% py.	E862157	9	5
BV15-29	68.58	70.1	Grey	QBS	DG	3	15	15% biotite, 85% qtz. Mixed grey and orangey brown. 15% veining - 10% qtz, 5% calcite. Oxidation = 3. Almost looks like different rock type but just increasing in oxidation and veining.	E862158	10	5
BV15-29	70.1	71.63	Grey	QBS	DG	3	18	15% biotite, 85% qtz. Orangey grey. Oxidation = 3. Calcite = 20%.	E862159	7	4
BV15-29	71.63	73.15	Grey	QBS	DP	2	5	red-orange lmnt on fracture surface. Oxidized.	E862161	-1	6
BV15-29	73.15	74.68	Black	QBS	DP	1	1	Black; 80% biotite; oxidation	E862162	-1	6
BV15-29	74.68	76.2	Grey	QBS	DP	0	1	Medium grey. 35% biotite. Qtzt bands 2-5mm.	E862163	-1	6
BV15-29	76.2	77.72	Grey	QBS	DP	0	2	Medium grey. 35% biotite. Qtz vein fragments to 5mm	E862164	-1	6
BV15-29	77.72	79.25	Grey	QBS	DP	1	0	Light grey 25% biotite, qtzt bands 2mm	E862165	-1	4
BV15-29	79.25	80.77	Grey	QBS	DP	1	1	Light grey 20% biotite, qtzt bands 2mm	E862166	-1	5
BV15-29	80.77	82.3	Grey	QBS	DP	1	1	Light grey 20% biotite, qtzt bands 2mm	E862167	-1	6
BV15-29	82.3	83.82	Grey	QBS	DP	1	1	Light grey 20% biotite, qtzt bands 2mm	E862168	-1	7
BV15-29	83.82	85.34	Grey	QBS	DP	2	3	Light grey 20% biotite, qtzt bands 2mm	E862169	-1	8
BV15-29	85.34	86.87	Grey	QBS	DP	3	1	Light grey 20% biotite, qtzt bands 2mm	E862170	7	12
BV15-29	86.87	88.39	Grey	QBS	DP	2	2	Light grey 20% biotite, qtzt bands 2mm	E862171	-1	8
BV15-29	88.39	89.92	Grey	QBS	DP	1	7	Light grey 20% biotite	E862172	-1	4
BV15-29	89.92	91.44	Grey	QBS	DP	1	2	Light grey 20% biotite	E862173	5	3
BV15-29	91.44	92.96	Grey	QBS	DP	0	0	Light grey 20% biotite	E862174	-1	5
BV15-29	92.96	94.49	Grey	QBS	DP	0	0	Medium grey	E862175	-1	4
BV15-29	94.49	96.01	Grey	QBS	DP	1	0.5	Medium grey	E862176	-1	4
BV15-29	96.01	97.54	Grey	QBS	DP	1	0	Medium grey	E862177	-1	7
BV15-29	97.54	99.06	Grey	QBS	DP	0	0	Medium grey	E862178	-1	7
BV15-29	99.06	100.58	Grey	QBS	DP	1	1	Medium grey	E862179	-1	6
BV15-29	100.58	102.11	Grey	QBS	DP	0	0	Medium grey qtz veinlet 2-5mm wide with blebs, po. First appearance of chlorite. Few pieces of qtzt.	E862181	-1	8
BV15-30	0	1.52		CAS					No Sample		

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-30	1.52	3.05	Grey	QBS	DG	0	0	Overburden to ~7 feet then bedrock started. Still a mixture of qtz vein, oxidized qtz and QBS so not entirely local material (i.e. Half overburden, half bedrock).	E862086	-1	9
BV15-30	3.05	4.57	Grey	QBS	DG	3	4	Biotite ~20%, 75% qtz. Moderate oxidation. ~3-5% vein material.	E862087	-1	4
BV15-30	4.57	6.1	Grey	QBS	DG	3	0.25	Biotite 20%, moderate oxidation. Trace qtz veins.	E862088	-1	4
BV15-30	6.1	7.62	Grey	QBS	DG	3	5	Biotite ~15%, 75% qtz. Qtz veins 5. Moderate oxidation.	E862089	-1	6
BV15-30	7.62	9.14	Grey	QBS	DG	2	1	Biotite ~15%, 84% qtz. Qtz vein 1%. Minimal oxidation.	E862091	-1	5
BV15-30	9.14	10.67	Grey	QBS	DG	2	3	Biotite 10%, 87% qtz. Qtz vein 3%. No sx, very strong well developed foliation. Oxidation is not pervasive - only select grains.	E862092	-1	5
BV15-30	10.67	12.19	Grey	QBS	DG	1	1	Biotite 10%, 89% qtz, qtz vein 1%. No sx.	E862093	-1	5
BV15-30	12.19	13.72	Grey	QBS	DG	0	3	Biotite 10%, 87% qtz, qtz vein 3%. Occurring in schist as euhedral crystals. Possible chlorite alteration developing as green discolouration of qtz bands.	E862094	-1	8
BV15-30	13.72	15.24	Grey	QBS	DG	2	5	Biotite 10%, 85% qtz, qtz vein 5%.	E862095	-1	6
BV15-30	15.24	16.76	Grey	QBS	DG	2	7	Biotite 10%, Qtz vein 7%, 83% qtz. Euhedral py forming fine grains 1mm in size. Also disseminated wisps in schist.	E862096	5	6
BV15-30	16.76	18.29	Grey	QBS	DG	3	15	Biotite 10%, 75% qtz, 15% qtz vein. Py observed in a qtz vein. Also possible pinhead size silver mineralization - sx. Moderate oxidation.	E862097	17	6
BV15-30	18.29	19.81	Grey	QBS	DG	3	60	Biotite 10%, 35% qtz. Qtz 90% of bedrock. Qtz vein ~60%. Moderate oxidation. Minimal chlorite alteration.	E862098	-1	4
BV15-30	19.81	21.34	Grey	QBS	DG	2	25	Biotite 10%, 65% quartz. Minimal oxidation. Qtz vein comprises 25% of rock--no sx in veins. Minor calcite with qtz veins.	E862200	30	11
BV15-30	21.34	22.86	Grey	QBS	DG	2	7	Biotite 10%, qtz veins 7%, 83% qtz. Minimal oxidation. Oxidation appears to occur along fractures. Minor calcite (~0.5%) along with qtz veins.	E862201	-1	8
BV15-30	22.86	24.38	Grey	QBS	DG	0	5	Biotite 10%, 85% qtz, qtz veins 5%. Py occurs disseminated within schist and within quartz veins. Subtle pinkish brown colour in qtz host rock bands representing weak potassic. Weak greenish colour to select qtz veins = chlorite.	E862202	-1	11
BV15-30	24.38	25.91	Grey	QBS	DG	2	10	10% biotite, 80% quartz, qtz veins 10%. Foliated qtz biotite schist. Note - water was hit in the run below, 85-90.	E862203	18	10
BV15-30	25.91	27.43	Grey	QBS	DG	2	15	Biotite 10%, Qtz veins 15%, 75% quartz. Py is coating the foliation planes and forming subhedral crystals on margins of veins in schist. Minimal oxidation.	E862204	7	5
BV15-30	27.43	28.96	Grey	QBS	DG	0	5	Biotite ~10%, qtz vein 5%, 85% quartz. Weak brownish pink colour to qtz in host rock.	E862205	-1	3
BV15-30	28.96	30.48	Grey	QBS	DG	0	10	Biotite 10%, qtz vein 10%. Pinkish brown colour of schist = potassic alteration. Chlorite alteration observed in green staining/discolouration of qtz vein. Rock continues to be strongly schistose.	E862206	-1	-1
BV15-30	30.48	32	Grey	QBS	DG	0	12	Biotite 10%, quartz 78%, qtz vein 12%. Py occurs as subhedral masses and crystals along vein margins as wisps aligned within foliation planes. Moderate potassic alteration.	E862207	-1	5
BV15-30	32	33.53	Grey	QBS	DG	0	3	Biotite 10%, Qtz vein 3%, 87% qtz. Potassic alteration pervasive, imparts a subtle pinkish brown colour. Moderate potassic alteration.	E862208	-1	3
BV15-30	33.53	35.05	Grey	QBS	DG	0	3	Biotite 15%, 82% qtz, 3% qtz vein. Potassic alteration is absent.	E862209	-1	4
BV15-30	35.05	36.58	Grey	QBS	DG	0	2	Biotite 15%, 83% qtz, qtz vein 2%. Minimal potassic alteration.	E862210	-1	6
BV15-30	36.58	38.1	Grey	QBS	DG	0	2	Biotite 15%, 83% qtz, qtz vein 2%. Minimal potassic alteration.	E862211	8	5
BV15-30	38.1	39.62	Grey	QBS	DG	0	2	Biotite 12%, Qtz vein 2%, 86% qtz. Chlorite observed in a single qtz vein mixed with py +/- po. Moderate potassic alteration. Minimal chlorite alteration and 2% py mineralization.	E862212	5	8
BV15-30	39.62	41.15	Grey	QBS	DG	0	1	Biotite 12%, qtz vein 1%, 87% qtz. Potassic alteration imparts a subtle brown colour to the qtz bands. Very minor oxidation of sulphides within one of the qtz veins. Potassic alteration is moderate, 2% py mineralization.	E862213	-1	9
BV15-30	41.15	42.67	Grey	QBS	DG	0	1	Biotite 7%, Qtz veins 1%, 92% qtz. Observed ~1mm bands of qtz separated by fine septa of biotite. Rare py is disseminated through the schist.	E862214	-1	4
BV15-30	42.67	44.2	Grey	QBS	DG	0	1	Biotite 7%, 92% quartz. Still a schist but texture is starting to look almost gneissic. 1-1.5mm thick bands of qtz separated by thin septa of biotite. ~1% qtz veins.	E862215	-1	5
BV15-30	44.2	45.72	Grey	QBS	DG	0	2	Biotite 10%, 88% quartz, qtz veining 2%. Minimal potassic alteration, 2% py mineralization. Py occurs as wisps in schist--disseminated, as well as subhedral crystals on margin of qtz vein.	E862216	-1	-1
BV15-30	45.72	47.24	Grey	QBS	DG	2	25	Biotite at 10% forming sub mm bands between 1-2mm thick qtz layers. Qtz vein: 25% of rock--some qtz veins exhibit a weak oxidation. Very weak calcite occurring within the qtz veins. 65% quartz. 0.5% py mineralization.	E862217	8	9
BV15-30	47.24	48.77	White	QBS	DG	0	65	Logged as Quartz Vein. 30% quartz, 2% carbonates. Minimal silicification alteration. Quartz vein with 35% QBS with weak silica alteration and weak chlorite alteration. Py occurs in vein material and as sub-euhedral crystals in the schist wall rock. Some qtz grains have fine veinlets of calcite. Biotite minor composition of host rock. Note: water becoming more of an issue--slowing progress. 2% chlorite, 2% silicification alteration. 5% py mineralization.	E862218	264	1370
BV15-30	48.77	50.29	Grey	QBS	DG	0	10	Silicified schist with 10% qtz veins. Py occurs in the schist and vein material. Biotite ~10%. Veins dramatically dropped off from previous interval but rock has a bleached silicified appearance--could have a weak calcite +/- dolomite alteration as well in the schist. 80% quartz, moderate silicification alteration, 8% py mineralization.	E862219	145	514
BV15-30	50.29	51.82	Grey	QBS	DG	0	20	Silicified schist. Same host rock but very bleached. Pyrite occurs in the altered host rock. Qtz veins comprise 20% of rock. Qtz also exhibits a pinkish brown (host rock qtz layers) reflecting moderate to locally intense potassic alteration. 70% quartz, moderate silicification alteration, moderate to intense potassic alteration. 4% py mineralization.	E862221	17	26
BV15-30	51.82	53.34	Grey	QBS	DG	0	15	Same schistose rock as above and has 15% qtz veins throughout sample. Schist is weak, to moderately potassically altered. Py occurs as anhedral-subhedral crystals. Qtz veins are observed parallel to schistosity. Moderate potassic alteration. 5% py mineralization.	E862222	9	21

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-30	53.34	54.86	Grey	QBS	DG	0	0	Biotite ~10%. Qtz bands are showing a weakly developing pinkish brown colour - potassic alteration. Schist hosts disseminated py. 85% quartz. Minimal potassic alteration. 3% pyrite mineralization. Note: Still dealing with water issues in the hole.	E862223	7	6
BV15-30	54.86	56.39	Grey	QBS	DG	0	12	Biotite ~10%. Qtz bands exhibit brownish pink colour: potassic alteration (moderate intensity). Pervasive in schist. Qtz veins ~12% of rock. No oxidation at all. 78% quartz. 3% py mineralization.	E862224	8	11
BV15-30	56.39	57.91	Grey	QBS	DG	0	12	Biotite ~10%. Qtz bands in schist exhibit moderate to strong potassic alteration. Py occurs as sub-euhedral 0.5-1mm crystals in schist. Qtz vein comprises 12% of rock.	E862225	-1	10
BV15-30	57.91	59.44	Grey	QBS	DG	0	12	Biotite 10%, Qtz vein 12%. Moderate potassic alteration. 5% pyrit mineralization.	E862226	8	14
BV15-30	59.44	60.96	Grey	QBS	DG	0	2	Biotite ~10%, qtz veins decrease to 2%. 88% quartz. Potassic alteration decreased to minimal. Subtle brownish colouration to rock. Pyrite occurs in schist.	E862227	6	8
BV15-31	0	1.52		CAS					No Sample		
BV15-31	1.52	3.05		CS	DG	4	0	Casing inserted to 10 and continuing to 15 feet. Chlorite rich schist. Strongly foliated. Rare feldspar rich layers banded with the chlorite. Rock is pervasively oxidized. Trace py (0.2%) mineralization. Moderately to intensely oxidized.	E862229	-1	55
BV15-31	3.05	4.57		CS	DG	4	0	Difficult to recognize protolith due to level of oxidation. Comprised of mainly chlorite defining schistosity layered with feldspar. Moderate to intense oxidation is pervasive. Casing continues to 15 feet and they are putting in another 5 feet.	E862231	-1	31
BV15-31	4.57	6.1		CS	DG	4	0	Oxidation is still moderate to strong but it is decreasing. Rare quartz vein fragments.	E862232	-1	17
BV15-31	6.1	7.62		CS	DG	4	0	Possible silicification. Rock is mainly chlorite and feldspar but it is harder than normal so possible silicification. Oxidation is moderate to intense, and pervasive; silicification is moderate.	E862233	-1	16
BV15-31	7.62	9.14	Green	CS	DG	4	0	Moderate to strong oxidation (pervasive). Rock is mainly green with white bands of feldspar. Rock seems very hard for chlorite schist so inferring moderate silicification. Casing extended down to 30 feet.	E862234	-1	13
BV15-31	9.14	10.67	Brown	CS	DG	4	2.5	Schistose fabric. 2-3% quartz veins. Moderate to intense oxidation.	E862235	-1	10
BV15-31	10.67	12.19		CS	DG	4	1	1% quartz veins. **Hole started to caving in as they started to drill 40-45. Decided to trip out and put more casing in hole. Moderate to intense oxidation.	E862236	-1	20
BV15-31	12.19	13.72	Brown	CS	DG	4	7	Schistose fabric through rock. Some veins (~7%). Oxidation is moderate to intense and pervasive.	E862237	-1	11
BV15-31	13.72	15.24	Grey	CS	DG	0	5	Schist. Quartz veins occur parallel to the fabric and comprise ~5% of the rock. Minimal silicification.	E862238	5	28
BV15-31	15.24	16.76	Grey	QBS	DG	3	3	Lost the chlorite and gained biotite. Biotite ~15%. Quartz veining 3%. Moderate oxidation.	E862239	-1	24
BV15-31	16.76	18.29	Grey	QBS	DG	3	7	Biotite schist with quartz. Biotite ~60%, quartz veining ~7%. Quartz veins are all moderately oxidized. **Casing was extended to 60 feet due to squeezing of the hole while drilling ~35-40 feet. Extreme oxidation possibly related to fracturing and broken ground? Moderate Oxidation.	E862240	6	58
BV15-31	18.29	19.81	Brown	QBS	DG	4	20	Qtz vein ~20%. Moderate to intense oxidation.	E862241	8800	490
BV15-31	19.81	21.34	Grey	QBS	DG	3	3	Qtz vein ~3%. Moderate oxidation.	E862251	149	140
BV15-31	21.34	22.86	Grey	QBS	DG	2	1	Oxidation is decreasing. Qtz veins decreasing to 1%. Weak chlorite alteration because of subtle green colour. Minimal oxidation, minimal chlorite alteration.	E862252	-1	59
BV15-31	22.86	24.38		CS	DG	2	0	Dominated by chlorite again with feldspar veining trace and oxidation is decreasing by minor degree. Minimal oxidation.	E862253	5	85
BV15-31	24.38	25.91	Green	CS	DG	2	1	1% qtz veins and they are oxidized (minimally). Well developed foliation.	E862254	-1	93
BV15-31	25.91	27.43	Green	CS	DG	2	0	No veining. Oxidation seems to be on fine surfaces (minimal).	E862255	6	106
BV15-31	27.43	28.96	Green	CS	DG	1	0	Oxidation is rare.	E862256	-1	43
BV15-31	28.96	30.48	Green	CS	DG	0	0	Very minimal oxidation.	E862257	11	107
BV15-31	30.48	32	Grey	QBS	DG	0	0	Colour changed to bluish grey and biotite with quartz is visible. No oxidation. No sx.	E862258	-1	35
BV15-31	32	33.53	Grey	QBS	DG	3	15	15% of qtz veins. Pyrite is disseminated and subhedral crystals 0.5-1mm in size within the qtz veins. Moderate oxidation. 2% pyrite mineralization.	E862259	-1	16
BV15-31	33.53	35.05	Grey	QBS	DG	2	15	15% of interval is qtz vein. Qtz vein material commonly oxidized. Some chlorite almost appear to be quartzite with no biotite visible. Qtz almost has a sugary texture in some pieces. Minimal oxidation, Minimal chlorite alteration, 3% pyrite mineralization.	E862261	-1	27
BV15-31	35.05	36.58	Grey	QBS	DG	0	0	Py occurs as sub-euhedral sub-mm crystals in host rock. Steel grey schist locally exhibits a greenish hue, possibly reflecting chlorite alteration. Minimal chlorite alteration, 2% pyrite mineralization.	E862262	-1	9
BV15-31	36.58	38.1		QBS	DG	0	25	Qtz vein: 25% of interval. Flecks of chlorite within the qtz vein. Pyrite within veins and schist. Minimal chlorite alteration, 4% pyrite mineralization.	E862263	-1	7
BV15-31	38.1	39.62	Green	QBS	DG	0	2	Minor biotite, essentially a qtz rich rock. Qtz vein: 2%. Patchy green colour that irregularly developed through the rock. Minimal chlorite alteration, 1% pyrite mineralization.	E862264	-1	21
BV15-31	39.62	41.15	Grey	QBS	DG	0	5	5% quartz vein. 2% pyrite mineralization.	E862265	-1	27
BV15-31	41.15	42.67	Grey	QZT	DG	0	7	7% qtz - light and dark grains - all quartz. Minimal chlorite alteration.	E862266	4700	6190
BV15-31	42.67	44.2	Grey	QZT	DG	0	30	30% quartz vein mixed with the very bleached and light grey quartzite. Unit was a dark blue grey and has become a very light grey. No pyrite in the veins. Moderate silicification. 5% pyrite mineralization.	E862267	47700	33400
BV15-31	44.2	45.72	Grey	QZT	DG	2	7	Mixture of dark grey quartzite and light grey bleached quartzite. Pyrite is in vein material and host rock. Qtz vein =7%. Oxidation is minimal and of select vein grains. 3% pyrite mineralization. Moderate silicification.	E862268	304	2590
BV15-31	45.72	47.24	Grey	QBS	DG	0	0	Very minor biotite but enough present to create a platy cleavage. Rock is very siliceous and is dominated by quartz. Rock shows a subtle green colour - possibly reflecting chlorite. Minimal chlorite alteration.	E862269	25	99

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-31	47.24	48.77	Grey	QZT	DG	0	0	Light to dark grey poorly maintained rock. Very poor schistose fabric developed. Rock generally looks siliceous. Light grey colour. Very localized oxidation. Minimal silicification and chlorite alteration. 3% pyrite mineralization.	E862270	495	4660
BV15-31	48.77	50.29	Grey	QZT	DG	0	0	Light grey. Disseminated pyrite (5%). Also see single fleck of silver coloured sx (aspy?). 0.2% aspy mineralization. Moderate silicification. **In run below water was intersected. Increasing in flow as they progress.	E862271	2200	8660
BV15-31	50.29	51.82	Grey	QZT	DG	0	0	Massive siliceous rock that is merately silicified. Silicification causes light grey colour: bleaching. 5% quartz veining with euhedral crystals - sub mm in size. Wall rock also contains finely disseminated pyrite. Very weak oxidation of select grains. Pervasive silicification. Moderate silicification, 7% pyrite mineralization. **Contending with a lot of water!!	E862272	1200	6440
BV15-31	51.82	53.34	Grey	QZT	DG	2	15	0.5-1mm euhedral pyrite disseminated throughout the quartzite. ~15% of interval comprised of qtz vein with no pyrite in vein material. Weak oxidation of select grains. Moderate pervasive silicification. Minimal oxidation; 12% pyrite mineralization.	E862273	1000	2320
BV15-31	53.34	54.86	Grey	QZT	DG	0	0	Rock is not quite as bleached as interval above. Grains show both dark and light grey clasts. The light grey clasts typically have disseminated sulphides. Also see 0.5-1mm thick pyrite veins. Pyrite is typically euhedral and 0.5-1mm crystals. Minimal to moderate silicification, 10% pyrite mineralization.	E862274	301	1420
BV15-31	54.86	56.39	Grey	QZT	DG	0	15	Vairably silicified; some dark grey, some light grey but all looks to be the same siliceous rock. Pyrite (12%) is disseminated and occurs as 0.5-1mm euhedral crystals within the host rock. Qtz veins comprise 15% of interval and rarely contain sx. No oxidation.	E862275	312	479
BV15-31	56.39	57.91	Grey	QZT	DP	1	7	As above, 40% dark grey. Pyrite cubes and along seams parallel to foliation. 7% quartz vein. Local lmnt. No calcite. Minimal oxidation, 7% quartz, minimal silicification. 4% pyrite mineralization.	E862276	76	142
BV15-31	57.91	59.44	Grey	QZT	DP	1	5	Variably silicified. 20-30% QBS; pyrite cubes are ~1mm. Wet sample. Patchy, bright green chlorite along qtz vein margins. Minimal oxidation and chlorite alteration. 5% quartz. Moderately silicified. 2% pyrite mineralization.	E862277	418	746
BV15-31	59.44	60.96	Grey	QZT	DP	1	4	15% QBS; very wet sample. 4% quartz. Minimal silicification. 2% pyrite mineralization.	E862278	283	585
BV15-31	60.96	62.48	Grey	QZT	DP	1	3	35% QBS; wet sample. 3% quartz. 2% pyrite mineralization.	E862279	201	584
BV15-31	62.48	64.01	Grey	QZT	DP	0	3	30% QBS; wet. Very fine disseminated specks po. 3% quartz. 2% pyrite mineralization. Trace po.	E862281	637	500
BV15-31	64.01	65.53	Grey	QBS	DP	1	2	Medium grey; 30% qtzt. Wet sample. 2% quartz. 0.5% po, 1.5% pyrite mineralization.	E862282	240	765
BV15-31	65.53	67.06	Grey	QBS	DP	2	3	Medium grey; 30% qtzt. Wet sample. % quartz. trace po, 2% pyrite mineralization.	E862283	165	546
BV15-31	67.06	68.58	Grey	QBS	DP	1	3	Medium grey; 15% qtzt. Wet sample. 3% quartz. 1% pyrite mineralization.	E862284	378	1210
BV15-31	68.58	70.1	Grey	QBS	DP	0	2	Medium grey; 10% qtzt. Wet sample. 2% quartz. 1% pyrite mineralization.	E862285	223	733
BV15-32	0	1.52		CAS				No Sample			
BV15-32	1.52	3.05	Brown	QBS	DG	4		10% biotite	E862354	17	84
BV15-32	3.05	4.57		QBS	DG	4	3	Oxidation is essentially pervasive. ~3% of interval comprised of qtz vein. No sx.	E862355	-1	9
BV15-32	4.57	6.1		QBS	DG	3		Qtz rich bands with minor biotite forming thin septa. 5-10% of grains are QMS. Oxidation is dropping off. No sx. 25' of casing set in hole.	E862356	-1	8
BV15-32	6.1	7.62		QZT	DP	2		60% QTZT, 40% QBS. No sulphides seen.	E862357	6	11
BV15-32	7.62	9.14		QZT	DP	3		As above; more oxidized; faint biotite alteration brownish cast.	E862358	7	21
BV15-32	9.14	10.67		QZT	DP	3		10% QBS bands to 3mm wide py as irregular blebs to 0.75mm.	E862359	243	1680
BV15-32	10.67	12.19		QZT	DP	4		5% QBS; more oxidized; no sulphides seen.	E862361	509	1630
BV15-32	12.19	13.72	Grey	QZT	DP	2		20% QBS bands; medium to light grey finely foliated rock	E862362	10	43
BV15-32	13.72	15.24	Grey	QZT	DP	2		Light Grey; rare very fine speck py	E862363	-1	16
BV15-32	15.24	16.76	Grey	QZT	DP	2		Light Grey; rare specks py	E862364	7	20
BV15-32	16.76	18.29	Grey	QZT	DP	1		Light grey; very rare py	E862365	-1	6
BV15-32	18.29	19.81	Grey	QZT	DP	2		Light grey; no sulphides seen	E862366	8	17
BV15-32	19.81	21.34	Grey	QBS	DP	1		30% QTZT; medium grey; no sulphides seen; qtzt bands to 4mm	E862367	-1	8
BV15-32	21.34	22.86	Grey	QZT	DP	1		20% QBS; light grey; no sulphides seen	E862368	-1	9
BV15-32	22.86	24.38	Grey	QZT	DP	1		30% QBS; light grey; very rare fine speck sulphide	E862369	-1	4
BV15-32	24.38	25.91	Grey	QBS	DP	0	0.5	Medium grey; no sulphides; calcite veinlets to 0.4mm wide	E862370	7	6
BV15-32	25.91	27.43	Grey	QZT	DP	1		Light grey; py veinlet 0.3mm wide along fault surface	E862371	6	31
BV15-32	27.43	28.96	Grey	QZT	DP	0		Light grey; no sulphides seen	E862372	7	5
BV15-32	28.96	30.48	Grey	QBS	DP	0		30% QTZT; medium grey; py mass 2x5mm rimmed on one side by cpy.	E862373	-1	-1
BV15-32	30.48	32	Grey	QBS	DP	1	1	Medium brownish grey; rare speck sulphide	E862374	-1	8
BV15-32	32	33.53	Grey	QBS	DP	1		Light grey; no sulphides seen granular, quartz-rich so ~ transitional to qtzt.	E862375	-1	-1
BV15-32	33.53	35.05	Grey	QBS	DP	0	0.5	Light grey --> transitional to qtzt; sulphides as small blebs.	E862376	-1	4
BV15-32	35.05	36.58	Grey	QBS	DP	1	0.5	Light grey; blebby sulphides	E862377	5	12
BV15-32	36.58	38.1	Grey	QBS	DP	0	0.2	Light brownish grey	E862378	-1	-1
BV15-32	38.1	39.62	Grey	QBS	DP	1		Light brownish grey; irregular blebby po masses	E862379	-1	5
BV15-32	39.62	41.15	Grey	QBS	DP	0	0.2	as above; one speck cpy seen beside larger po bleb.	E862381	-1	7
BV15-32	41.15	42.67	Grey	QBS	DP	0		Light brownish grey; qtzt bands to 2mm wide. Rare po specks.	E862382	9	5
BV15-32	42.67	44.2	Grey	QBS	DP	2	40	60% QBS, 40% qtz vein; traces po in QBS; pervasive yellow-orange faint lmnt stair on vein fragments; no sulphides seen within vein qtz.	E862383	-1	4

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-32	44.2	45.72	Grey	QBS	DP	1	7	Light grey; vein frags as for last interval - pervasive limonite throughout = no sulphides seen in vein quartz	E862384	-1	4
BV15-32	45.72	47.24	Grey	QBS	DP	1	4	Light grey' orange lmnt on some fracture sites within schist	E862385	5	4
BV15-32	47.24	48.77	Grey	QBS	DP	0	0	Light grey; much less biotite than seen higher in hole. ~ zero sulphides.	E862386	15	4
BV15-32	48.77	50.29	Grey	QBS	DP	0	2	Light grey; as above, except here vein qtz frags contain very fine pyrite	E862387	6	5
BV15-32	50.29	51.82	Grey	QBS	DP	1	0	Light brownish grey; lmnt but no sulphides seen.	E862388	5	6
BV15-32	51.82	53.34	Grey	QBS	DP	0		Light brownish grey; no sulphides seen	E862389	-1	5
BV15-32	53.34	54.86	Grey	QBS	DP	0		Light brownish grey; no sulphides seen	E862391	8	3
BV15-32	54.86	56.39	Grey	QBS	DP	1		as above but silt size material in sample light yellow-brown, limonite	E862392	-1	18
BV15-32	56.39	57.91	Grey	QBS	DP	1		as above; ~ less limonite fine in sample	E862393	6	27
BV15-32	57.91	59.44	Grey	QBS	DP	0		as above; limonite no more within fine size fraction	E862394	-1	4
BV15-32	59.44	60.96	Grey	QBS	DP	0	0.5	as above	E862395	-1	-1
BV15-32	60.96	62.48	Grey	QBS	DP	0		Light brownish grey; no sulphides seen	E862396	-1	-1
BV15-32	62.48	64.01	Grey	QBS	DP	0		Light brownish grey; granular qtz bands to 4mm; no sulphides seen	E862397	-1	-1
BV15-32	64.01	65.53	Grey	QBS	DP	0		As above	E862398	-1	6
BV15-32	65.53	67.06	Grey	QBS	DP	0		As above	E862399	6	11
BV15-32	67.06	68.58	Grey	QBS	DP	0	0.5	As above except for fine off-white quartz veinlet	E862400	12	54
BV15-32	68.58	70.1	Grey	QZT	DP	0		10% QBS as above. Qtzt pale grey, finely granular with lensoid sulphide mass parallel to foliation, 1mm wide by 8mm long	E862401	122	351
BV15-32	70.1	71.63	Grey	QZT	DP	0		30% QBS as above; sulphides seen but light poor	E862402	-1	8
BV15-32	71.63	73.15	Grey	QBS	DP	0	0.5	Medium brownish grey; interval 5% qtzt	E862403	-1	6
BV15-32	73.15	74.68	Grey	QBS	DP	0		As above; no sulphides seen	E862404	8	15
BV15-32	74.68	76.2	Grey	QBS	DP	0	0.5	10% qtzt; few small, irregular po blebs.	E862405	-1	4
BV15-32	76.2	77.72	Grey	QBS	DP	0		medium brownish grey; moderately well developed foliation throughout	E862406	-1	4
BV15-32	77.72	79.25	Grey	QBS	DP	1	0.5	35% qtzt; light brownish grey; brownish yellow lmnt on fracture surfaces; no sulphides seen	E862407	26	31
BV15-32	79.25	80.77	Grey	QZT	DP	0	5	10% QBS; well mineralized qtzt with say 5% pale grey vein qtz. Occasional subhedral py grains but mostly blebby sulphides	E862408	175	160
BV15-32	80.77	82.3	Grey	QZT	DP	0		20% QBS. Local 3% sulphides as blebs to 1.5x3mm but average ~0.4% sulphides combined	E862409	23	43
BV15-32	82.3	83.82	Grey	QBS	DP	0		Light brownish grey; qtzt bands to 2.5mm wide blebby po, lesser py.	E862410	-1	7
BV15-32	83.82	85.34	Grey	QBS	DP	0	2	20% Qtzt; py bands to 4mm wide	E862411	-1	5
BV15-32	85.34	86.87	Grey	QBS	DP	0	0.5	15% qtzt; irregular blebs, po most abundant sulphide	E862412	-1	5
BV15-32	86.87	88.39	Grey	QBS	DP	0	1	Medium brownish grey	E862413	-1	6
BV15-32	88.39	89.92	Grey	QBS	DP	0		Medium brownish grey; 10% Qtzt.	E862414	-1	5
BV15-32	89.92	91.44	Grey	QBS	DG	0		3% qtz veins. Single chip displayed qtz vein within schist parallell to the schistosity. Minor fine grained py within the qtz veins and the schist	E862415	-1	3
BV15-32	91.44	92.96	Grey	QBS	DG	0	2	Qtz veins ~2%. Biotite comprises 10-15% of interval. Qtz vein in a single chip is seen cross cutting the schistosity.	E862416	-1	4
BV15-32	92.96	94.49	Grey	QBS	DG	0	0.1	Trace qtz veins.	E862417	-1	5
BV15-32	94.49	96.01	Grey	QBS	DG	0		No qtz veins. Fine grained sx through schist. Qtz forms 1-3mm bands separated by fine septa of biotite	E862418	-1	5
BV15-32	96.01	97.54	Grey	QBS	DG	0	0.1	Trace qtz veins. Fine grained disseminated py.	E862419	10	5
BV15-32	97.54	99.06	Grey	QBS	DG	0	3	3-4% qtz veins. Qtz vein margin is parallel to the schistosity. Also see py, fine grained and disseminated, within schist and coarser, euhedral py along vein margins.	E862421	-1	9
BV15-32	99.06	100.58	Grey	QBS	DG	0		~30% of the interval exhibits an orange brown colour reflecting an increased level of oxidation. No qtz veining in this run. Py is disseminated and wispy within schist.	E862422	-1	9
BV15-32	100.58	102.11	Grey	QBS	DG	0		Oxidation is gone	E862423	-1	18
BV15-32	102.11	103.63	Grey	QBS	DG	0	3	Qtz veining comprises ~3% of grains. ***Changed to the denser bit***	E862424	-1	9
BV15-32	103.63	105.16	Grey	QBS	DG	0		No veining. No sulphides	E862425	-1	-1
BV15-32	105.16	106.68	Grey	QBS	DG	2	5	Qtz veining at 5%. Oxidation of both vein and schist increase in this interval. Py is fine grained and disseminated.	E862426	-1	13
BV15-32	106.68	108.2	Grey	QBS	DG	0	4	Qtz veining at ~4%. Trace oxidation. Pyrite occurs with schist - fine grained and dissminated. Also see rare euhedral cubes along vein margins in vein and schist.	E862427	-1	17
BV15-32	108.2	109.73	Grey	QBS	DG	0	4	Qtz veining at 4%.	E862428	-1	5
BV15-32	109.73	111.25	Grey	QBS	DG	0	3	Qtz veins: 3%. ~5% biotite. Py is fine grained and occurs in the schist.	E862429	-1	6
BV15-32	111.25	112.78	Grey	QBS	DG	0	3	Qtz veins 3%	E862431	-1	5
BV15-32	112.78	114.3	Grey	QBS	DG	0	3	Qtz veins @ 3%	E862432	-1	8
BV15-32	114.3	115.82	Grey	QBS	DG	0	2	Qtz veins ~2%	E862433	-1	4
BV15-32	115.82	117.35	Grey	QBS	DG	0	1	Qtz veins ~1%. Schist comprised of 1-5mm thick qtz rich bands separated by very fine septa of biotite. Biotite comprises 7% of rock. Py is fine grained and occurs within the schist and the vein.	E862434	-1	4
BV15-32	117.35	118.87	Grey	QBS	DG	0	1	Qtz veins ~1%. Sx decreasing. Fine grained and dissminated within schist.	E862435	-1	5
BV15-32	118.87	120.4	Grey	QBS	DG	0	0.1	Qtz veining absent. No sx.	E862436	-1	-1
BV15-32	120.4	121.92	Grey	QBS	DG	0		Trace qtz veins	E862437	-1	7
BV15-32	121.92	123.44	Grey	QBS	DG	0	0.1	No qtz veins. Sx increases - disseminated through the schist.	E862438	-1	-1

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-32	123.44	124.97	Grey	QBS	DG	0		Trace qtz veins. Fine grained disseminated sx through schist. ***Rig shut down 45min due to lightning in the area***	E862439	6	-1
BV15-32	124.97	126.49	Grey	QBS	DG	0		1% qtz veins - single 0.4cm vein observed in a grain	E862440	5	7
BV15-32	126.49	128.02	Grey	QBS	DG	0	3	Qtz veins comprise 3%	E862441	-1	4
BV15-32	128.02	129.54	Grey	QBS	DG	0	1	Qtz veins ~1%. Minor py, fine grained and disseminated within the schist	E862442	-1	8
BV15-32	129.54	131.06	Grey	QBS	DG	0	0.5	Fine grained, faintly foliated.	E862443	-1	8
BV15-32	131.06	132.59	Grey	QBS	DG	0	0.3	Qtzt band 3mm wide contains blebs po.	E862444	-1	4
BV15-32	132.59	134.11	Grey	QBS	DG	1		As above except limonite oxides on some fracture surfaces.	E862445	-1	8
BV15-32	134.11	135.64	Grey	QBS	DG	0		Medium brownish grey; more quartzite than above intervals. Also more biotite-altered. **20:35 bit done at 444.5' so pulling rods***	E862446	7	10
BV15-33	0	1.52		CAS					No Sample		
BV15-33	1.52	3.05	Grey	QBS	DG	3		Generally all one rock type but also see QMS. Interval most likely bedrock and overburden. Moderate oxidation. 1% pyrite mineralization.	E862447	9	36
BV15-33	3.05	4.57	Grey	QBS	DG			Oxidation drops off dramatically but still weak. Sx are disseminated within schist; are fine grained. A moderately developed fabric is present, possibly only moderately developed because of lack of mica. 95% quartz. Minimal oxidation, 0.3% pyrite mineralization.	E862448	-1	10
BV15-33	4.57	6.1	Grey	QMS	DG			Light greenish grey QMS with minor QBS grains. Subtle green colour could indicate a weak chlorite overprint. Minor pyrite disseminated irregularly through the schist. Minimal chlorite alteration. 0.3% pyrite mineralization.	E862449	-1	10
BV15-33	6.1	7.62	Green	QZT	DG		20	Grey green quartz. Qtz vein ~30% of rock. Pyrite is fine grained and in vein and quartzite. 0.5% pyrite mineralization.	E862451	-1	5
BV15-33	7.62	9.14	Grey	QBS	DG			~5-10% of interval consists of greenish-grey quartzite. Also no vein material in this run. Biotite is variable ~ 5-10%. 90% quartz, 0.3% pyrite mineralization.	E862452	-1	7
BV15-33	9.14	10.67	Grey	QBS	DG			~5-10% biotite in sub mm layers between 1-5mm quartz bands. No veining and no oxidation. Pyrite (0.5%) is disseminated and fine grained.	E862453	17	9
BV15-33	10.67	12.19	Grey	QBS	DG			10-15% biotite. No oxidation or veining. 85% quartz. 0.5% pyrite mineralization.	E862454	-1	10
BV15-33	12.19	13.72	Grey	QBS	DG	2		10% biotite. Oxidation appeared to start over lowermost 1-2 feet. Minimal oxidation. 90% quartz. 0.3% pyrite mineralization.	E862455	-1	15
BV15-33	13.72	15.24	Grey	QBS	DG	2	1	Qtz vein: 1%. Pyrite is fine grained and disseminated and aligned in foliation. 1.5% pyrite mineralization.	E862456	19	14
BV15-33	15.24	16.76	Grey	QBS	DG	2		No veining. Minimal oxidation. 90% quartz. 0.3% pyrite mineralization.	E862457	17	73
BV15-33	16.76	18.29	Grey	QBS	DG	0	3	3% quartz veins. Minimal oxidation. 87% quartz. 0.3% pyrite mineralization.	E862458	12	19
BV15-33	18.29	19.81	Grey	QBS	DG	0		~10-15% quartzite - greenish grey. 90% quartz. 0.2% pyrite mineralization.	E862459	13	11
BV15-33	19.81	21.34	Grey	QBS	DG	0		5-10% biotite. No quartz veining. Trace oxidation. Pyrite is fine grained and forms sub mm masses aligned within foliation plane. 0.5 pyrite mineralization.	E862461	-1	17
BV15-33	21.34	22.86	Grey	QBS	DG	0		5-10% biotite. No qtz veining. No oxidation. 90% quartz. 0.3% pyrite mineralization.	E862462	5	21
BV15-33	22.86	24.38	Grey	QBS	DG	0		5-8% biotite. 5% calcite veining. 88% quartz. 0.5 pyrite mineralization.	E862463	7	13
BV15-33	24.38	25.91	Grey	QBS	DG	2		~80% of chips are QBS; 5% calcite vein and 15% QMS. A light grey rock that iso contains minor biotite. Minimal oxidation. 85% quartz. 0.3% pyrite mineralization.	E862464	631	4200
BV15-33	25.91	27.43	Grey	QZT	DG		1	Mixture of greenish grey quartzite, calcite veins, quartz veins and QBS. Quartzite dominates at ~60%. Single chip 1x0.5cm is a quartz calcite vein fragment with two 1-2cm wide bands of sx - one band is euhedral pyrite and the other is fine grained silver sulphide - possible arsenopyrite and very fine grained but forming a massive veinlet parallel to the pyrite vein. Minimal chlorite alteration. 0.2po, 0.2% aspy mineralization.	E862465	31	513
BV15-33	27.43	28.96	Grey	QZT	DG		20	Greenish grey quartz within ~45-50% quartz calcite veining. Rare siliceous blue grey bands with possible very fine grained sx. Minimal chlorite alteration. 0.5% pyrite mineralization.	E862466	13	48
BV15-33	28.96	30.48	Green	QBS	DG		1.5	Qtz calcite veining ~ 3%. 7% of chips are greyish green quartzite. Minor pyrite disseminated within schist ~10% biotite. Rare +1cm quartzite clasts exhibit 0.5cm thick smokey grey bands that possibly could have very fine grained disseminated sx. 0.5% pyrite mineralization.	E862467	13	21
BV15-33	30.48	32	Green	QBS	DG		0.1	25% of chips are the grey-green quartzite. Massive with very poor foliation development. Trace quartz veining. 1% pyrite mineralization.	E862468	8	10
BV15-33	32	33.53	Grey	QBS	DG		7	7% of chips are quartz veins. Sx occur in the viens and within the schist. 1% pyrite mineralization.	E862469	15	21
BV15-33	33.53	35.05	Grey	QBS	DG			No veining and only trace pyrite (0.1%) mineralization.	E862470	13	15
BV15-33	35.05	36.58	Green	QBS	DG			Biotite ~10% of rock. No quartz veining. 15% of the chips exhibit a greenish grey colour representing a sub population of quartz. 90% quartz. Minimal chlorite alteration. 0.2% pyrite alteration.	E862471	24	55
BV15-33	36.58	38.1	Grey	QBS	DG		0.1	Trace quartz veins. 90% quartz. 0.3% pyrite mineralization.	E862472	-1	37
BV15-33	38.1	39.62	Grey	QBS	DG		3	3% quartz veins. 90% quartz. 0.3% pyrite mineralization.	E862473	7	31
BV15-33	39.62	41.15	Grey	QBS	DG		1	15-20% greenish grey quartzite. More difficult to see the pyrite within the QBS but there generally appears to be slightly more sx in the quartzite compared with the schist. 1% quartz veining. 1% pyrite mineralization.	E862474	19	60
BV15-33	41.15	42.67	Grey	QBS	DG		2	2% quartz veins. 5-7% greenish grey quartzite. 90% quartz. 0.3% pyrite mineralization.	E862475	5	21
BV15-33	42.67	44.2	Grey	QBS	DG		4	4% quartz veins +/- calcite veinlets. 90% quartz. 0.5% pyrite mineralization.	E862476	20	45
BV15-33	44.2	45.72	Grey	QBS	DG		0.1	Trace quartz veins with fine to medium grained euhedral pyrite (0.5% pyrite mineralization). 90% quartz.	E862477	23	36
BV15-33	45.72	47.24	Grey	QBS	DG		0.1	Trace quartz veins. 90% quartz. 0.3% pyrite mineralization.	E862478	15	17
BV15-33	47.24	48.77	Grey	QBS	DG		0.1	Trace quartz veins.	E862479	13	20

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-33	48.77	50.29	Grey	QBS	DG		0.1	Trace quartz veins. 90% quartz. 0.3% pyrite mineralization.	E862481	24	56
BV15-33	50.29	51.82	Grey	QBS	DG			The green colouration mentioned above is silicification. Can see a chip with a hair line fracture and the green colour is zoned around the healed hair line fracture. Still not common enough through the chips to be representative alteration throughout. 90% quartz. 0.3% pyrite mineralization.	E862482	5	17
BV15-33	51.82	53.34	Grey	QBS	DG		2.5	Quartz-calcite veins ~5%. Weak silicification causing s bleaching. 0.3% pyrite mineralization.	E862483	6	14
BV15-33	53.34	54.86	Grey	QBS	DG		3	Quartz calcite vein ~6%. 90% quartz. Minimal silicification. 0.3% pyrite mineralization.	E862484	-1	9
BV15-33	54.86	56.39	Grey	QZT	DG	3		Light greenish grey. No sx. Oxidation increases to moderate.	E862485	5	9
BV15-33	56.39	57.91	Grey	QZT	DG			Light grey with green flecks throughout quartzite. Poorly banded/foliated. Very weak chlorite alteration. 0.4% pyrite mineralization.	E862486	-1	16
BV15-33	57.91	59.44	Grey	QZT	DG			Irregularly developed pink staining in select chips. Weak chlorite and minimal potassic alteration. 0.3% pyrite mineralization. 95% quartz.	E862487	6	8
BV15-33	59.44	60.96	Grey	QBS	DG		2	~2% quartz vein. Biotite ~8%. 90% quartz. 0.3% pyrite mineralization.	E862488	-1	8
BV15-33	60.96	62.48	Grey	QBS	DG		1	Mixture of QBS - 60%, and light greenish grey quartzite. The schist exhibits a pinkish brown colour reflecting a weak potassic overprint. 2% quartz calcite veining. 0.3% pyrite mineralization.	E862489	-1	6
BV15-33	62.48	64.01	Grey	QZT	DG		2.5	Mixture of quartzite (80%Q, QBS (15%), Quartz calcite veining (5%). 0.3% pyrite mineralization.	E862491	-1	8
BV15-33	64.01	65.53	Grey	QZT	DG		5	Quartz calcite veining ~10%. 4% QBS. Pyrite observed coating grains discontinuously as well as random clots occurring within the quartzite. 0.5% pyrite mineralization.	E862492	-1	16
BV15-33	65.53	67.06	Grey	QBS	DG			No sx, no veining. 95% quartz.	E862493	-1	10
BV15-33	67.06	68.58	Grey	QBS	DG			No sx, no veining. 95% quartz.	E862494	-1	9
BV15-33	68.58	70.1	Grey	QBS	DG		1	No sx, 1% quartz veining. 95% quartz.	E862495	-1	8
BV15-33	70.1	71.63	Grey	QBS	DP		0.5	0.5% quartz veinlets; rare speck po. 90% quartz. Trace po mineralization.	E862496	-1	8
BV15-33	71.63	73.15	Grey	QBS	DP		0.5	0.5% quartz veinlets. 90% quartz. Trace biotite alteration. 0.1% py/po mineralization.	E862497	-1	7
BV15-33	73.15	74.68	Grey	QBS	DP		0.5	0.5% quartz veinlets; py/po specks; green chlorite (?? Epidote) appears. 90% quartz. Trace chlorite and biotite alteration. 0.1% po/py mineralization.	E862498	-1	6
BV15-33	74.68	76.2	Grey	QBS	DP			QBS to 249'. 249' into white coarse grained quartzite to traces green chlorite. No sulphides seen in quartzite. 90% quartz. 0.1% cpy, 0.3% po mineralization.	E862499	-1	10
BV15-33	76.2	77.72	White	QZT	DP			Medium grained, greenish white. 98% quartz. Weak chlorite/sericite/potassic alteration. 0.1% pyrite mineralization.	E862500	-1	6
BV15-33	77.72	79.25	White	QZT	DP			Medium grained, greenish white. 98% quartz. Weak chlorite and sericite. 0.1% pyrite mineralization.	E863001	-1	8
BV15-33	79.25	80.77	Grey	QZT	DP			30% QBS. 98% quartz. Weak chlorite alteration. 0.1% py/po mineralization.	E863002	-1	11
BV15-33	80.77	82.3	Grey	QBS	DP			30% quartzite. 97% quartz. Weak chlorite alteration. 0.1% pyrite mineralization.	E863003	-1	8
BV15-33	82.3	83.82	Grey	QBS	DP			20% quartzite. 95% quartz. Weak chlorite and biotite alteration. 0.5% py and 0.2% po mineralization.	E863004	6	9
BV15-33	83.82	85.34	Grey	QBS	DP			30% quartzite. 97% quartz. Weak chlorite alteration. 0.1% py/po mineralization.	E863005	-1	7
BV15-33	85.34	86.87	Grey	QBS	DP			No quartzite. 95% quartz. Trace biotite alteration. 0.1% py/py mineralization.	E863006	-1	11
BV15-33	86.87	88.39	Grey	QBS	DP				E863007	-1	12
BV15-33	88.39	89.92	Grey	QBS	DP			15% quartzite.	E863008	-1	9
BV15-33	89.92	91.44	Grey	QBS	DP			<not enough light to describe sample>	E863009	-1	13
BV15-33	91.44	92.96	Grey	QZT	DP			30% QBS.	E863010	-1	7
BV15-33	92.96	94.49	Grey	QZT	DP			5% QBS. 0.1% pyrite mineralization.	E863011	12	9
BV15-33	94.49	96.01	White	QZT	DP			Quartzite finely banded.	E863012	7	9
BV15-33	96.01	97.54	Grey	QZT	DP			Rare speck pyrite; quartzite medium to coarse grained. 100% quartz.	E863013	-1	10
BV15-33	97.54	99.06	Grey	QZT	DP			Medium grained pale grey. 100% quartz.	E863014	-1	8
BV15-33	99.06	100.58	Grey	QZT	DP			Pale grey, fine to medium grained. Trace potassic alteration. 0.1% pyrite mineralization.	E863015	-1	12
BV15-33	100.58	101.5	Grey	QBS	DP	1		15% quartzite. Very weak oxidation. 90% quartz. 0.1% pyrite mineralization.	E863016	-1	8
BV15-34	0	1.52		CAS					No Sample		
BV15-34	1.52	3.05	Orange	GR	DG	3		Medium to coarse grained granite. No foliation. Quartz feldspar and biotite are primary minerals. Minor component of schistose rocks so possibly could still be overburden. Moderate oxidation.	E863017	-1	8
BV15-34	3.05	4.57	Orange	GR	DG	3		Moderate oxidation. No ss.	E863018	-1	8
BV15-34	4.57	6.1	Grey	GR	DG	4		Moderate to strong oxidation. No sx. 10-20% schistose grains therefore could still be in overburden.	E863019	-1	6
BV15-34	6.1	7.62	Grey	QZT	DG	3		Rare chips of granite and mostly see quartzite with very minor layers with slight increase in biotite. Pyrite (0.3%) forms medium grained euhedral crystals.	E863021	-1	13
BV15-34	7.62	9.14	Grey	QZT	DG	4	0.1	Trace quartz veining. Moderate to intense oxidation. 0.1% pyrite mineralization.	E863022	-1	13
BV15-34	9.14	10.67	Grey	QBS	DG	1		15% biotite. No veining. Very weak oxidation. 85% quartz.	E863023/E863024	-1	5
BV15-34	10.67	12.19	Grey	QZT	DG	2		Light grey. No veining. Minimal oxidation. 0.1% py mineralization.			
BV15-34	12.19	13.72	Grey	QZT	DG	3		No veining. Moderate oxidation. 0.1% pyrite mineralization.	E863025	20	71
BV15-34	13.72	15.24	Grey	QZT	DG	2		No veining - ~25% of interval is QBS. Minimal oxidation. 99% quartz. 0.1% pyrite mineralization.	E863026	-1	10
BV15-34	15.24	16.76	Grey	QZT	DG			No veining. 40% QBS. Minimal oxidation. 95% quartz. 0.1% pyrite mineralization.	E863027	-1	8
BV15-34	16.76	18.29	Grey	QBS	DG			Mix of QBS (~75) and quartzite (25%). No veining.	E863028	-1	6
BV15-34	18.29	19.81	Grey	QBS	DG			25% quartzite. 95% quartz. 0.1% pyrite mineralization.	E863029	-1	8
BV15-34	19.81	21.34	Grey	QBS	DG	2		45% quartzite. Minimal oxidation. 95% quartz. 0.1% pyrite mineralization.	E863031	-1	4

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-34	21.34	22.86	Grey	QZT	DG			~45% QBS. Non magnetic. 98% quartz. 0.2% pyrite mineralization.	E863032	-1	8
BV15-34	22.86	24.38	Grey	QZT	DG			~30% QBS. Pyrite (0.3%) is fine grained and occurs irregularly. Trace oxidation. 98% quartz.	E863033	-1	5
BV15-34	24.38	25.91	Grey	QBS	DG			~20% quartzite. QBS consists of ~5% biotite. Biotite occurs more as irregular flecks throughout quartz instead of distinct continuous septa. Pyrite (0.3%) is pin head size and disseminated. 95% quartz.	E863034	-1	5
BV15-34	25.91	27.43	Grey	QZT	DG			~35% QBS. No veining. Quartzite typically has a greenish grey colour. The QBS is dark grey reflecting the biotite mixed with quartz. 98% quartz. 0.1% pyrite mineralization.	E863035	-1	5
BV15-34	27.43	28.96	Grey	QZT	DG			~35% QBS. 0.3% pyrite mineralization.	E863036	-1	6
BV15-34	28.96	30.48	Grey	QBS	DG			~40% quartzite. Non magnetic. 95% quartz.	E863037	6	9
BV15-34	30.48	32	Grey	QZT	DG			10% QBS. 98% quartz.	E863038	-1	11
BV15-34	32	33.53	Grey	QZT	DG			30% QBS. Non magnetic. 0.2% pyrite mineralization.	E863039	-1	6
BV15-34	33.53	35.05	Grey	QZT	DG	2		25% QBS. Non magnetic. Minimal oxidation. 98% quartz. 0.2% pyrite mineralization.	E863040	-1	6
BV15-34	35.05	36.58	Grey	QZT	DG	2		10% QBS. Minimal oxidation. 0.1% pyrite mineralization.	E863041	-1	8
BV15-34	36.58	38.1	Grey	QZT	DG	3		7% QBS. Moderate oxidation. 0.1% pyrite mineralization.	E863042	-1	7
BV15-34	38.1	39.62	Grey	QZT	DG	4		3% QBS. Pervasive oxidation in most chips. Sx have disappeared. Oxidation is intense.	E863043	-1	8
BV15-34	39.62	41.15	Grey	QZT	DG	3		10% QBS. Moderate oxidation.	E863044	-1	35
BV15-34	41.15	42.67	Grey	QZT	DG	3		~35% granite. Moderate oxidation.	E863045	9	26
BV15-34	42.67	44.2	Grey	QZT	DG	2	3	~3% quartz vein. Minimal oxidation. 98% quartz. 0.2% pyrite mineralization.	E863046	-1	34
BV15-34	44.2	45.72	White	QZT	DG		15	~15-20% quartz vein. Very difficult to see features because interval had a high clay content and stuck to the surface of the grains. Appeared to be a mixture of quartzite and quartz veins.	E863047	-1	92
BV15-34	45.72	47.24	White	QZT	DG		70	Sample estimated to have 70-80% quartz vein with remaining being light grey quartzite. Drilling through clay seam before/during intersecting vein. No sx. Just see clay and not specifically sericite. Clay looks white and chalky more like kaolinite. Intense Sericite alteration.	E863048	-1	28
BV15-34	47.24	48.77	Green	CS	DG		6	Chlorite schist. 12% quartz-calcite vein.	E863049	-1	10
BV15-34	48.77	50.29	Green	CS	DG		3.5	7% quartz calcite veining.	E863051	-1	8
BV15-34	50.29	51.82	Green	CS	DG		0.5	1% quartz calcite veining. Strong foliation in rock.	E863052	-1	7
BV15-34	51.82	53.34	Green	CS	DG		0.5		E863053	-1	3
BV15-34	53.34	54.86	Green	CS	DG		0.5		E863054	-1	10
BV15-34	54.86	56.39	Green	CS	DG		0.5		E863055	-1	8
BV15-34	56.39	57.91	Green	CS	DG		0.1	Trace quartz calcite veins.	E863056	-1	8
BV15-34	57.91	59.44	Green	CS	DG				E863057	-1	5
BV15-34	59.44	60.96	Green	CS	DG				E863058	-1	6
BV15-34	60.96	62.48	Green	CS	DG				E863059	-1	5
BV15-34	62.48	64.01	Green	CS	DG				E863061	-1	7
BV15-34	64.01	65.53	Green	CS	DG				E863062	7	6
BV15-34	65.53	67.06	Green	CS	DP				E863063	15	60
BV15-34	67.06	68.58	Green	CS	DP				E863064	-1	8
BV15-34	68.58	70.1	Green	CS	DP				E863065	-1	6
BV15-34	70.1	71.63	Green	CS	DP				E863066	-1	6
BV15-34	71.63	73.15	Green	CS	DP				E863067	-1	10
BV15-34	73.15	74.68	Green	CS	DP				E863068	-1	6
BV15-34	74.68	76.2	Green	CS	DP				E863069	-1	6
BV15-34	76.2	77.72	Green	CS	DP				E863070	-1	3
BV15-34	77.72	79.25	Green	CS	DP				E863071	-1	5
BV15-34	79.25	80.77	Green	CS	DP				E863072	8	41
BV15-34	80.77	82.3	Grey	QZT	DP			Light grey fine grained; weak biotite altered.	E863073	44	181
BV15-34	82.3	83.82	Grey	QZT	DP			As above but here only rare speck pyrite and less biotite than last interval.	E863074	-1	36
BV15-34	83.82	85.34		QZT	DP		40		E863075	-1	17
BV15-34	85.34	86.87	Grey	QZT	DP				E863076	-1	9
BV15-34	86.87	88.39		QZT	DP				E863077	7	16
BV15-34	88.39	89.92	Grey	QZT	DP			Pyrite very fine, sooty	E863078	54	217
BV15-34	89.92	91.44	Grey	QZT	DP			~5% qtz grains, medium grey, with pyrite as subhedral crystals to 1mm across plus dark grey sooty sulphides	E863079	79	240
BV15-34	91.44	92.96	Grey	QZT	DP			Pyrite evenly distributed	E863081	-1	11
BV15-34	92.96	94.49		QZT	DP			As above, weak sericite altered?	E863082	-1	5
BV15-34	94.49	96.01	White	QZT	DP			More pyrite than last interval	E863083	-1	7
BV15-34	96.01	97.54	White	QZT	DP			Traces of specks	E863084	5	6
BV15-34	97.54	99.06	White	QZT	DP				E863085	-1	5
BV15-34	99.06	100.58	White	QZT	DP			Traces of chlorite	E863086	-1	-1
BV15-34	100.58	102.11	White	QZT	DP				E863087	-1	-1
BV15-34	102.11	103.63	White	QZT	DP				E863088	6	33

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-34	103.63	105.16	White	QZT	DP				E863089	18	43
BV15-34	105.16	106.68		QZT	DP				E863091	56	170
BV15-34	106.68	108.2		QZT	DP				E863092	64	297
BV15-34	108.2	109.73	Grey	QZT	DP				E863093	13	61
BV15-34	109.73	111.25	Grey	QZT	DP				E863094	26	67
BV15-34	111.25	112.78	Grey	QZT	DP				E863095	-1	6
BV15-34	112.78	114.3	Grey	QZT	DP				E863096	6	15
BV15-34	114.3	115.82	Grey	QZT	DP				E863097	-1	12
BV15-34	115.82	117.35	Grey	QZT	DP				E863098	-1	5
BV15-34	117.35	118.87	Grey	QZT	DP				E863099	-1	6
BV15-34	118.87	120.4		QZT	DP				E863100	-1	7
BV15-34	120.4	121.92		QZT	DP				E863101	26	64
BV15-34	121.92	123.44	White	QZT	DP			Off-white to greenish. Wet sample	E863102	12	82
BV15-34	123.44	124.97	White	QZT	DP			Off-white to greenish. Wet sample	E863103	6	32
BV15-35	0	1.52		OB					No Sample		
BV15-35	1.52	3.05	Black	QBS	DP	1	5	QBS + Overburden	L399351	19	45
BV15-35	3.05	4.57	Grey	QZT	DP	1	0	Medium grey, 5% biotite	L399352	-1	18
BV15-35	4.57	6.1	Grey	QZT	DP	0	0	Medium grey, 7% biotite	L399353	20	13
BV15-35	6.1	7.62	Grey	QZT	DP	2	5	Medium grey, 7% biotite	L399354	-1	16
BV15-35	7.62	9.14	Grey	QZT	DP	2	20	Medium grey; 5% biotite	L399355	-1	14
BV15-35	9.14	10.67	Grey	QZT	DP	2	12	Light grey, very small sample	L399356	-1	9
BV15-35	10.67	12.19	Grey	QZT	DP	2	10	Light grey, very small sample	L399357	8	7
BV15-35	12.19	13.72	Grey	QZT	DP	2	3	Quartzite. Medium grey. Mica (muscovite) thick (1mm blebs), biotite 10%.	L399358	28	5
BV15-35	13.72	15.24	Grey	QZT	AM	2	5	Quartzite medium to dark grey. Biotite 5%, 30% medium grey QBS.	L399359	-1	5
BV15-35	15.24	16.76	Grey	QBS	AM	1	2	Grey. Quartz veins. 10% quartzite.	L399361	-1	9
BV15-35	16.76	18.29	Grey	QBS	AM	1	2	5% quartzite. Minor oxidation. Grey and brownish grey.	L399362	-1	8
BV15-35	18.29	19.81	Grey	QZT	AM	1	2	35% QBS. Medium grey. Quartz veins weak oxidation.	L399363	-1	6
BV15-35	19.81	21.34	Grey	QZT	AM	1	1	40% QBS. Medium grey. Quartz veins. Minor pyrite. Minor oxidation.	L399364	-1	21
BV15-35	21.34	22.86	Grey	QBS	AM	1	1	Medium grey. Quartz veins. Minor oxidation.	L399365	13	9
BV15-35	22.86	24.38	Grey	QBS	AM	1	1	10% Quartzite. Medium grey. Quartz veins. Minor oxidation.	L399366	37	20
BV15-35	24.38	25.91	Grey	QZT	AM	1	3	30% QBS. As above.	L399367	-1	19
BV15-35	25.91	27.43	Grey	QBS	AM	0	1	Dark greenish grey, minor py, bands to 2mm. Subhedral pyrite grains to 1.5m.	L399368	-1	6
BV15-35	27.43	28.96	Grey	QBS	AM	1	2	Dark grey, quartz veins. Minor oxidation. Minor Pyrite; 5% quartzite bands. Po blebs along margin greyish white quartz veinlet.	L399369	-1	8
BV15-35	28.96	30.48	Grey	QBS	AM	1	1	Dark greenish grey. Minor pyrite. Minor oxidation.	L399370	-1	9
BV15-35	30.48	32	Grey	QZT	AM	2	3	Quartzite (60%), QBS (40%). Minor pyrite, minor oxidation. Qtz veins.	L399371	8	48
BV15-35	32	33.53	Grey	QBS	AM	3	5	5% moderately oxidized quartz vein. 15% Quartzite.	L399372	18	136
BV15-35	33.53	35.05	Grey	QBS	AM	2	3	Water is found at 115'. Vein quartz fragments oxidized; orange limonite.	L399373	11	117
BV15-35	35.05	36.58	Grey	QBS	AM	1	6	Pyrite as irregular masses to 2mm across; oxidized vein quartz fragments as above.	L399374	36	141
BV15-35	36.58	38.1	Grey	QBS	AM	1	2	25% oxidized quartzite fragments.	L399375	36	57
BV15-35	38.1	39.62	Grey	QBS	AM	1	2	Quartzite 30%, QBS 70%.	L399376	15	47
BV15-35	39.62	41.15	Grey	QBS	AM	1	1	Less water, drier sample. 25% quartzite. Greenish grey chlorite alteration.	L399377	9	35
BV15-35	41.15	42.67	Grey	QBS	AM	1	3	Greenish grey, chlorite alteration. Quartz veins, pyrite.	L399378	-1	12
BV15-35	42.67	44.2	Grey	QBS	AM	1	2	Greenish grey, chlorite alteration. Quartz veins, pyrite. 3% quartzite.	L399379	-1	8
BV15-35	44.2	45.72	Grey	QBS	AM	2	2	Minor chlorite alteration. Quartz veins with pyrite, minor oxidation. 5% quartzite.	L399381	-1	10
BV15-35	45.72	47.24	Grey	VN	AM	2	80	Mainly quartz vein. 10% quartzite; 10% QBS. Traces sooty grey sulphide (? Sulphosalts).	L399382	830	290
BV15-35	47.24	48.77	Grey	QZT	AM	2	2	Quartzite. Minor quartz veins. Minor silicification.	L399383	1120	643
BV15-35	48.77	50.29	Grey	QZT	AM	1	5	Light grey, quartz + pyrite veins. Moderate silicification.	L399384	352	745
BV15-35	50.29	51.82	Grey	QZT	AM	1	2	Light grey to light greenish grey, quartz veins. 25% QBS.	L399385	53	102
BV15-35	51.82	53.34	Grey	QZT	AM	1	5		L399386	1040	4930
BV15-35	53.34	54.86	Grey	QZT	AM	1	10		L399387	2780	10000
BV15-35	54.86	56.39	Grey	QZT	AM	1	10		L399388	584	2810
BV15-35	56.39	57.91	Grey	QZT	AM	1	5	Pyrite (fine grained) disseminated/micro veins.	L399389	358	1560
BV15-35	57.91	59.44	Grey	QZT	AM	2	10		L399391	849	1890
BV15-35	59.44	60.96	Grey	QZT	AM	1	5	More indurated than above.	L399392	844	1000
BV15-35	60.96	62.48	Grey	QZT	AM	0	10	Some grey; very finely disseminated sulphides.	L399393	927	1530
BV15-35	62.48	64.01	Grey	QZT	AM	0	15	Some grey; very finely disseminated sulphides.	L399394	2130	1130
BV15-35	64.01	65.53	Grey	QZT	AM	1	5	Moderate vein content. Pyrite (cubic) in vein.	L399395	603	1220
BV15-35	65.53	67.06	Grey	QBS	AM	1	3	35% quartzite.	L399396	736	2320

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-35	67.06	68.58	Grey	QBS	AM	1	3	Dark grey, quartz + pyrite veins. Silicified; 30% quartzite.	L399397	870	2450
BV15-35	68.58	70.1	Grey	QZT	AM	1	3	35% QBS. Medium grey.	L399398	224	576
BV15-35	70.1	71.63	Grey	QBS	AM	0	5	Dark grey, quartz + pyrite veins. Cubic pyrite, silicified quartzite. 40% quartzite.	L399399	241	710
BV15-35	71.63	73.15	Grey	QZT	AM	0	5	Light grey silicified. Sooty, very fine grained pyrite, disseminated.	L399400	860	2640
BV15-35	73.15	74.68	Grey	QBS	AM	0	2	Dark greenish grey with grey silica quartz veins. 10% light grey quartzite.	L399401	578	1550
BV15-35	74.68	76.2	Grey	QBS	AM	1	2	25% quartzite which contains 3% disseminated pyrite as subhedral crystals.	L399402	430	1950
BV15-35	76.2	77.72	Grey	QBS	AM	0	3	Fine grained cubic pyrite. 30% quartzite.	L399403	257	1200
BV15-35	77.72	79.25	Grey	QZT	AM	0	3	White (milky) quartz veins. 40% QBS. Pyrite disseminated.	L399404	634	2370
BV15-35	79.25	80.77	Grey	QZT	AM	1	2	40% QBS. Vein quartz off-white, vitreous.	L399405	351	1300
BV15-35	80.77	82.3	Grey	QZT	AM	1	2	Quartzite (60%) QBS (40%). Quartz veins. Minor pyrite.	L399406	202	1750
BV15-35	82.3	83.82	Grey	QBS	AM	1	2	Light grey, quartz veins, disseminated pyrite 30% quartzite	L399407	187	1210
BV15-35	83.82	85.34	Grey	QBS	AM	1	2	Dark grey, quartz veins. Pyrite 40% quartzite.	L399408	332	1530
BV15-35	85.34	86.87	Grey	QZT	AM	1	2	Light grey, quartz veins, pyrite disseminated, pyrite in veins. 40% QBS.	L399409	129	575
BV15-35	86.87	88.39	Grey	QZT	AM	1	5	Mostly quartzite. QBS (30%), silicified. Bluish very fine disseminated sulphides present.	L399410	149	609
BV15-35	88.39	89.92	Grey	QBS	AM	1	3	Quartz vein with oyrite; light grey, silicification. 30% Quartzite.	L399411	144	703
BV15-35	89.92	91.44	Grey	QZT	AM	1	2	20% QBS.	L399412	930	2240
BV15-35	91.44	92.96	Grey	QBS	AM	0	6	Contact zone. Quartzite QBS. Silicification, quartz veins with pyrite. 30% quartzite vein quartz; vitreous, glassy.	L399413	214	868
BV15-35	92.96	94.49	Grey	QBS	AM	0	2	Dark grey QBS with white and grey quartz and pyrite veins. 30% quartzite.	L399414	439	1170
BV15-35	94.49	96.01	Grey	QBS	AM	0	3	Dark grey QBS with quartz and pyrite veins. Minor silicification. 25% Quartzite.	L399415	173	784
BV15-35	96.01	97.54	Grey	QBS	AM	0	2	Dark grey QBS with quartz and pyrite veins. Minor silicification. 25% Quartzite.	L399416	117	508
BV15-35	97.54	99.06	Grey	QZT	AM	0	3	Medium grey. 25% quartzite. Sericite alteration?	L399417	269	1220
BV15-35	99.06	100.58	Grey	QZT	AM	1	2	Greenish grey to light grey minor silicification. Pyrite dissemination.	L399418	303	1520
BV15-35	100.58	102.11	Grey	QBS	AM	0	2	20% quartzite.	L399419	123	571
BV15-35	102.11	103.63		QBS					No Sample		
BV15-35	103.63	105.16	Grey	QBS	DP	0	15		L399421	100	420
BV15-35	105.16	106.68	Grey	QBS	DP	0	10	Weak chlorite. Medium grey.	L399422	31	134
BV15-35	106.68	108.2	Grey	QBS	DP	0	5	Weak chlorite. Medium brownish grey. Huge pebble vein quartz with pyrite.	L399423	21	68
BV15-35	108.2	109.73		QBS	DP	0	5	Dark greenish grey, pyrite as subhedral cubes to 3mm; also dusty bluish grey mineral (?gn or aspy) in vein quartz which contain trace chlorite.	L399424	-1	19
BV15-35	109.73	111.25		QBS	DP	1	2	Much less vein quartz than above	L399425	-1	18
BV15-35	111.25	112.78		QBS	DP	1	1	Much less vein quartz than above	L399426	-1	19
BV15-35	112.78	114.3		QBS	DP	1	1		L399427	31	175
BV15-35	114.3	115.82		QBS	DP	1	1		L399428	-1	25
BV15-35	115.82	117.35		QBS	DP	1	0.5		L399429	-1	14
BV15-35	117.35	118.87	Brown	QBS	DP	1	0	Chlorite-quartz altered bands	L399431	-1	9
BV15-35	118.87	120.4	Brown	QBS	DP	1	0	pyrite subhedral grains to 0.5mm	L399432	11	106
BV15-35	120.4	121.92		QZT	DP	0	2	20% QBS; light grey	L399433	136	531
BV15-35	121.92	123.44	Grey	QZT	DP	0	2	5% QBS	L399434	-1	21
BV15-35	123.44	124.97	Grey	QZT	DP	0	1	Light Grey	L399435	-1	10
BV15-35	124.97	126.49	Grey	QZT	DP	0	0	Light Grey	L399436	6	17
BV15-35	126.49	128.02	Grey	QZT	DP	0	0	Light Grey	L399437	114	73
BV15-35	128.02	129.54	Grey	QZT	DP	0	3	Light Grey	L399438	18	23
BV15-35	129.54	131.06	Grey	QZT	DP	0	2	Light Grey	L399439	-1	10
BV15-35	131.06	132.59	Grey	QZT	DP	0	1	Light Grey	L399440	17	30
BV15-35	132.59	134.11	Grey	QZT	DP	0	2	Light Grey	L399441	-1	14
BV15-35	134.11	135.64	Grey	QZT	DP	0	2	Light Grey	L399442	-1	6
BV15-35	135.64	137.16	Grey	QZT	DP	0	1	Light Grey	L399443	11	36
BV15-35	137.16	138.68	Grey	QZT	DP	1	1	Light grey; pyrite crystals to 2mm	L399444	416	1850
BV15-35	138.68	140.21	Grey	QZT	DP	0	1	Light grey; pyrite crystals to 2mm	L399445	392	1640
BV15-35	140.21	141.73		QZT	DP	0	0	Light brownish grey; more biotite	L399446	48	218
BV15-35	141.73	143.26		QZT	DP	0	20	Off-white; 5% QBS	L399447	14	68
BV15-35	143.26	144.78		QZT	DP	0	15	Off-white 40% light brownish grey 60%	L399448	10	55
BV15-35	144.78	146.3		QZT	DP	1	20	off-white to light grey.	L399449	280	1310
BV15-35	146.3	147.83		QZT	DP	0	20	10% QBS; light grey	L399451	161	775
BV15-35	147.83	149.35		QZT	DP	0	20	Light grey; pyrite grains subhedral to 0.75mm. Similar to 465-470' depth above.	L399452	180	795
BV15-35	149.35	150.88		QZT	DP	1	3	Light grey; dusty bluish sooty sulphides present in trace amounts. Nice looking intervals.	L399453	175	845
BV15-35	150.88	152.4	Grey	QZT	DP	0	5	Light greenish grey, fine grained as above. Cube galena 1mm across.	L399454	93	419
BV15-35	152.4	153.92	Grey	QZT	DP	0	3	Light brownish grey fine grained.	L399455	111	553
BV15-35	153.92	155.45	Grey	QZT	DP	0	3	Light brownish grey fine grained.	L399456	69	312

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-35	155.45	156.97	Grey	QZT	DP	0	2	Light brownish grey fine grained.	L399457	346	3110
BV15-35	156.97	158.5	Grey	QZT	AM	0	1	Grey and light brownish grey. Quartz vein with pyrite.	L399458	58	303
BV15-35	158.5	160.02	Grey	QZT	AM	0	1	Dark grey and ight greenish grey. Quartz vein + pyrite (c)	L399459	155	680
BV15-35	160.02	161.54	Grey	QZT	AM	0	2	Dark grey and ight greenish grey. Quartz vein + pyrite (c)	L399461	286	2410
BV15-35	161.54	163.07	Grey	QZT	AM	0	3	Greenish grey, chlorite/sericite? alteration. Quartz and fine pyrite + galena (?) veins possibly cpy traces(?)	L399462	188	940
BV15-35	163.07	164.59	Grey	QZT	AM	0	2	Greenish grey, chlorite/sericite? alteration. Dark grey pieces. Milky quartz. Minor pyrite.	L399463	106	530
BV15-35	164.59	166.12	Grey	QZT	AM	0	1	Greenish grey, dark grey, chlorite/sericite, minor pyrite, quartz veins. Pyrite associated with chlorite/sericite (?) alteration. Possibly very minor galena.	L399464	20	122
BV15-35	166.12	167.64	Grey	QZT	AM	0	2	Greenish grey chlorite/sericite alteration pieces, dark grey with biotite (?). Milky quartz, light grey quartz with minor pyrite.	L399465	93	494
BV15-35	167.64	169.16	Grey	QZT	AM	0	2	Greenish grey chlorite/sericite alteration. Dark gray pices with brown, pyrite and po in quartz veins and chlorite/sericite alteration.	L399466	142	767
BV15-36	0	7.62		OB				No Sample			
BV15-36	7.62	9.14		QBS	DP	0	0	No chips viewed. None for the tray as minimal sample volume.	L399467	28	157
BV15-36	9.14	10.67	Grey	QBS	DP	4	40	Medium grey schist; not well washed.	L399468	20	202
BV15-36	10.67	12.19	Grey	QBS	DP	2	20	10% Quartzite	L399469	24	173
BV15-36	12.19	13.72	Grey	QBS	DP	2	10	very fine grained, dense, quartzite w. Sooty.	L399470	7	62
BV15-36	13.72	15.24	Grey	QBS	DP	2	5	very fine grained, dense, quartzite w. Sooty.	L399471	-1	28
BV15-36	15.24	16.76	Grey	QBS	DP	2	2	very fine grained, dense, quartzite w. Sooty.	L399472	11	63
BV15-36	16.76	18.29	Grey	QBS	DP	2	1	very fine grained, dense, quartzite w. Sooty.	L399473	7	49
BV15-36	18.29	19.81	Grey	QBS	DP	2	1	very fine grained, dense, quartzite w. Sooty.	L399474	12	46
BV15-36	19.81	21.34	Grey	QBS	DP	1	0	very fine grained, dense, quartzite w. Sooty.	L399475	6	30
BV15-36	21.34	22.86	Grey	QBS	DP		0.5	very fine grained, dense, quartzite w. Sooty.	L399476	15	43
BV15-36	22.86	24.38	Grey	QBS	DP	1	1	very fine grained, dense, quartzite w. Sooty.	L399477	8	33
BV15-36	24.38	25.91	Grey	QBS	DP	1	0	very fine grained, dense, quartzite w. Sooty.	L399478	20	78
BV15-36	25.91	27.43	Grey	QBS	DP	1	0	very fine grained, dense, quartzite w. Sooty.	L399479	8	81
BV15-36	27.43	28.96		QBS	DP	1	0	10% Quartzite	L399481	17	88
BV15-36	28.96	30.48		QBS	DP	1	0	20% Quartzite	L399482	18	142
BV15-36	30.48	32		QBS	DP	2	0	20% Quartzite	L399483	-1	101
BV15-36	32	33.53		QZT	DP	2	1	10% QBS	L399484	88	129
BV15-36	33.53	35.05		QZT	DP	3	0	20% QBS	L399485	27	95
BV15-36	35.05	36.58		QZT	DP	2	0		L399486	103	85
BV15-36	36.58	38.1		QZT	DP	0	0		L399487	93	113
BV15-36	38.1	39.62		QZT	DP	2	0		L399488	258	534
BV15-36	39.62	41.15		QZT	DP	2	0		L399489	358	507
BV15-36	41.15	42.67		QZT	DP	1	0		L399491	155	361
BV15-36	42.67	44.2		QZT	DP	1	10	Fine seams pyrite in vein	L399492	328	790
BV15-36	44.2	45.72		QZT	DP	1	10	Pyrite disseminated in vein material	L399493	494	2150
BV15-36	45.72	47.24		QBS	DP	1	4	20% quartzite	L399494	220	161
BV15-36	47.24	48.77		QBS	DP	1	4	10% Quartzite	L399495	45	58
BV15-36	48.77	50.29		QBS	DP	1	1	WET	L399496	39	53
BV15-36	50.29	51.82		QBS	DP	1	1	DRY	L399497	16	16
BV15-36	51.82	53.34		QBS	DP	1	1		L399498	16	27
BV15-36	53.34	54.86		QZT	DP	0	1	20% QBS	L399499	46	55
BV15-36	54.86	56.39		QBS	DP	0	0	30% Quartzite	L399500	47	72
BV15-36	56.39	57.91		QZT	DP	1	2	Light grey, fine grained, very fine disseminated pyrite	B00261701	94	199
BV15-36	57.91	59.44		QZT	DP	1	0	Light grey, fine grained, very fine disseminated pyrite	B00261702	1040	258
BV15-36	59.44	60.96		QBS	DP	0	2	30% Quartzite	B00261703	407	113
BV15-36	60.96	62.48		QZT	DP	1	1	10% QBS	B00261704	197	153
BV15-36	62.48	64.01		QZT	DP	1	1	10% QBS	B00261705	182	195
BV15-36	64.01	65.53		QZT	DP	1	2	10% QBS	B00261706	137	160
BV15-36	65.53	67.06		QBS	DP	1	1	5% Quartzite	B00261707	413	272
BV15-36	67.06	68.58		QBS				No Sample			
BV15-36	68.58	70.1		QBS	DP	1	1	20% Quartzite	B00261709	51	151
BV15-36	70.1	71.63		QBS	DP	1	1	10% Quartzite	B00261710	22	60
BV15-36	71.63	73.15		QBS	DP	1	0.5	30% quartzite; fine grained, light grey	B00261711	58	30
BV15-36	73.15	74.68		QBS	DP	0	5	light grey	B00261712	56	106
BV15-36	74.68	76.2		QBS	DP	0	5	light grey	B00261713	48	57
BV15-36	76.2	77.72		QBS	DP	1	2	light grey	B00261714	38	62

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-36	77.72	79.25		QBS	DP	0	1	light grey	800261715	32	45
BV15-36	79.25	80.77		QBS	DP	0	4		800261716	39	36
BV15-36	80.77	82.3		QBS	DP	0	1	light brownish grey; fine grained	800261717	18	25
BV15-36	82.3	83.82		QBS	DP	0	0	light grey	800261718	11	16
BV15-36	83.82	85.34		QBS	DP	0	1	light grey	800261719	58	18
BV15-36	85.34	86.87		QZT	DP	0	1	light grey; 30% QBS	800261721	214	37
BV15-36	86.87	88.39		QBS	DP	0	0.5	30% Quartzite	800261722	34	27
BV15-36	88.39	89.92		QZT	DP	1	0.5		800261723	43	26
BV15-36	89.92	91.44		QZT	DP	1	0.5		800261724	254	1820
BV15-36	91.44	92.96		QZT	DP	4	2	10% QBS	800261725	168	616
BV15-36	92.96	94.49		QZT	DP	1	2	5% QBS	800261726	234	474
BV15-36	94.49	96.01		QBS	DP	1	1		800261727	33	96
BV15-36	96.01	97.54		QBS	DP		1		800261728	15	41
BV15-36	97.54	99.06		QBS	DP		0.5	10% indurated quartzite	800261729	14	49
BV15-36	99.06	100.58		QBS	DP	1	0		800261731	12	32
BV15-36	100.58	102.11		QBS	DP	1	0		800261732	16	31
BV15-36	102.11	103.63		QBS	DP	1	2		800261733	20	23
BV15-36	103.63	105.16		QBS	DP	1	1	Brownish grey, fine grained	800261734	19	39
BV15-36	105.16	106.68		QBS	DP	1	0.5	Brownish grey, fine grained	800261735	11	36
BV15-36	106.68	108.2	Grey	QBS	AM	1	0.5	Brownish grey, fine grained	800261736	6	16
BV15-36	108.2	109.73	Grey	QBS	AM	0	1	Brownish grey, fine grained	800261737	10	9
BV15-36	109.73	111.25	Grey	QBS	AM	0	2	Brownish grey, greenish grey; quartz was with pyrite, cpy, po	800261738	5	19
BV15-36	111.25	112.78	Grey	QBS	AM	0	2	Brownish grey, greenish grey; quartz was with pyrite, cpy, po	800261739	9	32
BV15-36	112.78	114.3	Grey	QBS	AM	0	3	Greenish grey to brownish grey. Pyrite associated with quartz vein.	800261740	30	44
BV15-36	114.3	115.82	Grey	QBS	AM	0	1	Brownish grey, quartz vein with pyrite, minor po	800261741	-1	12
BV15-36	115.82	117.35	Grey	QBS	AM	0	1	Greenish grey to brownish grey. Chl/ser (?). Py associated with quartz vein and chl/ser alteration	800261742	33	311
BV15-36	117.35	118.87	Grey	QZT	AM	0	5	Quartzite with py (cubic). Light grey. Mineralized.	800261743	585	1200
BV15-36	118.87	120.4	Grey	QBS	AM	0	1	Brownish grey, minor greenish, minor quartz + pyrite veins.	800261744	48	92
BV15-36	120.4	121.92	Grey	QBS	AM	0	1	Brownish grey, minor greenish (chl alteration). 10% quartzite. Minor quartz + pyrite veins.	800261745	14	14
BV15-36	121.92	123.44	Grey	QBS	AM	0	1	Brownish grey, minor greenish, chlorite alteration + pyrite. Qtz + Py veins. Moderately magnetic.	800261746	-1	11
BV15-36	123.44	124.97	Grey	QBS	AM	0	0.5	Greenish grey and brownish grey. Moderate chlorite alteration. Minor pyrite.	800261747	-1	9
BV15-36	124.97	126.49	Grey	QBS	AM	0	0.5	Brownish grey and greenish grey, chlorite alteration	800261748	-1	9
BV15-36	126.49	128.02	Grey	QBS	AM	0	0.5	Brownish grey, minor quartz and pyrite veins/chlorite alteration	800261749	-1	8
BV15-36	128.02	129.54	Grey	QBS	AM	0	0.5	Brownish grey, minor quartz and pyrite veins/chlorite alteration	800261751	-1	8
BV15-36	129.54	131.06	Grey	QBS	AM	0	0.5	Brownish grey, minor quartz and pyrite veins/chlorite alteration	800261752	7	12
BV15-36	131.06	132.59	Grey	QBS	AM	0	1	Brownish grey and greenish grey, chlorite at minor quartz veins.	800261753	-1	12
BV15-36	132.59	134.11	Grey	QBS	AM	0	2	Brownish grey, chlorite alteration, quartz + pyrite veins, weakly magnetic.	800261754	148	1750
BV15-36	134.11	135.64	Grey	QBS	AM	0	1	Brownish grey, chlorite alteration. Quartz veins with pyrite, po in veins	800261755	-1	21
BV15-36	135.64	137.16	Grey	QBS	AM	0	1	Brownish grey, chlorite alteration, minor quartz + pyrite veins.	800261756	-1	10
BV15-36	137.16	138.68	Grey	QBS	AM	0	1	Brownish grey, chlorite alteration, minor quartz + pyrite veins.	800261757	-1	19
BV15-36	138.68	140.21	Grey	QBS	AM	0	2	Brownish grey + greenish grey, chlorite alteration	800261758	10	20
BV15-36	140.21	141.73	Grey	QZT	AM	1	3	5% QBS	800261759	50	117
BV15-36	141.73	143.26	Grey	QZT	AM	0	2	3% QBS	800261761	7	6
BV15-36	143.26	144.78	Grey	QZT	AM	0	1	Light grey	800261762	7	9
BV15-36	144.78	146.3	Grey	QBS	AM	0	2	Brownish grey; minor quartz veins and chlorite	800261763	17	14
BV15-36	146.3	147.83	Grey	QBS	AM	0	1	Brownish grey, 5% quartzite	800261764	-1	6
BV15-36	147.83	149.35	White	QZT	AM	0	1	White to light greenish grey, minor QBS, min py	800261765	11	9
BV15-36	149.35	150.88	Grey	QZT	AM	0	3	Light grey to medium grey to white; 5% QBS	800261766	143	522
BV15-36	150.88	152.4	Grey	QBS	AM	0	1	Greenish grey, minor veins. 5% quartzite	800261767	31	81
BV15-36	152.4	153.92	Grey	QBS	AM	0	0.5	Brownish grey and greenish grey	800261768	6	8
BV15-36	153.92	155.45	Grey	QBS	AM	0	1	Brownish grey, minor chlorite and qtz+py veins	800261769	12	32
BV15-36	155.45	156.97	Grey	QBS	AM	0	1	Brownish grey, minor chlorite and qtz+py veins	800261770	47	91
BV15-36	156.97	158.5	Grey	QBS	AM	0	1	Brownish grey, minor chlorite alteration, minor quartz veins. Weak Pyrite and po traces.	800261771	7	3
BV15-36	158.5	160.02	Grey	QBS	DP	0	2	As above except no po seen	800261772	12	18
BV15-36	160.02	161.54	Grey	QZT	DP	0	0.5	Light grey, fine grained.	800261773	10	11
BV15-36	161.54	163.07	Grey	QZT	DP	0	0.5	30% QBS	800261774	12	9
BV15-36	163.07	164.59		QBS	DP	0	0.5	Light brownish grey. 25% quartzite; py bleb 1.5mm	800261775	129	334
BV15-36	164.59	166.12	Grey	QZT	DP	0	0.5	30% QBS; occ. Py-rich (30%) grains to 5mm across.	800261776	9	11
BV15-36	166.12	167.64		QZT	DP	0	0.5	Light greenish grey, fine grained.	800261777	25	20

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-36	167.64	169.16		QZT	DP	1	0	Light greenish grey, fine grained.	800261778	52	191
BV15-36	169.16	170.69		QZT	DP	0	0	Light greenish grey, fine grained.	800261779	7	15
BV15-37	0	7.62		OB					No Sample		
BV15-37	7.62	9.14	Grey	QBS	DP	2	0	Light grey, fine grained.	800261781	-1	9
BV15-37	9.14	10.67	Grey	QBS	DP	2	0	Light grey, fine grained.	800261782	-1	5
BV15-37	10.67	12.19	Grey	QBS	DP	2	3	Medium grey, fine grained; vein quartz fragments pervasive, laminations	800261783	-1	9
BV15-37	12.19	13.72	Grey	QZT	DP	3	0.5	30% QBS	800261784	-1	11
BV15-37	13.72	15.24		QBS	DP	2	1	20% Quartzite	800261785	-1	7
BV15-37	15.24	16.76		QBS	DP	2	0.5	medium grey, fine grained	800261786	-1	8
BV15-37	16.76	18.29	Grey	QBS	DP	2	0	10% quartzite	800261787	-1	9
BV15-37	18.29	19.81		QZT	DP	3	1	5% QBS (chlorite alteration, weakly)	800261788	25	240
BV15-37	19.81	21.34		QZT	DP	2	2		800261789	10	140
BV15-37	21.34	22.86	Grey	QBS	DP	2	0.5	medium grey	800261791	6	29
BV15-37	22.86	24.38		QBS	DP	2	0		800261792	-1	21
BV15-37	24.38	25.91		QBS	DP	2	0		800261793	-1	21
BV15-37	25.91	27.43		QBS	DP	2	0		800261794	-1	33
BV15-37	27.43	28.96		QBS	DP	2	0.5	20% quartzite	800261795	43	24
BV15-37	28.96	30.48		QBS	DP	2	0		800261796	-1	35
BV15-37	30.48	32		QBS	DP	1	0	10% quartzite	800261797	-1	46
BV15-37	32	33.53		QBS	DP	2	0		800261798	-1	31
BV15-37	33.53	35.05	Grey	QBS	DP	1	0	40% quartzite; greenish grey	800261799	-1	29
BV15-37	35.05	36.58		QBS	DP	1	0	5% oxidized quartzite chips	800261800	-1	63
BV15-37	36.58	38.1		QBS	DP	1	0	30% quartzite	800261801	-1	77
BV15-37	38.1	39.62		QBS	DP	1	0		800261802	-1	15
BV15-37	39.62	41.15		QBS	DP	1	0		800261803	-1	10
BV15-37	41.15	42.67		QBS	DP	1	0		800261804	-1	7
BV15-37	42.67	44.2		QBS	DP	1	0	weak epidote alteration	800261805	21	56
BV15-37	44.2	45.72	White	QBS	DP	1	0	30% off-white quartzite	800261806	-1	38
BV15-37	45.72	47.24		QBS	DP	1	0	10% quartzite	800261807	-1	17
BV15-37	47.24	48.77		QZT	DP	1	0	10% QBS; likely casing not set deep enough or fault zone; bit grabbing, slow	800261808	-1	27
BV15-37	48.77	50.29		QBS	DP	0	0	10% quartzite; pyrite as fracture coating patches to 2mm across.	800261809	31	22
BV15-37	50.29	51.82		QBS	DP	1	0		800261810	13	19
BV15-37	51.82	53.34		QBS	DP	1	0	3% quartzite; soupy; small sample	800261811	-1	38
BV15-37	53.34	54.86		QBS	DP	1	0.3		800261812	-1	37
BV15-37	54.86	56.39		QBS	DP	1	0	Likely very weak epidote - altered.	800261813	-1	52
BV15-37	56.39	57.91	Grey	QBS	DP	1	0.5	Another possible fault; broken ground, high water flow as @ 160' depth. QBS + Quartzite (20%). Greenish grey. Chlorite altered. Minor pyrite.	800261814	55	38
BV15-37	57.91	59.44	Grey	QBS	AM	1	1	Another possible fault; broken ground, high water flow as @ 160' depth. QBS + Quartzite (20%). Greenish grey. Chlorite altered. Minor pyrite.	800261815	9	24
BV15-37	59.44	60.96	Grey	QBS	AM	1	1	Another possible fault; broken ground, high water flow as @ 160' depth. QBS + Quartzite (20%). Greenish grey. Chlorite altered. Minor pyrite.	800261816	324	9
BV15-37	60.96	62.48	Grey	QBS	AM	1	1	Quartzite (20%) + QBS (80%). Light grey to greenish grey.	800261817	-1	8
BV15-37	62.48	64.01	Grey	QBS	AM	1	0.3	Greenish grey, chlorite altered, minor pyrite.	800261818	14	5
BV15-37	64.01	65.53	Grey	QBS	AM	0	0.3	QBS (80%), Qtzt (20%). Chlorite alteration, very minor pyrite	800261819	26	9
BV15-37	65.53	67.06	Grey	QZT	AM	0	0.3	QBS (60%), Qtzt (40%). Chlorite alteration, very minor pyrite	800261821	-1	11
BV15-37	67.06	68.58	Grey	QZT	AM	0	0.3	QBS (20%), Qtzt (20%). Chlorite alteration, very minor pyrite	800261822	32	37
BV15-37	68.58	70.1	Grey	QBS	AM	0	0.5	QBS (20%), Qtzt (20%). Chlorite alteration, minor pyrite	800261823	20	23
BV15-37	70.1	71.63	Grey	QBS	AM	0	0	QBS (20%), Qtzt (20%). Chlorite alteration, minor pyrite	800261824	5	25
BV15-37	71.63	73.15	Grey	QZT	AM	0	2	Moderate pyrite	800261825	67	650
BV15-37	73.15	74.68	Grey	QZT	AM	0	2		800261826	496	3340
BV15-37	74.68	76.2	Grey	QZT	AM	1	3		800261827	177	1530
BV15-37	76.2	77.72	Grey	QZT	AM	1	2		800261828	634	8450
BV15-37	77.72	79.25	Grey	QZT	AM	1	5	pyrite subhedral crystals (drill hole is caving in)	800261829	319	2360
BV15-37	79.25	80.77	Grey	QZT	AM	1	4	Mostly quartzite, 25% QBS, quartz + pyrite veins, minor ? (sph?)	800261831	323	1750
BV15-37	80.77	82.3	Grey	QZT	AM	1	1	25% QBS	800261832	152	1210
BV15-37	82.3	83.82	Grey	QZT	AM	1	0.5	30% QBS	800261833	98	636
BV15-37	83.82	85.34	Grey	QZT	AM	1	0.3	30% QBS	800261834	66	274
BV15-37	85.34	86.87	Grey	QZT	AM	1	2	quartzite with quartz veins, minor chlorite alteration, sphalerite?	800261835	53	276
BV15-37	86.87	88.39	Grey	QZT	AM	1	2	quartzite with quartz veins, minor chlorite alteration, sphalerite?	800261836	619	661

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-37	88.39	89.92	Grey	QZT	AM	1	3	quartzite with quartz veins, minor chlorite alteration, sphalerite?	800261837	352	5490
BV15-37	89.92	91.44	Grey	QZT	AM	1	1	quartzite with quartz veins, minor chlorite alteration, sphalerite?	800261838	126	2910
BV15-37	91.44	92.96	Grey	QZT	AM	1	2	Quartzite, minor QBS, quartz + pyrite veins.	800261839	136	1120
BV15-37	92.96	94.49	Grey	QZT	AM	1	5	Quartzite, quartz + pyrite veins, sphalerite vein?	800261840	170	897
BV15-37	94.49	96.01	Grey	QZT	AM	1	10	Quartzite, quartz + pyrite veins, sphalerite vein?	800261841	206	2100
BV15-37	96.01	97.54	Grey	QZT	AM	1	7	Quartzite, quartz + pyrite veins, sphalerite vein?	800261842	204	2200
BV15-37	97.54	99.06	Grey	QZT	AM	1	3	Quartzite (80%), QBS (20%), Quartz veins, sphalerite, pyrite	800261843	96	791
BV15-37	99.06	100.58	Grey	QBS	AM	1	2	Quartzite (80%), QBS (20%), Quartz veins, pyrite, minor sphalerite	800261844	49	447
BV15-37	100.58	102.11	Grey	QBS	AM	1	3	Quartzite (40%), grey quartz veins with pyrite, sphalerite	800261845	44	307
BV15-37	102.11	103.63	Grey	QBS	AM	0	5	QBS, dark greenish grey, quartz veins + pyrite, sphalerite	800261846	33	328
BV15-37	103.63	105.16	Grey	QBS	AM	1	1	QBS, minor quartzite. Dark greenish grey; pyrite, sphalerite	800261847	12	92
BV15-37	105.16	106.68	Grey	QBS	DP	1	2	Dark greenish grey; very fine grained.	800261848	6	89
BV15-37	106.68	108.2		QZT	DP	1	1	30% QBS; pyrite bands 1mm along quartz veinlet margin	800261849	5	39
BV15-37	108.2	109.73		QBS	DP	1	0.5	brown biotite-altered bands to 2mm wide.	800261851	-1	41
BV15-37	109.73	111.25		QBS	DP	1	0.3	brown biotite-altered bands to 2mm wide.	800261852	-1	34
BV15-37	111.25	112.78		QBS	DP	1	0.2	brown biotite-altered bands to 2mm wide; pyrite blebs to 2mm	800261853	44	217
BV15-37	112.78	114.3		QZT	DP	0	0.5	Nice looking; pyrite disseminated subhedral grains and traces	800261854	45	368
BV15-37	114.3	115.82		QZT	DP	1	0.5	Not as good looking as above	800261855	42	328
BV15-37	115.82	117.35		QZT	DP	2	0.5	Nice-looking; weak chlorite and biotite. ~0.5% pyrite very fine disseminated, sooty.	800261856	205	2600
BV15-37	117.35	118.87		QZT	DP	1	0.5	***Very good looking***	800261857	554	4810
BV15-37	118.87	120.4		QZT	DP	1	0.5	Weak clay - (sericite?) altered locally.	800261858	121	1290
BV15-37	120.4	121.92		QZT	DP	1	1	15% QBS	800261859	21	145
BV15-37	121.92	123.44		QZT	DP	1	0.5	5% QBS, not sericite altered	800261861	53	355
BV15-37	123.44	124.97		QZT	DP	1	0.5	5% QBS, not sericite altered	800261862	39	250
BV15-37	124.97	126.49	Grey	QZT	DP	1	0.5	Light greenish grey, finely granular	800261863	68	425
BV15-37	126.49	128.02	Grey	QZT	DP	0	0	Light greenish grey, finely granular	800261864	73	360
BV15-37	128.02	129.54	Grey	QZT	DP	0	0	Light greenish grey, finely granular	800261865	15	98
BV15-37	129.54	131.06	Grey	QZT	DP	1	0	Light greenish grey, finely granular	800261866	-1	42
BV15-37	131.06	132.59	Grey	QZT	DP	1	0	Light grey, more silica indurated	800261867	630	2560
BV15-37	132.59	134.11	Grey	QZT	DP	1	2	Light grey, more silica indurated	800261868	553	2700
BV15-37	134.11	135.64	Grey	QZT	DP	1	0	Light grey, more silica indurated	800261869	492	2460
BV15-37	135.64	137.16	Grey	QZT	DP	0	0.5	Light grey, more silica indurated	800261870	301	1280
BV15-37	137.16	138.68	Grey	QZT	DP	1	0	Light grey, more silica indurated	800261871	48	121
BV15-37	138.68	140.21		QZT	DP	1	0	457' possible fault; drill but dropped ~25cm very quickly. Two large pebble sized chips kept as representative sample.	800261872	12	167
BV15-38	0	9.14		OB					No Sample		
BV15-38	9.14	10.67	Grey	QBS	DP	1	0.5		800261873	-1	11
BV15-38	10.67	12.19	Grey	QBS	DP	1	0	medium greyish green, fine grained	800261874	-1	11
BV15-38	12.19	13.72		QZT	DP	2	0	medium greyish green, fine grained	800261875	-1	24
BV15-38	13.72	15.24		QZT	DP	2	0		800261876	-1	21
BV15-38	15.24	16.76		QZT	DP	2	0		800261877	-1	20
BV15-38	16.76	18.29		QBS	DP	2	0	40% quartzite	800261878	-1	18
BV15-38	18.29	19.81		QZT	DP	2	0	20% QBS	800261879	-1	16
BV15-38	19.81	21.34		QZT	DP	2	0		800261881	-1	9
BV15-38	21.34	22.86		QZT	DP	3	0		800261882	-1	11
BV15-38	22.86	24.38		QZT	DP	2	0	10% QBS	800261883	-1	9
BV15-38	24.38	25.91		QZT	DP	2	0	30% QBS	800261884	-1	14
BV15-38	25.91	27.43		QZT	DP	2	0	15% QBS	800261885	-1	7
BV15-38	27.43	28.96		QZT	DP	1	0	10% QBS	800261886	-1	6
BV15-38	28.96	30.48		QZT	DP	1	0.5	10% QBS; weak epidote	800261887	-1	5
BV15-38	30.48	32		QZT	DP	1	1	10% QBS; weak epidote alteration	800261888	-1	-1
BV15-38	32	33.53		QZT	DP	1	0	30% QBS	800261889	-1	4
BV15-38	33.53	35.05		QBS	AM	1	1	20% quartzite. Pyrite associated with biotite (?)	800261891	-1	27
BV15-38	35.05	36.58		QBS	AM	0	2	QBS with quartz veins and pyrite in veins and biotite	800261892	-1	7
BV15-38	36.58	38.1		QBS	AM	0	1	QBS with quartz veins and pyrite in veins and biotite	800261893	6	6
BV15-38	38.1	39.62		QBS	AM	0	0.5	QBS, epidote, chlorite, pyrite	800261894	-1	5
BV15-38	39.62	41.15		QBS	AM	0	1	QBS, minor pyrite and quartz veins	800261895	-1	20
BV15-38	41.15	42.67		QBS	AM	0	1	QBS, minor pyrite and quartz veins	800261896	-1	12
BV15-38	42.67	44.2	Green	QBS	AM	1	1	Minor epidote, chlorite altered. Pyrite associated with biotite (quartz veins)	800261897	-1	25
BV15-38	44.2	45.72	Green	QBS	AM	1	1	QBS with minor quartz + pyrite veins	800261898	-1	15

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-38	45.72	47.24	Green	QBS	AM	1	1	QBS 80%, Quartzite 10%. Chlorite, epidote, pyrite, quartz veins	800261899	-1	74
BV15-38	47.24	48.77	Green	QBS	AM	0	0	QBS, minor epidote and pyrite	800261900	-1	89
BV15-38	48.77	50.29	Green	QBS	AM	0	0.5	QBS (80%), Quartzite (20%), chlorite, epidote, pyrite	800261901	-1	28
BV15-38	50.29	51.82	Green	QBS	AM	1	1	QBS, minor chlorite	800261902	-1	7
BV15-38	51.82	53.34	Green	QBS	AM	1	0.5	QBS 60%, Quartzite 40% , minor chlorite, quartz + pyrite vein	800261903	-1	9
BV15-38	53.34	54.86	Green	QBS	AM	0	0.5	QBS, minor chlorite	800261904	-1	9
BV15-38	54.86	56.39	Green	QBS	AM	1	1	QBS 80%, Quartzite 20%, quartz veins	800261905	-1	8
BV15-38	56.39	57.91	Grey	QBS	AM	1	1	Quartzite 30%, quartz veins + minor ..?	800261906	-1	8
BV15-38	57.91	59.44	Grey	QBS	AM	1	1	40% quartzite	800261907	-1	5
BV15-38	59.44	60.96	Grey	QZT	AM	1	1	30% QBS	800261908	-1	10
BV15-38	60.96	62.48	Green	QBS	AM	0	1	QBS 70%, Quartzite 30%, quartz + pyrite veins	800261909	-1	8
BV15-38	62.48	64.01	Grey	QZT	AM	1	2	Quartzite 70%, QBS 30%, quartz + pyrite veins	800261910	-1	7
BV15-38	64.01	65.53	Grey	QZT	AM	0	3	Quartzite 70%, QBS 30%, quartz + pyrite veins	800261911	-1	5
BV15-38	65.53	67.06	Grey	QBS	AM	1	1	Quartzite (25%), quartz veins + pyrite. Fault (?) caving hole in at 215'.	800261912	6	48
BV15-38	67.06	68.58	Grey	QZT	AM	1	2	Quartzite 60%, QBS 40%, quartz + pyrite veins, chlorite alteration	800261913	-1	44
BV15-38	68.58	70.1	Green	QZT	AM	1	1	Mostly quartzite (80%), QBS 20%, quartz + pyrite veins	800261914	-1	17
BV15-38	70.1	71.63	Green	QBS	AM	1	1	Minor oxides. Quartz + pyrite veins.	800261915	-1	26
BV15-38	71.63	73.15	Grey	QBS	AM	1	1	chlorite altered, minor epidote. Pyrite + quartz veins.	800261916	-1	25
BV15-38	73.15	74.68	Grey	QBS	AM	1	1	QBS 80%, Quartzite 20%, quartz + pyrite veins....?	800261917	-1	15
BV15-38	74.68	76.2	Grey	QBS	AM	1	1	Quartzite 80%, QBS 20%, Quartz + pyrite veins	800261918	-1	12
BV15-38	76.2	77.72	Grey	QZT	AM	2	2	Quartzite with quartz + pyrite veins, 40% QBS, minor oxidation	800261919	-1	19
BV15-38	77.72	79.25	Grey	QBS	AM	1	1	QBS (80%), Quartzite (20%), quartz + pyrite veins	800261921	-1	18
BV15-38	79.25	80.77	Grey	QBS	AM	1	1	40% Quartzite, minor biotite. Chlorite alteration. Quartz + pyrite veins	800261922	-1	36
BV15-38	80.77	82.3	Grey	QBS	AM	1	2	40% Quartzite, minor biotite. Chlorite alteration. Quartz + pyrite veins	800261923	-1	18
BV15-38	82.3	83.82	Grey	QBS	AM	1	2	Minor chlorite alteration. Pyrite associated with biotite and quartz veins.	800261924	-1	14
BV15-38	83.82	85.34	Grey	QBS	AM	1	2	QBS 70%, Quartzite 30%. Quartz veins with pyrite. Chlorite alteration.	800261925	-1	22
BV15-38	85.34	86.87	Grey	QZT	AM	0	1	Banded quartzite. Biotite bands. Minor chlorite alteration.	800261926	-1	18
BV15-38	86.87	88.39	Grey	QZT	AM	0	2	Banded quartzite. Quartz + pyrite veins. Minor chlorite alteration. Silica?	800261927	-1	13
BV15-38	88.39	89.92	Grey	QZT	AM	1	3	Light grey quartzite with pyrite (sph?). Silicification (?) and sericite alteration (?). Probably mineralized zone.	800261928	-1	7
BV15-38	89.92	91.44	Grey	QZT	AM	1	2	Quartzite 80%, QBS 20%, minor chlorite alteration. Pyrite in veins.	800261929	-1	11
BV15-38	91.44	92.96	Grey	QBS	AM	0	2	Light grey, 30% quartzite, quartz + pyrite veins. Silica (?).	800261931	-1	11
BV15-38	92.96	94.49	Grey	QBS	AM	1	1	Light grey. Quartzite 40%. Quartz + pyrite veins.	800261932	-1	25
BV15-38	94.49	96.01	Grey	QZT	AM	0	3	Light grey. Quartz + pyrite veins. Silicification (?). Sericite (?).	800261933	-1	14
BV15-38	96.01	97.54	Grey	QZT	AM	0	2	Light grey. Minor biotite. Quartz + pyrite veins.	800261934	-1	22
BV15-38	97.54	99.06	Grey	QBS	AM	0	0.5	QBS (80%), Quartzite (20%). Minor chlorite alteration.	800261935	-1	15
BV15-38	99.06	100.58	Grey	QBS	AM	0	0.5	Dark green. Minor quartz veins + pyrite.	800261936	-1	10
BV15-38	100.58	102.11	Green	QBS	AM	0	0.5	Dark green. Minor quartz veins + pyrite.	800261937	-1	8
BV15-38	102.11	103.63	Grey	QBS	AM	0	0.5	QBS, minor quartzite (20%). Chlorite alteration. Minor pyrite.	800261938	-1	3
BV15-38	103.63	105.16	Grey	QBS	AM	0	0.5		800261939	-1	9
BV15-38	105.16	106.68	Green	QBS	AM	0	0.5	Dark green; minor chlorite alteration, pyrite associated with biotite	800261940	7	10
BV15-38	106.68	108.2	Green	QBS	AM	0	0.7	Dark green; minor chlorite alteration, pyrite on fracture surfaces.	800261941	-1	9
BV15-38	108.2	109.73	Green	QBS	DP	1	0.5	Dark green; minor chlorite alteration, pyrite on fracture surfaces, quartzite 15%	800261942	-1	9
BV15-38	109.73	111.25	Green	CS	DP	0	0.5	Dark green; pyrite as blebs. 25% chlorite	800261943	-1	4
BV15-38	111.25	112.78	Brown	QZT	DP	1	0.5	Bit change at 365'. 20% QBS. 10% chlorite schist in this interval; light grey-brown.	800261944	-1	17
BV15-38	112.78	114.3	Brown	QBS	DP	0	1	30% quartzite. Medium greyish brown	800261945	-1	10
BV15-38	114.3	115.82	Brown	QBS	DP	1	1	20% quartzite. Medium greyish brown. Possible trace blue/grey aspy? In quartz veinlet.	800261946	-1	7
BV15-38	115.82	117.35		QZT	DP	1	0.5	30% QBS	800261947	-1	8
BV15-38	117.35	118.87		QZT	DP	1	0.5	40% QBS; less chlorite altered than above. Also weakly epidote altered.	800261948	-1	7
BV15-38	118.87	120.4	Grey	QZT	DP	0	0.5	30% QBS. Medium brownish grey colour. Py as subround blebs to 1mm across.	800261949	-1	11
BV15-38	120.4	121.92	Grey	QZT	DP	0	0.1	40% QBS. As above except more QBS.	800261951	-1	9
BV15-38	121.92	123.44		QZT	DP	0	0.2	10% QBS. Weakly epidote altered. Pyrite as irregular masses rarely to 3mm across. Fine biotite rich bands 1.5mm wide through quartzite.	800261952	-1	3
BV15-38	123.44	124.97		QZT	DP	1	0.3	5% QBS.	800261953	-1	22
BV15-38	124.97	126.49		QZT	DP	1	0.5	15% QBS. 1cm quartzite fragment in chips.	800261954	-1	19
BV15-38	126.49	128.02	Brown	QZT	DP	1	0.3	30% dark greyish brown QBS. Pyrite as euhedral grain max 1.5mm across to locally sooty very fine disseminated.	800261955	-1	6
BV15-38	128.02	129.54	Brown	QZT	DP	0	0.3	QBS as above	800261956	-1	11
BV15-38	129.54	131.06	Brown	QZT	DP	0	0.2	30% QBS as for 415-420'. Trace bluish aspy?	800261957	-1	9
BV15-38	131.06	132.59	Brown	QZT	DP	1	0.5	20% dark grey brown QBS	800261958	-1	11
BV15-39	0	1.52		OB					No Sample		

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-39	1.52	3.05	Grey	QZT	AM	3	1	Quartzite and overburden. Moderate to intense oxidation. Chlorite alteration?	800261959	9	9
BV15-39	3.05	4.57	Grey	QZT	AM	2	1	Quartzite, ox. Minor quartz veins. Brownish grey.	800261961	7	10
BV15-39	4.57	6.1	Grey	QZT	AM	2	1	Quartzite (80%), QBS (20%). Chlorite alteration. Quartz veins.	800261962	-1	18
BV15-39	6.1	7.62	Grey	QZT	AM	2	2	Quartzite (90%), QBS (10%). Chlorite alteration, quartz veins.	800261963	-1	15
BV15-39	7.62	9.14	Grey	QZT	AM	2	2	Very limited amount of sample. Quartz veins.	800261964	-1	14
BV15-39	9.14	10.67	Grey	QZT	AM	2	2	Quartzite. Minor chlorite alteration. Minor pyrite.	800261965	10	21
BV15-39	10.67	12.19	Grey	QZT	AM	2	2	Quartzite. Minor QBS. Quartz veins. Minor chlorite (?). Ox.	800261966	6	12
BV15-39	12.19	13.72	Grey	QZT	AM	1	1	Very little amount of sample. Greenish to brownish grey.	800261967	-1	15
BV15-39	13.72	15.24	Grey	QZT	AM	1	1	Brownish grey, quartz veins. Pyrite.	800261968	-1	9
BV15-39	15.24	16.76	Grey	QZT	AM	1	1	Brownish grey, quartz veins. Very minor pyrite.	800261969	-1	10
BV15-39	16.76	18.29	Grey	QZT	AM	1	1		800261970	-1	8
BV15-39	18.29	19.81	Grey	QZT	AM	1	1	Brownish grey, quartz veins, ox.	800261971	-1	14
BV15-39	19.81	21.34	Grey	QZT	AM	1	1	Brownish grey, quartz veins, ox.	800261972	-1	14
BV15-39	21.34	22.86	Grey	QZT	AM	1	1	Brownish grey, quartz veins, ox.	800261973	-1	13
BV15-39	22.86	24.38	Grey	QZT	AM	2	0.5	Brownish grey, quartz veins, ox.	800261974	12	32
BV15-39	24.38	25.91	Grey	QZT	AM	1	0.5	Brownish grey, quartz veins, ox.	800261975	10	32
BV15-39	25.91	27.43	Grey	QZT	AM	1	0.5	Brownish grey, quartz veins, ox.	800261976	8	39
BV15-39	27.43	28.96	Grey	QZT	AM	1	1	Brownish grey, quartz veins, ox.	800261977	5	37
BV15-39	28.96	30.48	Grey	QZT	AM	1	1	Brownish grey, minor pyrite	800261978	-1	19
BV15-39	30.48	32	Grey	QZT	AM	1	2	Quartzite, greenish to brownish grey, minor pyrite	800261979	-1	29
BV15-39	32	33.53	Grey	QZT	AM	1	2	Quartzite, greenish to brownish grey, minor pyrite	800261981	7	20
BV15-39	33.53	35.05	Grey	QZT	AM	1	2	Quartzite, greenish to brownish grey, minor pyrite	800261982	-1	7
BV15-39	35.05	36.58	Grey	QBS	AM	1	2	QBS 60%, Quartzite 40%, chlorite alteration, minor pyrite	800261983	-1	11
BV15-39	36.58	38.1	Grey	QZT	AM	1	1	Quartzite, greenish grey	800261984	-1	23
BV15-39	38.1	39.62	Grey	QZT	AM	1	1	Quartzite, greenish grey, minor pyrite	800261985	9	18
BV15-39	39.62	41.15	Grey	QBS	AM	1	3	Dark greenish grey	800261986	8	10
BV15-39	41.15	42.67	Grey	QBS	AM	1	3	Light grey. Minor silicification (?), pyrite with quartz (?)	800261987	-1	6
BV15-39	42.67	44.2	Grey	QBS	AM	2	1	Greenish grey, chlorite alteration	800261988	-1	11
BV15-39	44.2	45.72	Grey	QBS	AM	1	2	Greenish grey, chlorite alteration, pyrite disseminated, minor quartzite (?)	800261989	11	11
BV15-39	45.72	47.24	Grey	QBS	AM	1	1	Greenish grey, chlorite alteration, pyrite disseminated, minor quartzite (?). Very little sample, problem with hole.	800261991	-1	10
BV15-39	47.24	48.77	Grey	QBS	AM	1	1	Greenish grey, chlorite alteration, pyrite disseminated, minor quartzite (?). Very little sample, problem with hole.	800261992	6	23
BV15-39	48.77	50.29	Grey	QBS	AM	2	1	Minor chlorite alteration. Quartz bands. Moderate oxidation.	800261993	18	25
BV15-39	50.29	51.82	Grey	QBS	AM	1	1	Minor chlorite alteration. Quartz bands.	800261994	-1	13
BV15-39	51.82	53.34	Grey	QBS	AM	1	2	Minor chlorite alteration. Quartz bands.	800261995	17	19
BV15-39	53.34	54.86	Grey	QBS	AM	1	1	Minor chlorite alteration. Quartz bands.	800261996	21	27
BV15-39	54.86	56.39	Grey	QBS	AM	1	1	Minor chlorite alteration. Quartz bands.	800261997	12	11
BV15-39	56.39	57.91	Grey	QBS	AM	1	1	Minor chlorite alteration. Minor quartz veins.	800261998	20	17
BV15-39	57.91	59.44	Grey	QBS	AM	1	0.5	Quartz bands, minor chlorite alteration. Quartz veins.	800261999	47	91
BV15-39	59.44	60.96	Grey	QZT	AM	1	2	Quartzite 80%, QBS 20%. Quartz veins. Minor oxidation.	800262000	184	156
BV15-39	60.96	62.48	Grey	QBS	AM	1	1	Quartz bands in QBS; minor chlorite	800312001	121	50
BV15-39	62.48	64.01	Grey	QBS	AM	1	2	Quartz bands in QBS; minor chlorite	800312002	59	53
BV15-39	64.01	65.53	Grey	QZT	AM	1	1	Greenish grey to light green minor oxidation. Minor pyrite.	800312003	28	78
BV15-39	65.53	67.06	Grey	QBS	AM	1	1	Dark greenish grey	800312004	32	106
BV15-39	67.06	68.58	Grey	QZT	AM	1	1	Greenish grey. Mild oxidation.	800312005	78	168
BV15-39	68.58	70.1	Grey	QZT	AM	1	4	Grey colour to brownish. Quartz veins. Minor pyrite.	800312006	96	344
BV15-39	70.1	71.63	Grey	QBS	AM	1	1	Quartz veins in QBS. Pyrite	800312007	88	289
BV15-39	71.63	73.15	Grey	QBS	AM	1	2	QBS with quartz bands, quartz veins, minor pyrite	800312008	81	100
BV15-39	73.15	74.68	Grey	QZT	AM	1	2	Quartzite with quartz veins. Minor oxidation. 30% QBS	800312009	234	136
BV15-39	74.68	76.2	Grey	QZT	AM	1	3	Quartzite with quartz veins. Minor oxidation. 10% QBS	800312010	733	1540
BV15-39	76.2	77.72	Grey	QZT	AM	2	5	Quartzite with quartz veins. Minor oxidation. 10% QBS	800312011	896	1970
BV15-39	77.72	79.25	Grey	QZT	AM	1	5	Light grey. Cubic pyrite.	800312012	539	756
BV15-39	79.25	80.77	Grey	QZT	AM	1	5	Light grey. Cubic pyrite.	800312013	345	398
BV15-39	80.77	82.3	Grey	QZT	AM	1	3	20% QBS	800312014	207	443
BV15-39	82.3	83.82	Grey	QZT	AM	0	3	20% QBS	800312015	167	378
BV15-39	83.82	85.34	Grey	QZT	AM	0	3	Light grey. 25% QBS	800312016	219	412
BV15-39	85.34	86.87	Grey	QBS	AM	0	2	QBS. Chlorite alteration. Quartz vein with pyrite.	800312017	60	63
BV15-39	86.87	88.39	Grey	QBS	AM	0	4	QBS. Chlorite alteration. Quartz vein with pyrite.	800312018	81	77
BV15-39	88.39	89.92	Grey	QBS	AM	1	2	QBS. Chlorite alteration. Quartz vein with pyrite.	800312019	39	48
BV15-39	89.92	91.44	Grey	QBS	AM	0	0.5	QBS, dark greenish grey, minor quartz and pyrite veins	800312021	60	88

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-39	91.44	92.96	Grey	QBS	AM	0	0.5	Dark greenish grey, minor quartz veins	800312022	58	85
BV15-39	92.96	94.49	Grey	QBS	AM	1	2	Dark greenish grey, quartz veins with pyrite. 20% quartzite.	800312023	53	82
BV15-39	94.49	96.01	Grey	QBS	AM	1	1	Dark greenish grey, quartz veins with pyrite. 3% quartzite.	800312024	29	28
BV15-39	96.01	97.54	Grey	QBS	AM	1	1	Dark greenish grey, quartz veins with pyrite. 4% quartzite.	800312025	40	42
BV15-39	97.54	99.06	Grey	QZT	AM	1	3	Light grey quartz vein with disseminated pyrite	800312026	467	1000
BV15-39	99.06	100.58	Grey	QBS	AM	1	2	Dark grey vein or silicified rock with fine grained pyrite. 10% quartzite.	800312027	52	127
BV15-39	100.58	102.11	Grey	QBS	AM	1	3	Dark grey, QBS with light grey quartz and pyrite veins, greenish alteration (chlorite/sericite?), minor oxidation.	800312028	60	139
BV15-39	102.11	103.63	Grey	QBS	AM	0	3		800312029	32	61
BV15-39	103.63	105.16	Grey	QBS	DG	0	5	More vein material in the bag/bucket than chip tray. Appeared to see stratification in the bucket of layers of white chips ~3-5cm thick, almost like veins from the rock. These layers were ~70-80% quartz and I observed ~4 of them in the 5 gallon bucket. 10% quartzite. **Driller did a rod cap count and he thinks the hole should be 5ft deeper so a 355'. I will not change anything but an error was observed here.	800312031	25	29
BV15-39	105.16	106.68	Grey	QBS	DG	1	1	Quartz vein % dropped right off compared to previous sample. Pyrite occurs in quartz vein and schist.	800312032	22	52
BV15-39	106.68	108.2	Grey	QBS	DG	2	3	Dominated by schist but 10-20% of interval comprised of quartzite.	800312033	29	39
BV15-39	108.2	109.73	Grey	QBS	DG	0	3	Pyrite in vein and schist. **Driller said footage could be right if the 5' above ground is taken into consideration. The top of hole assemble is 5 ft above ground and driller contend that in his estimate.	800312034	17	52
BV15-39	109.73	111.25	Grey	QBS	DG	1	2	~10-20% quartzite	800312035	28	64
BV15-39	111.25	112.78	Grey	QZT	DG	1	3	30% QBS. Pyrite occurs in vein and host rock. Rare biotite layers - sub mm define banding with quartzite.	800312036	58	84
BV15-39	112.78	114.3	Grey	QBS	DG	1	2	10-20% quartzite. Very minor oxidation of quartz fragments.	800312037	35	45
BV15-39	114.3	115.82	Grey	QBS	DG	0	5	30% quartzite. Veins are white to grey.	800312038	54	67
BV15-39	115.82	117.35	Grey	QZT	DG	0	2	Minor chips of QBS. Veins drop dramatically. Rocks are non-magnetic.	800312039	40	38
BV15-39	117.35	118.87	Grey	QBS	DG	0	1	Single chip appears to show a quartz vein cutting through wall rock - subparallel to foliation but foliation is not well developed in rock.	800312040	42	29
BV15-39	118.87	120.4	Grey	QZT	DG	0	3	Quartz veins are white, irregularly milky, rarely grey. Pyrite occurs both in veins in wall rock. Pyrite is typically fine grained and forming wispy patches.	800312041	34	32
BV15-39	120.4	121.92	Grey	QZT	DG	0	3	30% QBS	800312042	44	50
BV15-39	121.92	123.44	Grey	QBS	DG	0	10	~20-40% quartzite. White to smokey grey quartz veins. Pyrite occurs in QBS and vein material. Observed stratification in the bucket ~10cm seen of salt and pepper/white reflecting increasing concentration of quartz. I think the bucket is reflecting the actual quartz in the ground!? No idea how thick it actually could be, 20-30 cm?	800312043	134	68
BV15-39	123.44	124.97	Grey	QBS	DG	0	2	Veins are milky white to grey. Sulphides mainly in the schist.	800312044	34	23
BV15-39	124.97	126.49	Grey	QBS	DG	0	4	30% quartzite. Pyrite in schist and quartz veins.	800312045	45	31
BV15-39	126.49	128.02	Grey	QBS	DG	0	2	20-30% quartzite. Veins (not all) are adopting a subtle pink colour. Not sure if it could be a potassic alteration.	800312046	26	11
BV15-39	128.02	129.54	Grey	QZT	DG	0	3	Quartzite is irregularly coloured pink and green in the proximity to veins and within the quartz veins. Could reflect chlorite or potassic but only weak alteration if the case. Pyrite observed in vein material.	800312047	77	15
BV15-39	129.54	131.06	Grey	QZT	DG	0	3	Sulphides occur both in vein and wall rock.	800312048	67	20
BV15-39	131.06	132.59	Grey	QZT	DG	0	4	Veins are both milky white and smokey grey. Sulphides occur in veins and wall rock. Very different to discern silicification. Looks like unaltered quartzite. Another "seam" of quartz was observable in the bucket. Looks to be ~4-6cm thick measured within the bucket.	800312049	42	24
BV15-39	132.59	134.11	Grey	QZT	DG	0	3	Vein are dominantly smokey grey in colour.	800312051	23	24
BV15-39	134.11	135.64	Grey	QBS	DG	0	2	20-30% quartzite. Pyrite decreasing.	800312052	20	17
BV15-39	135.64	137.16	Grey	QZT	DG	0	2		800312053	34	22
BV15-39	137.16	138.68	Grey	QZT	DG	2	3	Big difference is the oxidation level has increased noticeably. Rock got harder and the return was white. QBS and biotite for that matter is essentially absent. Possibly increase in oxidation represents a fault structure.	800312054	42	40
BV15-39	138.68	140.21	Grey	QZT	DG	2	2	Oxidation still moderate but not quite as intense as above. Slight increase in QBS chips to 10-15% of interval.	800312055	38	37
BV15-39	140.21	141.73	Grey	QZT	DG	1	1	Pyrite increasing but primarily in the quartzite and not in the veins. Veins are white to grey.	800312056	23	27
BV15-39	141.73	143.26	Grey	QZT	DG	1	3	Pyrite in veins and within quartzite.	800312057	45	31
BV15-39	143.26	144.78	Grey	QZT	DG	1	1	Single chip observed a 3-4mm wide vein within quartzite and vein is parallel to banding in quartzite.	800312058	34	55
BV15-39	144.78	146.3	Grey	QZT	DG	0	3	Subtle brown colour of some quartzite grains due to minor biotite content. Pyrite occurs both within quartz veins and in quartzite wall rock.	800312059	43	96
BV15-39	146.3	147.83	Grey	QZT	DG	0	1	Pyrite mostly occurs within the quartzite. Pyrite rare to absent within the veins.	800312061	27	30
BV15-39	147.83	149.35	Grey	QZT	DG	0	1	Biotite content increasing in most chips. Still ~20% quartzite chips. Quartz veins look fairly "dry" with no sulphides and no alteration.	800312062	26	11
BV15-39	149.35	150.88	Grey	QZT	DG	0	2	Pyrite content decreasing. No pyrite in veins. Veins are typically white.	800312063	38	17
BV15-39	150.88	152.4	Grey	QZT	DG	0	2	Only rare pyrite with quartz veins. Pyrite is rarely disseminated and also occurs along fractures within quartzite - i.e. Coating flat surfaces.	800312064	48	157
BV15-39	152.4	153.92	Grey	QZT	DG	0	5	Considered shutting hole down but veins increasing and pyrite remained consistent. Pyrite occurs in both veins and wallrock.	800312065	69	20
BV15-39	153.92	155.45	Grey	QBS	DG	0	2	20-30% quartzite.	800312066	53	17
BV15-39	155.45	156.97	Grey	QBS	DG	0	2	10-20% Quartzite	800312067	48	22

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-40	0	1.52		CAS					No Sample		
BV15-40	1.52	3.05	Grey	QBS	AM	1	5	Greenish grey. QBS >>>>Quartzite (10%). Quartz veins. Oxidation. Chlorite alteration.	800312068	16	40
BV15-40	3.05	4.57	Grey	QBS	AM	1	5	Greenish grey. QBS >>>>Quartzite (10%). Quartz veins. Oxidation. Chlorite alteration.	800312069	-1	25
BV15-40	4.57	6.1	Grey	QBS	AM	1	3	Greenish grey. QBS>Quartzite (30%). Quartz veins, oxidation	800312070	-1	24
BV15-40	6.1	7.62	Grey	QBS	AM	1	1	Greenish grey, minor quartz veins, chlorite alteration. Water at 24'.	800312071	5	66
BV15-40	7.62	9.14	Grey	QBS	AM	1	2	QBS>>Quartzite (20%). Greenish grey, quartzite, oxidation.	800312072	-1	58
BV15-40	9.14	10.67	Grey	QBS	AM	2	2	QBS>>Quartzite (40%). Greenish grey to brownish. Quartz veins, oxidation.	800312073	-1	52
BV15-40	10.67	12.19	Grey	QZT	AM	2	5	Quartzite (80%), QBS (20%). Brownish grey quartz veins.	800312074	52	130
BV15-40	12.19	13.72	Grey	QZT	AM	1	5	Quartzite (80%), QBS (20%). Brownish grey quartz veins.	800312075	314	158
BV15-40	13.72	15.24	Grey	QZT	AM	2	5	Brownish grey quartz veins.	800312076	467	261
BV15-40	15.24	16.76	Grey	QZT	AM	2	5	Brownish grey quartz veins.	800312077	346	591
BV15-40	16.76	18.29	Grey	QZT	AM	2	5		800312078	509	755
BV15-40	18.29	19.81	Grey	QZT	AM	2	5	Quartzite with quartz veins, moderate oxidation. Iron oxide with quartz. End of casing;	800312079	423	549
BV15-40	19.81	21.34	Grey	QZT	AM	1	5	Quartzite with quartz veins. Minor QBS with chlorite alteration.	800312081	188	253
BV15-40	21.34	22.86	Grey	QZT	AM	1	5	Brownish grey and white, minor QBS (10%) quartz veins. Possible void between 73 and 75'.	800312082	23	104
BV15-40	22.86	24.38	Grey	QZT	AM	1	5	Brownish grey to white, minor QBS (10%) Quartz and iron oxide veins. Rock quality is low, they will put more casing.	800312083	346	365
BV15-40	24.38	25.91		QZT	DP	2	0	Very small sample	800312084	27900	92
BV15-40	25.91	27.43	Grey	QBS	DP	1	0.5	Medium greenish grey, fine grained	800312085	2100	30
BV15-40	27.43	28.96	Grey	QBS	DP	1	0.5	Medium greenish grey, fine grained. Biotite rich brown bands to 3mm wide	800312086	253	25
BV15-40	28.96	30.48		QBS	DP	1	1		800312087	413	21
BV15-40	30.48	32		QBS	DP	1	1		800312088	13	23
BV15-40	32	33.53		QBS	DP	1	1		800312089	16	22
BV15-40	33.53	35.05		QBS	DP	1	0.5		800312091	15	27
BV15-40	35.05	36.58		QBS	DP	1	4	quartz veinlets to 5mm+ wide.	800312092	-1	24
BV15-40	36.58	38.1		QBS	DP	1	10	quartz veinlets to 5mm+ wide; small soupy sample	800312093	149	32
BV15-40	38.1	39.62		QBS	DP	2	30	30% vein quartz, often pervasive lmnt stained. "Vein" may be mostly quartzite.	800312094	143	27
BV15-40	39.62	41.15	Brown	QZT	DP	2	0	10% QBS; one subhedral grain pyrite seen. Light yellow-brown; pervasive limonite. Wet sample.	800312095	-1	41
BV15-40	41.15	42.67		QZT	DP	2	0	30% QBS; soupy messy sample	800312096	6	70
BV15-40	42.67	44.2		QBS	DP	1	0	30% Quartzite; soupy messy sample	800312097	-1	59
BV15-40	44.2	45.72		QBS	DP	1	0	30% Quartzite; soupy messy sample	800312098	-1	45
BV15-40	45.72	47.24		QBS	DP	1	0	20% Quartzite; soupy messy sample	800312099	7	54
BV15-40	47.24	48.77		QBS	DP	2	0.5	10% quartzite	800312100	-1	87
BV15-40	48.77	50.29		QBS	DP	2	0.2	20% Quartzite	800312101	-1	46
BV15-40	50.29	51.82		QBS	DP	1	0.2	10% indurated quartzite; pyrite blebs	800312102	-1	20
BV15-40	51.82	53.34		QBS	DP	2	0.2	10% indurated quartzite; pyrite blebs	800312103	-1	56
BV15-40	53.34	54.86	Grey	QBS	DP	2	0.2	Medium greenish grey; pyrite as blebs to 1.5mm and as subhedral grains	800312104	-1	61
BV15-40	54.86	56.39	Grey	QBS	DP	2	0.5	5% quartzite	800312105	-1	26
BV15-40	56.39	57.91	Grey	QBS	DP	2	1	Medium greenish grey	800312106	-1	24
BV15-40	57.91	59.44		QZT	DP	1	0.5	30% QBS	800312107	6	15
BV15-40	59.44	60.96		QBS	DP	1	1	not as muddy samples as above	800312108	-1	14
BV15-40	60.96	62.48		QBS	DP	1	0.5	not as muddy samples as above	800312109	-1	9
BV15-40	62.48	64.01		QBS	DP	1	1	Weak epidote altered. Pyrite cubes to 2.5mm.	800312110	11	13
BV15-40	64.01	65.53		QBS	DP	1	2	Pyrite crystals mainly biotite altered quartz veinlet selvages.	800312111	-1	12
BV15-40	65.53	67.06		QZT	DP	1	3	35% QBS. Locally weak epidote altered.	800312112	-1	7
BV15-40	67.06	68.58		QZT	DP	1	2	25% QBS. Locally weak epidote altered.	800312113	-1	7
BV15-40	68.58	70.1		QZT	DP	1	2	15% QBS. Locally weak epidote altered.	800312114	-1	9
BV15-40	70.1	71.63		QZT	DP	1	4	20% QBS. More epidote alteration than above.	800312115	-1	10
BV15-40	71.63	73.15		QZT	DP	1	2	10-15% biotite throughout pyrite as blebs to 3mm across.	800312116	-1	8
BV15-40	73.15	74.68		QBS	DP	1	2	Banded pyrite grain 4mm. Weakly epidote altered.	800312117	5	10
BV15-40	74.68	76.2	Brown	QBS	DP	1	2	Very weakly epidote altered. Dark brown.	800312118	-1	5
BV15-40	76.2	77.72		QBS	DP	1	2	Pyrite blebs mainly on margin quartz veinlets	800312119	-1	8
BV15-40	77.72	79.25		QBS	DP	1	2		800312121	-1	9
BV15-40	79.25	80.77	Brown	QBS	DP	1	1	Dark greyish brown; 15% light grey quartzite.	800312122	-1	12
BV15-40	80.77	82.3	Grey	QBS	DP	1	2	Dark greenish grey. Very fine grained blebby pyrite.	800312123	-1	10
BV15-40	82.3	83.82	Grey	QBS	DP	0	1	Dark greenish grey. Very fine grained blebby pyrite.	800312124	-1	7
BV15-40	83.82	85.34	Grey	QBS	DP	1	1	Dark brown grey. Very fine grained blebby pyrite. 15% quartzite which contains much more pyrite than QBS.	800312125	-1	12
BV15-40	85.34	86.87		QZT	DP	1	1	30% QBS at top of interval.	800312126	100	84
BV15-40	86.87	88.39		QZT	DP	0	1	3% QBS chips. Bright green mineral (chlorite? Malachite?) seen in a few places.	800312127	505	1820
BV15-40	88.39	89.92	Brown	QZT	DP	0	1	35% medium - dark brown QBS. More bright green mineral traces seen.	800312128	387	1010

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-40	89.92	91.44		QBS	DP	0	2	30% interval well mineralized, indurated quartzite. Subhedral dogtooth calcite crystals in vein along fracture in QBS.	800312129	235	452
BV15-40	91.44	92.96		QZT	DP	0	1	40% QBS	800312131	156	345
BV15-40	92.96	94.49		QBS	DP	0	1	10% quartzite	800312132	54	153
BV15-40	94.49	96.01		QZT	DP	1	1	35% QBS	800312133	111	757
BV15-40	96.01	97.54		QZT	DP	1	1	10% QBS	800312134	47	95
BV15-40	97.54	99.06		QBS	DP	0	0.5		800312135	47	433
BV15-40	99.06	100.58		QZT	DP	0	1	25% QBS; weak epidote alteration	800312136	23	34
BV15-40	100.58	102.11		QZT	DP	0	1	40% QBS	800312137	97	555
BV15-40	102.11	103.63		QZT	DP	1	1	Fault (?) at 337' where ground is badly broken. 30% QBS. Handful pebble sized chips collected. Good looking rock.	800312138	134	630
BV15-40	103.63	105.16		QZT	DP	1	2	10% QBS; another fault with pebbles. 345' pulling rods.	800312139	48	449
BV15-41	0	9.14	Grey	OB	DP	2	5	Overburden: QBS frags, medium grey, fine grained. Clear grey vein quartz chips. No sulphides seen.	800312140	118	39
BV15-41	9.14	10.67	Grey	QBS	DP	1	3	Overburden: QBS frags, medium grey, fine grained. Clear grey vein quartz chips. No sulphides seen.	800312141	9	18
BV15-41	10.67	12.19	Grey	QBS	DP	1	0.5	Medium Grey.	800312142	6	9
BV15-41	12.19	13.72	Grey	QBS	DP	1	0.5	20% clean, finely granular, watery grey quartzite.	800312143	11	19
BV15-41	13.72	15.24		QBS	DP	0	0	Tiny sample so did not extract any chips.	800312144	8	18
BV15-41	15.24	16.76	Grey	QBS	DP	1	0.5	40% quartzite	800312145	-1	25
BV15-41	16.76	18.29	Grey	QBS	DP	0	0.5	58' into water. A new record. 35% quartzite.	800312146	6	16
BV15-41	18.29	19.81	Grey	QBS	DP	0	0.3	20% quartzite.	800312147	10	12
BV15-41	19.81	21.34	Grey	QBS	DP	0	0.2	35% quartzite which contains 2% disseminated pyrite as subhedral crystals	800312148	7	9
BV15-41	21.34	22.86	Grey	QZT	DP	0	0.2	20% QBS; light grey.	800312149	7	6
BV15-41	22.86	24.38	Grey	QZT	DP	1	0	10% QBS; light grey.	800312151	-1	9
BV15-41	24.38	25.91	Grey	QZT	DP	1	0	10% QBS; light grey; vitreous.	800312152	8	8
BV15-41	25.91	27.43	Grey	QZT	DP	1	0	5% QBS; light grey; vitreous.	800312153	20	13
BV15-41	27.43	28.96	Grey	QZT	DP	1	0	3% QBS; light grey; vitreous.	800312154	-1	4
BV15-41	28.96	30.48	Grey	QZT	DP	1	0	3% QBS; light grey; vitreous.	800312155	-1	5
BV15-41	30.48	32	Grey	QZT	DP	1	0	10% QBS; light grey; vitreous.	800312156	9	4
BV15-41	32	33.53	Grey	QBS	DP	0	0	10% quartzite; QBS fresh.	800312157	11	5
BV15-41	33.53	35.05	Grey	QZT	DP	1	0	15% QBS; light grey, vitreous	800312158	-1	8
BV15-41	35.05	36.58	Grey	QZT	DP	1	0.2	10% QBS; pale grey, vitreous	800312159	5	10
BV15-41	36.58	38.1	Grey	QZT	DP	0	0.2	10% QBS; pale grey, vitreous	800312161	11	4
BV15-41	38.1	39.62	Grey	QZT	DP	0	0	10% QBS; pale grey, vitreous	800312162	8	-1
BV15-41	39.62	41.15		QBS	DP	0	0.2	35% quartzite	800312163	47	3
BV15-41	41.15	42.67		QZT	DP	0	0.2	15% QBS	800312164	5	7
BV15-41	42.67	44.2		QZT	DP	0	0	15% QBS	800312165	-1	5
BV15-41	44.2	45.72		QZT	DP	0	0.2	10% QBS	800312166	8	9
BV15-41	45.72	47.24		QZT	DP	0	0.3		800312167	7	7
BV15-41	47.24	48.77		QZT	DP	0	0.1		800312168	-1	4
BV15-41	48.77	50.29		QZT	DP	0	0.2	Badly broken ground; likely fault. Pyrite as very finely disseminated specks, streaky. Good looking rock.	800312169	-1	6
BV15-42	0	3.05		OB					No Sample		
BV15-42	3.05	4.57	Grey	QBS	DP	1	0	20% quartzite; QBS medium greenish grey. Finely banded; weakly epidote altered.	800312170	2100	6
BV15-42	4.57	6.1		QBS	DP	2	0.2	35% quartzite; quartzite oxidized.	800312171	22	11
BV15-42	6.1	7.62		QBS	DP	1	0.3	10% quartzite; quartzite oxidized.	800312172	-1	5
BV15-42	7.62	9.14		QBS	DP	2	0.3	10% quartzite; quartzite oxidized.	800312173	182	9
BV15-42	9.14	10.67		QZT	DP	1	0.3	40% quartzite; quartzite oxidized.	800312174	-1	6
BV15-42	10.67	12.19		QBS	DP	1	0.2	20% quartzite; quartzite oxidized.	800312175	-1	6
BV15-42	12.19	13.72	Grey	QBS	DP	0	0.2	Dark grey, fresh. 5% quartzite.	800312176	-1	7
BV15-42	13.72	15.24	Grey	QBS	DP	1	0.2	Dark grey, fresh. 35% quartzite.	800312177	-1	7
BV15-42	15.24	16.76	Grey	QBS	DP	1	0.2	Dark grey, fresh. 20% quartzite.	800312178	-1	10
BV15-42	16.76	18.29	Grey	QBS	DP	1	0.2	Dark grey, fresh. 5% quartzite.	800312179	-1	11
BV15-42	18.29	19.81	Grey	QBS	DP	1	1	Dark grey, fresh. 10% quartzite.	800312181	-1	7
BV15-42	19.81	21.34	Grey	QBS	DP	0	0.2	Dark grey, fresh. 5% quartzite.	800312182	-1	7
BV15-42	21.34	22.86	Grey	QBS	DP	1	1	Dark grey, fresh. 5% quartzite.	800312183	-1	10
BV15-42	22.86	24.38	Grey	QBS	DP	0	1	Dark grey, fresh. 5% quartzite.	800312184	-1	8
BV15-42	24.38	25.91	Grey	QBS	DP	0	1	Dark grey, fresh. 5% quartzite.	800312185	10	9
BV15-42	25.91	27.43	Grey	QBS	DP	0	2	Dark grey, fresh. 5% quartzite.	800312186	-1	10
BV15-42	27.43	28.96	Grey	QBS	DP	0	1	Dark grey, fresh. 5% quartzite.	800312187	-1	9
BV15-42	28.96	30.48	Grey	QBS	DP	0	1	Dark grey, fresh. 5% quartzite.	800312188	-1	10
BV15-42	30.48	32	Grey	QBS	DP	0	2	Dark grey, fresh. 5% quartzite.	800312189	-1	10
BV15-42	32	33.53	Grey	QBS	DP	1	2	5% Quartzite. Greenish grey	800312191	-1	7

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-42	33.53	35.05	Grey	QBS	DP	0	0.2	Medium greenish grey, finely banded. Small, soupy wet sample. Pyrite very fine disseminated mostly along quartz veinlet margins.	800312192	-1	11
BV15-42	35.05	36.58	Grey	QBS	DP	1	0.4	As above except here weakly oxidized	800312193	-1	9
BV15-42	36.58	38.1		QBS	DP	1	0.5	Fresher; as above except here weakly oxidized.	800312194	-1	8
BV15-42	38.1	39.62	Grey	QBS	DP	0	0.7	Medium greenish grey; pyrite very fine disseminated	800312195	-1	9
BV15-42	39.62	41.15	Grey	QBS	DP	0	1	Medium greenish grey; pyrite very fine disseminated	800312196	-1	5
BV15-42	41.15	42.67	Grey	QBS	DP	0	1	Medium greenish grey; pyrite very fine disseminated	800312197	-1	8
BV15-42	42.67	44.2	Grey	QBS	DP	0	2	Medium greenish grey; pyrite very fine disseminated	800312198	-1	9
BV15-42	44.2	45.72	Grey	QBS	DP	0	1	Medium greenish grey; pyrite very fine disseminated; 15% light grey fine grained quartzite	800312199	-1	7
BV15-42	45.72	47.24	Grey	QBS	DP	0	0.5	10% quartzite as above.	800312200	-1	6
BV15-42	47.24	48.77		QZT	DP	0	0.3	5% QBS	800312201	-1	5
BV15-42	48.77	50.29		QZT	DP	0	0.3	5% QBS	800312202	-1	6
BV15-42	50.29	51.82	Grey	QZT	DP	0	0.5	Quartzite minor QBS (10%)	800312203	-1	5
BV15-42	51.82	53.34	Grey	QBS	AM	0	0.5	QBS minor quartzite (20%), chlorite alteration, pyrite.	800312204	-1	7
BV15-42	53.34	54.86	Grey	QBS	AM	0	0.5	QBS minor quartzite (20%), chlorite alteration, pyrite.	800312205	-1	5
BV15-42	54.86	56.39	Grey	QBS	AM	0	0.5	QBS minor quartzite (20%), chlorite alteration, pyrite.	800312206	-1	3
BV15-42	56.39	57.91	Grey	QZT	AM	0	0.5	Quartzite minor QBS (30%), chlorite alteration, epidote alteration.	800312207	-1	4
BV15-42	57.91	59.44	Grey	QZT	AM	0	1	Quartzite minor QBS (20%), quartz veins, chlorite alteration, epidote	800312208	-1	4
BV15-42	59.44	60.96	Grey	QZT	AM	0	1	Quartzite minor QBS (20%), quartz veins, chlorite alteration, epidote	800312209	-1	-1
BV15-42	60.96	62.48	Grey	QBS	AM	0	1	Quartzite (20%), Quartz veins, chlorite alteration, epidote alteration. Minor pyrite, weakly magnetic	800312210	-1	4
BV15-42	62.48	64.01	Grey	QZT	AM	0	1	Quartzite minor QBS (10%). Chlorite alteration, minor epidote (?). Quartz veins and pyrite in veins and biotite.	800312211	-1	4
BV15-42	64.01	65.53	Grey	QZT	AM	0	1	Quartzite minor QBS (10%). Chlorite alteration, minor epidote (?). Quartz veins and pyrite in veins and biotite.	800312212	-1	3
BV15-42	65.53	67.06	Grey	QBS	AM	0	1	Quartzite (20%), chlorite alteration, minor epidote (?), quartz veins + po + py	800312213	-1	4
BV15-42	67.06	68.58	Grey	QZT	AM	0	1	Quartzite (20%), chlorite alteration, minor epidote (?), quartz veins + po + py. 20% QBS.	800312214	-1	6
BV15-42	68.58	70.1	Grey	QZT	AM	0	1	Quartzite (20%), chlorite alteration, minor epidote (?), quartz veins + po + py. 20% QBS.	800312215	-1	7
BV15-42	70.1	71.63	Grey	QZT	AM	0	1	Quartzite (20%), chlorite alteration, minor epidote (?), quartz veins + po + py. 10% QBS.	800312216	-1	-1
BV15-42	71.63	73.15	Grey	QZT	AM	0	0.5	Quartzite, quartz veins, minor pyrite, chlorite, epidote alteration	800312217	-1	-1
BV15-42	73.15	74.68	Grey	QZT	AM	0	1	Quartzite, quartz veins, minor pyrite, chlorite, epidote alteration, 10% QBS	800312218	-1	4
BV15-42	74.68	76.2	Grey	QZT	AM	0	1	Quartzite, quartz veins, minor pyrite, chlorite, epidote alteration, 5% QBS	800312219	-1	3
BV15-42	76.2	77.72	Grey	QZT	AM	0	3	Quartzite, chlorite alteration, minor QBS, Quartz + pyrite veins	800312221	-1	4
BV15-42	77.72	79.25	Grey	QBS	AM	0	0.5	QBS, minor quartzite, chlorite alteration, minor pyrite.	800312222	-1	5
BV15-42	79.25	80.77	Grey	QBS	AM	0	0.5	QBS, minor quartzite, chlorite alteration, minor pyrite.	800312223	-1	4
BV15-42	80.77	82.3	Grey	QBS	AM	0	0.5	QBS, minor quartzite, chlorite alteration, minor pyrite.	800312224	-1	5
BV15-42	82.3	83.82	Grey	QBS	AM	0	0.5	QBS, minor quartzite, chlorite alteration, minor pyrite.	800312225	-1	6
BV15-42	83.82	85.34	Grey	QBS	AM	0	1	QBS minor quartzite (10%). Chlorite alteration, minor pyrite. Weakly magnetic.	800312226	-1	4
BV15-42	85.34	86.87	Grey	QBS	AM	0	1	QBS minor quartzite (10%). Chlorite alteration, minor pyrite. Weakly magnetic.	800312227	-1	4
BV15-42	86.87	88.39	Grey	QBS	AM	0	0.5	QBS minor quartzite (10%). Chlorite alteration, minor pyrite. Weakly magnetic.	800312228	-1	7
BV15-42	88.39	89.92	Grey	QBS	AM	0	0.5	QBS minor quartzite (10%). Chlorite alteration, minor pyrite. Weakly magnetic.	800312229	-1	6
BV15-42	89.92	91.44	Grey	QBS	AM	0	1	QBS 60%, Quartzite 40%. Quartz veins, chlorite alteration.	800312231	91	713
BV15-42	91.44	92.96	Grey	QZT	AM	0	80	Light grey quartz vein, minor QBS; indurated; pyrite as subhedral grains	800312232	1310	3350
BV15-42	92.96	94.49	Grey	QZT	AM	0	0	Light grey, silica, sericite (?), pyrite	800312233	2140	5330
BV15-42	94.49	96.01	Grey	QZT	AM	0	0	Light grey, silica, sericite (?), pyrite	800312234	1430	6050
BV15-42	96.01	97.54	Grey	QBS	AM	0	10	QBS with minor quartzite (10%), silica, sericite, sa, chlorite alteration. Quartz veins with pyrite.	800312235	99	226
BV15-42	97.54	99.06	Grey	QBS	AM	0	10	QBS, quartz veins, chlorite alteration	800312236	38	75
BV15-42	99.06	100.58	Grey	QBS	AM	0	1	Greenish grey, minor pyrite, chlorite, epidote; quartz veins weakly magnetic.	800312237	10	20
BV15-42	100.58	102.11	Grey	QBS	AM	0	0.5	Greenish grey, minor pyrite, chlorite, epidote; quartz veins weakly magnetic.	800312238	7	12
BV15-42	102.11	103.63	Grey	QBS	AM	0	1	QBS, minor quartz + pyrite veins, chlorite alteration, epidote (?).	800312239	10	18
BV15-42	103.63	105.16	Grey	QBS	AM	0	0.5	QBS, minor quartz + pyrite veins, chlorite alteration, epidote (?).	800312240	15	24
BV15-42	105.16	106.68	Grey	QBS	AM	0	1	QBS (70%), Quartzite (30%), quartz veins with pyrite.	800312241	8	14
BV15-42	106.68	108.2	Grey	QZT	AM	0	0	Quartzite 60%, QBS 40%. Quartz + pyrite veins. Weakly magnetic.	800312242	12	31
BV15-42	108.2	109.73	Grey	QZT	AM	0	0	QBS; chlorite-epidote alteration. Minor quartz+pyrite veins. 20% QBS.	800312243	-1	8
BV15-42	109.73	111.25	Grey	QBS	AM	0	0.7	QBS; chlorite-epidote alteration. Minor quartz+pyrite veins. 20% QBS.	800312244	14	12
BV15-42	111.25	112.78	Grey	QBS	AM	0	3	QBS, chlorite, quartz veins, minor pyrite. Quartzite 30%.	800312245	18	19
BV15-42	112.78	114.3	Grey	QBS	AM	0	2	QBS, chlorite, quartz veins, minor pyrite. Quartzite 30%.	800312246	16	17
BV15-42	114.3	115.82	Grey	QBS	AM	0	2	QBS>Quartzite. Chlorite alteration. Quartz veins, minor pyrite	800312247	13	14
BV15-42	115.82	117.35	Grey	QBS	AM	0	0	Greenish grey, chlorite alteration, qtz veins + pyrite	800312248	11	11
BV15-42	117.35	118.87	Grey	QBS	AM	0	0	Greenish grey, chlorite alteration, qtz veins + pyrite	800312249	13	12
BV15-42	118.87	120.4	Grey	QBS	AM	0	1	Greenish grey, chlorite alteration, qtz veins + pyrite	800312251	16	33
BV15-42	120.4	121.92	Grey	QBS	AM	0	1	Quartzite (30%). Chlorite alteration. Quartz + pyrite veins.	800312252	19	23

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
BV15-42	121.92	123.44	Grey	QBS	AM	0	1	QBS > Quartzite (10%). Chlorite alteration, quartz + pyrite veins.	B00312253	62	247
BV15-42	123.44	124.97	Grey	QZT	AM	0	10	Quartzite 60%, QBS 40%, quartz + pyrite veins	B00312254	75	108
BV15-42	124.97	126.49	Grey	QZT	AM	0	10	Quartzite 60%, QBS 40%, quartz + pyrite veins	B00312255	53	70
BV15-42	126.49	128.02	Grey	QBS	AM	0	1	Greenish grey, chlorite, epidote, minor quartz veins.	B00312256	10	17
BV15-42	128.02	129.54	Grey	QBS	AM	0	0.5	Greenish grey, chlorite, epidote, minor quartz veins.	B00312257	7	13
BV15-42	129.54	131.06	Grey	QBS	AM	0	0.5	QBS (70%) > Quartzite (30%), minor quartz veins, minor pyrite	B00312258	14	11
BV15-42	131.06	132.59	Grey	QBS	AM	0	0.5	Greenish grey, chlorite, epidote, quartz veins	B00312259	10	7
BV15-42	132.59	134.11	Grey	QBS	AM	0	0.5	QBS (70%), Quartzite (30%), quartz veins, chlorite, minor pyrite	B00312261	-1	10
BV15-42	134.11	135.64	Grey	QBS	AM	0	1	QBS (70%), Quartzite (30%), quartz veins, chlorite, minor pyrite	B00312262	11	12
BV15-42	135.64	137.16	Grey	QBS	AM	0	1	QBS (70%), Quartzite (30%), quartz veins, chlorite, minor pyrite	B00312263	11	12
BV15-42	137.16	138.68	Green	QBS	AM	0	2	QBS (60%), Quartzite (40%), chlorite, epidote and calcite occurrence (prophyllitic alteration) minor quartz and pyrite.	B00312264	13	10
BV15-42	138.68	140.21	Grey	QZT	AM	0	1	Quartzite (60%), QBS (40%). Chlorite alteration, disseminated pyrite, quartz and pyrite veins.	B00312265	10	12
BV15-42	140.21	141.73	Grey	QZT	AM	0	2	Quartzite (60%), QBS (40%), quartz and pyrite veins. Pyrite micro veins.	B00312266	30	17
BV15-42	141.73	143.26		QZT	DP	0	2	Quartzite (60%), QBS (25%), quartz and pyrite veins. Pyrite micro veins.	B00312267	14	13
BV15-42	143.26	144.78		QBS	DP	0	1	30% quartzite	B00312268	9	28
BV15-42	144.78	146.3		QBS	DP	0	3	Weakly epidote-altered. Pyrite blebs to 3.5mm within quartz veinlet material.	B00312269	141	168
BV15-42	146.3	147.83		QBS	DP	0	2	20% quartzite. Weak epidote alteration.	B00312270	-1	6
BV15-42	147.83	149.35	Grey	QBS	DP	0	2	Weakly epidote-altered. Dark greenish grey	B00312271	12	7
BV15-42	149.35	150.88	Grey	QBS	DP	0	2	Weakly epidote-altered. Dark greenish grey	B00312272	-1	9
BV15-42	150.88	152.4	Grey	QBS	DP	0	1	Dark greenish grey; 5% quartzite.	B00312273	9	9
BV15-42	152.4	153.92	Grey	QBS	DP	0	2	Dark greenish grey; 5% quartzite.	B00312274	-1	15
BV15-42	153.92	155.45	Grey	QBS	DP	0	2	Dark greenish grey; 5% quartzite.	B00312275	5	8
BV15-42	155.45	156.97	Grey	QBS	DP	0	2	Dark greenish grey; 5% quartzite.	B00312276	8	8
BV15-42	156.97	158.5	Grey	QBS	DP	0	2	Dark greenish grey; 5% quartzite.	B00312277	7	8
BV15-42	158.5	160.02	Grey	QBS	DP	0	1	Dark greenish grey; 5% quartzite.	B00312278	-1	8
BV15-42	160.02	161.54	Grey	QBS	DP	0	1	Dark greenish grey; 5% quartzite.	B00312279	-1	4
BV15-42	161.54	163.07	Grey	QBS	DP	0	3	Dark greenish grey; 5% quartzite.	B00312281	-1	7
BV15-42	163.07	164.59	Grey	QBS	DP	0	3	Dark greenish grey; 5% quartzite.	B00312282	-1	9
BV15-42	164.59	166.12	Grey	QBS	DP	0	1	Dark greenish grey; 5% quartzite.	B00312283	9	9
BV15-42	166.12	167.03	Grey	QBS	DP	0	1	Dark greenish grey; 5% quartzite; one subhedral pyrite grain 2.5mm across.	B00312284	10	14
TG15-01	0	1.52		CAS	DG	3	0	Mixture of QBS, CS, QMS. Inseting one more run of 5 foot casing. Moderate oxidation.	No Sample		
TG15-01	1.52	3.05		CAS	DG	3	0	Mixture of QBS, CS, QMS. Inseting one more run of 5 foot casing. Moderate oxidation.	E862286	9	38
TG15-01	3.05	4.57	Orange	QMS	DG	3	20	Qtz muscovite schist. Qtz veins comprise 20% of interval. Wall rock is commonly oxidized. Pyrite occurs with qtz veins. Possible trace silver coloured fine grained sx within single qtz vein grain. Moderate oxidation. 75% quartz, 2% pyrite mineralization.	E862287	10	10
TG15-01	4.57	6.1	Orange	QMS	DG	3	20	Qtz veins = 20% of rock. These veins are typically white while the schist is orange. Muscovite copmries ~5-10% of rock forming very thin septa between the 1-2mm thick quartz bands. Moderate oxidation.	E862288	-1	10
TG15-01	6.1	7.62	Orange	QMS	DG	3	20	Qtz veins comprise ~20% - some vein material oxidized, some no. No sx. Moderate oxidation, 75% quartz.	E862289	-1	6
TG15-01	7.62	9.14	Orange	QMS	DG	3	25	Qtz veins ~25% of interval. Dominant colour is orange reflecting oxidation but there is also light grey with patchy green staining reflecting a weak chlorite alteration. Possible pin head silvery sx. Moderate oxidation, minimal chlorte alteration. 2% pyrite mineralization.	E862291	-1	10
TG15-01	9.14	10.67	Grey	QMS	DG	2	2	Muscovite very minor @5% of rock. Mainly consists of layers of qtz. Pyrite is fine grained and irregular throughout the rock. Qtz vein: 2%. Some of the qtz bands exhibit a brown colour. Minimal oxidation, 1 % pyrite mineralization.	E862292	-1	13
TG15-01	10.67	12.19	Grey	QBS	DG	0	0	Qtz bands are 1-3mm thick and biotite are 0.25-0.5mm. Also see biotite forming very thin septa between quartz bands. Pyrite occurs in schist. Very small pin head domains of sx. Biotite 10%. 90% quartz. Minimal chlorite alteration. 1% pyrite mineralization, 0.1 po mineralization.	E862293	9	10
TG15-01	12.19	13.72	Grey	QBS	DG	0	3	Qtz veining comprises 3% of interval. Biotite ~10%. Within one vein fragment there is py but also possible silvery grey sx - trace so 0.5%. 95% quartz. 1% pyrite, 0.5%po mineralization.	E862294	24	11
TG15-01	13.72	15.24	Grey	QBS	DG	0	0.1	Trace qtz veining. Two populations of colour, dark grey almost black and light grey. Light grey seems to be associated with minor qtz veining, so possible silicification. ~5% bitoite; very much a quartz rich rock. 95% quartz. 1% pyrite, 0.5% po mineralization.	E862295	14	9
TG15-01	15.24	16.76	Grey	QBS	DG	0	12	Qtz veining ~12%. Minimal chlorite alteration. 2% pyrite, 0.5% po mineralization. **During this run we got hit by a major hail and rain storm.	E862296	20	45
TG15-01	16.76	18.29	White	QBS	DG	4	7	Assuming it is the qtz biotite schist but cannot tell because of the level of oxidation and amount of clay in the interval. Half of chips comprised of clay-mud masses ~0.5-1cm in size. Oxidation went from nothing to strong. Possible fault zone intersected. Veining comprises ~7% of interval. Intense oxidation. Moderate clay alteration. 1% pyrite mineralization.	E862297	28	164

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
TG15-01	18.29	19.81	Grey	QBS	DG	2	35	Whitish grey mixture of weakly to moderately silicified schist. Qtz veins ~35% of interval. Veins show a weak level of oxidation. Pyrite occurs in veins and schist. Interesting because a vein system was intersected on margin of fault. Minimal oxidation. Moderate silicification alteration. 2% pyrite mineralization.	E862298	9	42
TG15-01	19.81	21.34	Grey	QBS	DG	0	15	Qtz veins 15%. Minor chlorite occurs in the veins. 755 quartz. Minimal chlorite alteration. 1% pyrite mineralization.	E862299	-1	21
TG15-01	21.34	22.86	Grey	QBS	DG	0	3	Qtz veining ~3%. Biotite 2%. 92% quartz. 0.5% pyrite mineralization.	E862300	67	821
TG15-01	22.86	24.38	Grey	QBS	DG	0	7	Qtz veins: 7% / Biotite ~3%. 90% quartz. 0.5% pyrite mineralization.	E862301	8	23
TG15-01	24.38	25.91	Grey	QBS	DG	0	10	Qtz veins ~10%. Pyrite and po within the veins and purite with host schist. 0.5% pyrite and 0.3% po mineralization. **30 minute shut down for lightning in area. Water was encountered as soon as hole was restored.	E862302	-1	12
TG15-01	25.91	27.43	Grey	QBS	DG	0	10	Very minor biotite except for rare bands/septa. Qtz veins ~10%. 1% pyrite and 0.5% po mineralization.	E862303	9	24
TG15-01	27.43	28.96	Grey	QBS	DG	0	3	3% qtz veins. 0.5% pyrite and 1% po mineralization.	E862304	11	17
TG15-01	28.96	30.48	Grey	QBS	DG	0	10	Mixed grey and white colour rock reflecting qtz veins and host schist. Minimal silicification. 0.5% pyrite mineralization.	E862305	-1	6
TG15-01	30.48	32	Grey	QBS	DG	0	0	Very light grey silicified schist. Hard to make out original fabric due to intense silicification. *Changed to no silica--very siliceous, almost quartzite. Same for 105 below. 0.5% pyrite mineralization.	E862306	5	5
TG15-01	32	33.53	Grey	QBS	DG	0	0	Some very minor oxidation. Weak schistosity still observed so interpreting to be silicified schist. ~1% biotite. 97% quartz. 2% pyrite mineralization.	E862307	-1	5
TG15-01	33.53	35.05	Grey	QZT	DG	2	0	Cannot recognize what was interpreted as schistosity above. There interpreted to be massive quartzite. Rock is still very light grey and bleached looking but interpreted silica alteration could actually be a different rock type. Py is fine grained and disseminated., 3%. Weak oxidation is developing in unit.	E862308	-1	7
TG15-01	35.05	36.58	Grey	QMS	DG	0	0	Quartz rich muscovite schist. Observe very thin septa of muscovite. Muscovite ~2% of rock. 96% quartz, 2% pyrite mineralization.	E862309	8	10
TG15-01	36.58	38.1	Grey	QZT	DG	0	0	Quartzite - 60% mixed with QBS. Rock has a septa and salt and pepper colour due to mixed lithologies. Pyrite is dissminated through the quartzite. Still can see a minor imprint of muscovite imparting a weak schistosity. 2% pyrite mineralization.	E862310	36	507
TG15-01	38.1	39.62	Grey	QBS	DG	0	0	Pyrite forming wisps aligned in foliation and also as veinlets 0.5-1mm wide. No alteration or oxidation. Slight green colour reflecting possible weak chlorite alteration. Crossed a lithology contact again. 3% pyrite mineralization.	E862311	26	13
TG15-01	39.62	41.15	Grey	QBS	DG	0	0	Very siliceous rock with little biotite. Mainly quartz. Exhibits a slight green trace - possible chlorite alteration? Pyrite is dissminated within rock. Very rare qtz veins with pyrite preserved along vein margins. 4% pyrite mineralization.	E862312	106	143
TG15-01	41.15	42.67	Grey	QMS	DG	0	0	Dominated (slightly!) by a 55-60% qtz-muscovite schist. Mainly quartz with very minor mica layers. Mixed with dark grey/black QBS. Colour is a salt and pepper. 4% pyrite mineralization.	E862313	19	42
TG15-01	42.67	44.2	Grey	QZT	DP	1	1	Light grey; dusty sooty pyrite (?); sample entirely Qtzt--> only 1 fragment QBS seen. Granular siliceous rock. Trace chlorite and sericite alteration. 0.5% pyrite mineralization.	E862314	50	114
TG15-01	44.2	45.72	Grey	QZT	DP	1	1	QBS 60%, 40% qtzt. Medium grey. Chlorite alteration in QBS but not seen in quartzite. Trace chlorite and sericite alteration. 0.5% py and 0.5% po mineralization.	E862315	259	525
TG15-01	45.72	47.24	Grey	QBS	DP	1	0	Light grey. Zero QBS. Py as sooty streaks (90%) and subhedral crystals (10%). Pyrite streaks can be perpendicular to foliation. Trace carbonation. 3% pyrite mineralization.	E862316	103	163
TG15-01	47.24	48.77	Grey	QZT	DP	0	0	Light grey - much less pyrite than previous interval. Here pyrite sooty, very fine dissminated. Trace carbonates, trace biotite and sericite alteration. 0.5% pyrite mineralization.	E862317	288	899
TG15-01	48.77	50.29	Grey	QZT	DP	0	0		E862318	40	124
TG15-01	50.29	51.82	Grey	QZT	DP	0.1	0	Light grey; pyrite as spots ~1mm across with faint margins. Trace oxidation (1mm on one fracture site), Trace sericite alteration. 0.5% pyrite mineralization.	E862319	130	229
TG15-01	51.82	53.34	Grey	QZT	DP	0	0	Light grey; pyrite mostly sooty. Less sulphide than last interval. Weak biotite and sericite alteration. 0.3% pyrite mineralization.	E862321	37	90
TG15-01	53.34	54.86	Grey	QZT	DP	0.1	0	Light grey; pyrite sooty, very fine disseminated and spots. Rock has greenish (chlorite?) tinge. Trace oxidation and carbonates. Weak chlorite and sericite alteration. 0.5% pyrite mineralization.	E862322	11	8
TG15-01	54.86	56.39	Grey	QZT	DP	0	0	Qtzt 75%, 25% QBS. Light grey qtzt. Medium greenish (chlorite) grey QBS. 0.5% carbonates. Trace chlorite, weak sericite alteration. 0.4% pyrite alteration.	E862323	10	16
TG15-01	56.39	57.91	Grey	QZT	DP	0	0	QBS 10%. Light to medium grey; pyrite sooty, very fine disseminated. Weak 1% carbonates, weak chlorite alteration. 1.5% pyrite mineralization.	E862324	9	25
TG15-01	57.91	59.44	Grey	CS	DP	0	0	Medium green-grey. Off-white calcite veinlets to 0.5mm wide. 0.4% carbonates, trace py/po mineralization.	E862325	7	13
TG15-01	59.44	60.96	Grey	QZT	DP	0	0	Medium grey; say 10% very fine biotite. Rare po. Trace chlorite and biotite alteration. Trace po mineralization.	E862326	8	7
TG15-01	60.96	62.48	Grey	QZT	DP	0	0	Light greenish grey; py as irregular masses, rare subhedral crystal. Trace biotite alteration. 2% pyrite mineralization.	E862327	12	6
TG15-01	62.48	64.01	Grey	QZT	DP	0	0	Medium grey; po very fine dissminated; water seen at 209' but sample OK. Mostly damp, not wet. Weak biotite alteration. Trace pyrite, 2% po mineralization.	E862328	89	8
TG15-01	64.01	65.53	Grey	QBS	DP	0	0.5	Damp sample. No sulphides seen within vein quartz. Poor light. 0.5% quartz. Weak biotite alteration, 0.5% po mineralization. **Stopped at 212.5 feet to change bit.	E862329	82	21
TG15-01	65.53	67.06	Grey	QZT	DP	0	0	Light brownish grey. No sulphides seen. Weak biotite alteration.	E862331	7	-1
TG15-01	67.06	68.58	Grey	QZT	DP	0	0	10% QBS. No sulphides. Weak biotite alteration.	E862332	9	4
TG15-01	68.58	70.1	Grey	QZT	DP	0	0	0.5% carbonates, weak biotite alteration. Trace py/po mineralization.	E862333	240	1350
TG15-01	70.1	71.63	Grey	QBS	DP	0	0	10% qtzt; medium brownish grey. Weak biotite alteration. Trace po mineralization.	E862334	13	7

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
TG15-01	71.63	73.15	Grey	QZT	DP	0	0	Light brownish grey; ~uniform. Weak biotite alteration.	E862335	7	6
TG15-01	73.15	74.68	Grey	QZT	DP	0	0	Medium brownish grey; no sulphides seen. Weak biotite alteration.	E862336	5	8
TG15-01	74.68	76.2	Grey	QZT	DP	0	0	30% QBS. No sulphides. Weak to minimal biotite alteration.	E862337	10	9
TG15-01	76.2	77.72	Grey	QZT	DP	0	0	20% QBS. Damp Sample. Minimal biotite alteration. Trace po alteration.	E862338	20	21
TG15-01	77.72	79.25	Grey	QZT	DP	0	0	Qtzt 50%, QBS 50%. Damp Sample. Minimal biotite alteration.	E862339	28	7
TG15-01	79.25	80.77	Grey	QZT	DP	0	0	Qtzt 90%, QBS 10%. Damp Sample. Minimal Biotite alteration.	E862340	12	4
TG15-01	80.77	82.3	Grey	QZT	DP	0	0	Damp. Medium to light brownish grey; no sulphides. Trace chlorite, weak biotite alteration.	E862341	10	4
TG15-01	82.3	83.82	Grey	QBS	DP	0	0	Damp. Minimal biotite alteration.	E862342	8	4
TG15-01	83.82	85.34	Grey	QZT	DP	0	0	70% qtzt, 30% QBS. Medium brownish grey; sulphides as irregular wispy masses to 0.75mm across. Minimal biotite alteration. 0.1% py/po mineralization.	E862343	-1	6
TG15-01	85.34	86.87	Grey	QBS	DP	0	0	Medium to dark brownish grey with qtzt bands to 3mm wide. Muddy sample. Minimal biotite alteration. 0.1 pyrite, 0.1 po mineralization.	E862344	-1	11
TG15-01	86.87	88.39	Grey	QZT	DP	0	0	Light brownish grey. Muddy. Minimal biotite. 0.3% po mineralization.	E862345	-1	6
TG15-01	88.39	89.92	Grey	QZT	DP	0	0	As above - muddy. Minimal biotite, 0.3% po mineralization.	E862346	8	11
TG15-01	89.92	91.44	Grey	QZT	DP	0	0	As above - muddy. Minimal biotite, 0.3% po mineralization.	E862347	24	8
TG15-01	91.44	92.96	Grey	QZT	DP	0	0	As above - muddy. Minimal biotite, 0.2% po mineralization.	E862348	5	7
TG15-01	92.96	94.49	Grey	QZT	DP	0	0	Light grey and muddy. Po blebs, specks to 1mm. Weak biotite alteration. 0.1% pyrite, 0.5% po mineralization.	E862349	18	7
TG15-01	94.49	96.01	Grey	QZT	DG	0	0	Light grey. Very subtle layering observed in the 1-2mm scale. Light grey, massive quartz layers with interlayered dark brown-grey quartz layers possible with mica in trace - 0.5% amounts. 0.5% pyrite mineralization.	E862351	13	8
TG15-01	96.01	97.54	Brown	QZT	DG	0	0	Massive sulphide-pyrrhotite comprising 60-70% of the chips. Mixed and visibly layered with quartzite and QBS. Select larger grains and 1.5cm in size show layers between massive sx and greenish black quartzite. In that sample layer of po is at least 0.75cm thick. Numerous other 1cm size grains comprising 100% po. The chips of massive sx are actually ~75-85% sx and 15-25% greenish quartz, forming medium to coarse grained crystals. Layer also comprised of 3-4% quartz vein fragments. The greenish colouring interpreted to be weak chlorite alteration.	E862352	15	8
TG15-01	97.54	99.06	Grey	QBS	DG	0	0	Mixed population of grians; population of brownish grey quartz biotite schist with very minor biotite (2-4%) and pale green quartzite. 2-4% po disseminated within both rock types. Quartz vein ~3-4% of interval. Minimal chlorite alteration. 3% po mineralization.	E862353	79	12
YCS15-01	0	1.52		CAS					No Sample		
YCS15-01	1.52	3.05	Grey	QBS	DG	3			E863104	-1	19
YCS15-01	3.05	4.57	Grey	QBS	DG	3		Strong foliation. Quartz bands ~ 1-3mm thick	E863105	-1	13
YCS15-01	4.57	6.1	Grey	QBS	DG	2	1	There is light grey quartzite mixed with the QBS chips.	E863106	-1	29
YCS15-01	6.1	7.62	Grey	QBS	DG	2		Non magnetic	E863107	-1	23
YCS15-01	7.62	9.14	Grey	QBS	DG	2			E863108	5	43
YCS15-01	9.14	10.67	Grey	QBS	DG	2		Non magnetic	E863109	-1	34
YCS15-01	10.67	12.19	Grey	QBS	DG	3			E863110	-1	29
YCS15-01	12.19	13.72	Black	ARG	DG			Appears to be mainly quartz feldspathic but cannot make out individual grains. Could have fine grained biotite +/- chlorite.	E863111	7	29
YCS15-01	13.72	15.24	Black	ARG	DG	1		Fine grained	E863112	-1	35
YCS15-01	15.24	16.76	Grey	ARG	DG	2		Can subtly distinguish quartz and feldspar grains	E863113	5	53
YCS15-01	16.76	18.29	Grey	QBS	DG	2			E863114	-1	125
YCS15-01	18.29	19.81	Grey	QBS	DG	3			E863115	1200	2340
YCS15-01	19.81	21.34	Grey	QBS	DG				E863116	5000	4070
YCS15-01	21.34	22.86	Grey	QBS	DG				E863117	26	63
YCS15-01	22.86	24.38	Grey	QBS	DG				E863118	14	26
YCS15-01	24.38	25.91	Grey	QBS	DG		3	Minor sx grains observed (py). Could have been mod silicified QBS or a light grey quartzite	E863119	15	33
YCS15-01	25.91	27.43	Grey	QBS	DG		0.1	Trace quartz vein chips	E863121	8	18
YCS15-01	27.43	28.96	Grey	QBS	DG				E863122	6	8
YCS15-01	28.96	30.48	Grey	QBS	DG				E863123	7	18
YCS15-01	30.48	32	Grey	QBS	DG			Rare chlorite rich grains	E863124	-1	13
YCS15-01	32	33.53	Grey	QBS	DG		0.1	Rare quartz veins occurs	E863125	5	41
YCS15-01	33.53	35.05	Grey	QBS	DG		2		E863126	5	39
YCS15-01	35.05	36.58	Grey	QBS	DG		3		E863127	-1	16
YCS15-01	36.58	38.1	Grey	QBS	DG				E863128	15	11
YCS15-01	38.1	39.62	Grey	QBS	DG		2	2% quartz vein. Fine grained disseminated pyrite occurs only in the veins.	E863129	6	22
YCS15-01	39.62	41.15	Grey	QBS	DG		0.1		E863131	-1	41
YCS15-01	41.15	42.67	Grey	QBS	DG				E863132	6	17
YCS15-01	42.67	44.2	Grey	QBS	DG		5	5% smokey grey quartz veins mixed with QBS. Rare greenish grey quartz grains - quartzite.	E863133	12	15
YCS15-01	44.2	45.72	Grey	QBS	DG		2	Trace sx with the vein, no sx in the schist.	E863134	-1	9
YCS15-01	45.72	47.24	Grey	QBS	DG		4		E863135	-1	18

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-01	47.24	48.77	Grey	QBS	DG		0.1	Trace quartz vein. 40% of interval comprised of greenish grey quartzite. Very rare biotite crystals within this translucent rock. Trace sx in the quartzite.	E863136	-1	7
YCS15-01	48.77	50.29	Grey	QZT	DG				E863137	-1	6
YCS15-01	50.29	51.82	Grey	QBS	DG		5	~35% Quartzite, 5% quartz veining. Trace sx in the vein.	E863138	-1	8
YCS15-01	51.82	53.34	Grey	QBS	DG			25% of chips comprised of greenish grey quartzite. Select quartz and vein material also show a green colour. Almost epidote cp;pir but too bright green chlorite but very hard so possibly epidote.	E863139	-1	6
YCS15-01	53.34	54.86	Grey	QBS	DG		4	4% quartz veins. Same almost pistachio green colour. Most likely epidote. Trace sx in the veins.	E863140	-1	8
YCS15-01	54.86	56.39	Grey	QBS	DG		0.1	Trace quartz veins. Minor chlorite in select vein material. 3% quartzite.	E863141	-1	11
YCS15-01	56.39	57.91	Grey	QBS	DG		3	3% quartz veins with trace sx within vein. No oxidation. No sx within schist. Biotite comprises 15-20% of the rock.	E863142	5	6
YCS15-01	57.91	59.44	Grey	QBS	DG		1	1% quartz veins. Weak pervasive chlorite alteration.	E863143	-1	9
YCS15-01	59.44	60.96	Grey	QBS	DG		1	1% quartz veins. 5% grey green quartzite.	E863144	5	8
YCS15-01	60.96	62.48	Grey	QBS	DG		3	3% quartz veins.	E863145	-1	10
YCS15-01	62.48	64.01	Green	QBS	DG		2	2% quartz veins	E863146	-1	10
YCS15-01	64.01	65.53	Green	CS	DG		0.1	Chlorite schist. Trace quartz veins.	E863147	-1	6
YCS15-01	65.53	67.06	Grey	QZT	DG			35% of interval is chlorite schist. The quartzite is light grey and in some chips resembles quartz vein but looks to be foliated and like it is comprised of grains. Trace sx.	E863148	-1	6
YCS15-01	67.06	68.58	Grey	QZT	DG			Light grey quartzite with no sx, 35% of interval is QBS.	E863149	-1	6
YCS15-01	68.58	70.1	Grey	QZT	DG			Quartz chlorite schist. 25% of chips are light grey quartzite. Quartzite contains trace pyrite.	E863151	-1	7
YCS15-01	70.1	71.63	White	CS	DG		40	40% quartz veining +/- quartzite mixed in. Rock is a salt and pepper but with green and white	E863152	-1	18
YCS15-01	71.63	73.15	Green	CS	DG			Veins absent. Chips comprised of quartz and chlorite	E863153	-1	10
YCS15-01	73.15	74.68	Green	CS	DG		2	2% quartz veins with ~1% pyrite only in the vein material.	E863154	26	407
YCS15-01	74.68	76.2	Green	CS	DG			No veining. Trace py.	E863155	-1	15
YCS15-01	76.2	77.72	Green	CS	DG			No veining. Trace py.	E863156	6	10
YCS15-01	77.72	79.25		QZT	DG			30% chlorite schist. Difficult to separate what is quartz vein and what is quartzite. Looks like there could be ~5-10% quartz vein with py and unknown red-brown mineral. Also could be a pinhead silvery sx mixed with the red-brown mineral. Could be a hematite stain on the quartz.	E863157	-1	6
YCS15-01	79.25	80.77		QZT	DG			Quartz chlorite schist. Grains are dominantly quartz with flecks of chlorite throughout. Red-brown staining on select chips could represent a weak hematite stain/alteration.	E863158	-1	6
YCS15-01	80.77	82.3		QZT	DG			Mainly quartz grains with a 10-15% black inclusions - looks like chlorite. Rare red-brown staining of select grains. NO sx.	E863159	6	7
YCS15-01	82.3	83.82		QZT	DG			No sulphides seen	E863161	-1	9
YCS15-01	83.82	85.34		QZT	DP			No sulphides	E863162	-1	-1
YCS15-01	85.34	86.87		QZT	DP			Trace py, chl	E863163	-1	5
YCS15-01	86.87	88.39		QZT	DP	1		Oxidized (~10% of chips, stained by orange limn).	E863164	9	11
YCS15-01	88.39	89.92		QZT	DP	3		Orange stained limonite in folds; less biotite than above. Zone?	E863165	-1	16
YCS15-01	89.92	91.44		QZT	DP	2		Slightly less oxidized than previous interval.	E863166	8	-1
YCS15-01	91.44	92.96		QBS	DP	2		Still some limonite-stained silt-size drill return.	E863167	-1	-1
YCS15-01	92.96	94.49	Grey	QBS	DP			Dark grey, fresh. 1% quartz veinlets.	E863168	6	-1
YCS15-01	94.49	96.01	Grey	QBS	DP			30% quartzite with ~5% fresh biotite as seen higher in hole.	E863169	-1	-1
YCS15-01	96.01	97.54	Grey	QBS	DP			As above	E863170	-1	-1
YCS15-01	97.54	99.06	Grey	QBS	DP			As above	E863171	5	-1
YCS15-01	99.06	100.58		QZT	DP			No Sample			
YCS15-01	100.58	102.11		QZT	DP			20% QBS fragments; traces red hem dusty disseminated in quartzite.	E863172	-1	34
YCS15-01	102.11	103.63		QBS	DP			Weak chlorite; traces pyrite	E863173	-1	4
YCS15-01	103.63	105.16		QZT	DP			10% quartzite	E863174	-1	-1
YCS15-01	105.16	106.68		QZT	DP			3% biotite, 1% chlorite, traces hematite, sericite.	E863175	-1	-1
YCS15-01	106.68	108.2	Grey	QZT	DP			Light grey, fine to medium grained, fresh	E863176	-1	-1
YCS15-01	108.2	109.73	Grey	QZT	DP			As above	E863177	10	-1
YCS15-01	109.73	111.25	Grey	QZT	DP			As above except here trace pyrite seen associated with very weak green chlorite (sericite?) alteration.	E863178	7	4
YCS15-01	111.25	112.78		QZT	DP			Pervasive weak chlorite altered but no sulphides seen	E863179	-1	-1
YCS15-01	112.78	114.3		QZT	DP			Less chlorite than last interval. No sulphides seen.	E863181	-1	7
YCS15-01	114.3	115.82		QZT	DP			As above	E863182	-1	5
YCS15-01	115.82	117.35		QZT	DP			As above except rare traces red hematite.	E863183	13	6
YCS15-01	117.35	118.87		QZT	DP			1% biotite flakes; still no sulphides seen.	E863184	-1	5
YCS15-01	118.87	120.4	Grey	QZT	DP			Pale grey, fine grained; no sulphides	E863185	8	-1
YCS15-01	120.4	121.92	Grey	QZT	DP			Pale grey, fine grained; no sulphides	E863186	6	3
YCS15-01	121.92	123.44	Grey	QZT	DP			As above except here traces red hematite and rare chlorite	E863187	6	4
YCS15-01	123.44	124.97	Grey	QZT	DP			As above; more hematite (0.2%)	E863188	-1	-1
YCS15-01	124.97	126.49	Grey	QZT	DP			As above	E863189	5	17
YCS15-01	126.49	128.02		QZT	DP			1% chlorite, 0.5% biotite; rare hematite; no sulphides	E863191	-1	27

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-01	128.02	129.54		CS	DP			~0.5% chlorite, 1% biotite; no sulphides seen nor hematite seen	E863192	-1	8
YCS15-01	129.54	131.06	Green	CS	DP			Light green; pervasive chlorite alteration. Say 75-90% quartz so maybe better described as quartzite. No sulphides.	E863193	6	11
YCS15-01	131.06	132.59	Grey	QZT	DP			65% interval as for previous; 35% pale brownish white quartzite. No sulphides seen.	E863194	-1	4
YCS15-01	132.59	134.11		QBS	DP			Pale brownish grey fine grained, no sulphides. Approximately 5% QBS.	E863195	-1	11
YCS15-01	134.11	135.64	Grey	QBS	DP			40% CS as from 425-430'. No sulphides seen.	E863196	10	24
YCS15-01	135.64	137.16	Grey	CS	DP			Medium brownish grey; fines in return light brown; fresh, no sulphides.	E863197	16	155
YCS15-01	137.16	138.68	Grey	CS	DP			40% QBS as above	E863198	7	23
YCS15-01	138.68	140.21	Green	CS	DP			5% QBS. Medium grey-green, no sulphides.	E863199	5	14
YCS15-01	140.21	141.73		QBS	DP			10% QBS	E863200	20	37
YCS15-01	141.73	143.26	Grey	QBS	DP			Dark brownish grey; fresh; no sulphides seen.	E863201	19	76
YCS15-01	143.26	144.78		QBS	DP			As above except 0.5% calcite veinlets	E863202	20	193
YCS15-01	144.78	146.3		QBS	DP			No carb veinlets; fresh.	E863203	20	130
YCS15-01	146.3	147.83		QBS	DP			As above	E863204	7	109
YCS15-01	147.83	149.35		QBS	DP			As above	E863205	9	69
YCS15-01	149.35	150.88		QBS	DP			As above	E863206	98	1630
YCS15-01	150.88	152.4		QBS	DP			As above	E863207	6	78
YCS15-01	152.4	153.92		QBS	DP			As above	E863208	6	47
YCS15-01	153.92	155.45		QBS	DP			As above	E863209	-1	44
YCS15-01	155.45	156.97	Grey	QZT	DP			Interval 35% pale grey, fine grained mineralized quartzite.	E863210	-1	51
YCS15-01	156.97	158.5	White	QZT	DP			60% pale greenish white quartzite; 40% medium brownish grey QBS; no sulphides.	E863211	-1	42
YCS15-01	158.5	160.02	White	QZT	DP			Pale greenish white; no sulphides seen	E863212	8	46
YCS15-01	160.02	161.54	White	QZT	DP			As above	E863213	-1	15
YCS15-01	161.54	163.07	White	QZT	DP			As above except here 5% QBS	E863214	5	8
YCS15-01	163.07	164.59		QZT	DP			3% QBS; no sulphides	E863215	-1	8
YCS15-01	164.59	166.12		QZT	DP			No QBS; rare bleb pyrite 1.5mm across	E863216	-1	14
YCS15-01	166.12	167.64	White	QBS	DP			Pale greenish white	E863217	-1	8
YCS15-01	167.64	169.16	Green	QZT	DP			Interval 35% greenish quartzite as above--quartzite contains 0.2% blebby pyrite.	E863218	-1	16
YCS15-02	0	1.52		CAS					No Sample		
YCS15-02	1.52	3.05	Grey	QBS	DG	3			E863219	13	324
YCS15-02	3.05	4.57	Grey	QBS	DG	3			E863221	28	801
YCS15-02	4.57	6.1	Orange	QBS	DG	4			E863222	6	131
YCS15-02	6.1	7.62	Orange	QMS	DG	4		Patchy weak silicification of selected chips.	E863223	-1	48
YCS15-02	7.62	9.14	Grey	QBS	DG	2	3	3% quartz vein	E863224	-1	43
YCS15-02	9.14	10.67	Grey	QBS	DG	2	3	20% quartzite, 3% quartz vein	E863225	-1	15
YCS15-02	10.67	12.19	Grey	QZT	DG		2	25% QBS, 2% quartz vein	E863226	-1	8
YCS15-02	12.19	13.72	Grey	QBS	DG			3% quartzite	E863227	-1	12
YCS15-02	13.72	15.24	Grey	QBS	DG			No sulphides. Non-magnetic.	E863228	-1	10
YCS15-02	15.24	16.76	Grey	QBS	DG			No sulphides. Non-magnetic.	E863229	-1	8
YCS15-02	16.76	18.29	Grey	QBS	DG		1		E863231	-1	8
YCS15-02	18.29	19.81	Grey	QBS	DG		0.5		E863232	5	7
YCS15-02	19.81	21.34	Grey	QBS	DG		0.5	Slightly foliated.	E863233	-1	10
YCS15-02	21.34	22.86	Grey	QBS	DG			No sulphides in any chips. Non-magnetic.	E863234	10	10
YCS15-02	22.86	24.38	Grey	QBS	DG	2		Non-magnetic	E863235	-1	14
YCS15-02	24.38	25.91	Grey	QBS	DG	1		Very weakly oxidized on selected chip surfaces. Non-magnetic.	E863236	6	12
YCS15-02	25.91	27.43	Grey	QBS	DG	1			E863237	-1	23
YCS15-02	27.43	28.96	Grey	QBS	DG	1	2		E863238	-1	13
YCS15-02	28.96	30.48	Grey	QBS	DG			Trace oxidation.	E863239	6	19
YCS15-02	30.48	32	Grey	QBS	DG	2	1		E863240	-1	56
YCS15-02	32	33.53	Grey	QBS	DG	1	2	~20% of chips are quartzite, 2% quartz veins.	E863241	-1	47
YCS15-02	33.53	35.05	Grey	QBS	DG	1	0.5		E863242	-1	62
YCS15-02	35.05	36.58	Grey	QBS	DG		0.5		E863243	-1	20
YCS15-02	36.58	38.1	Grey	QBS	DG		1		E863244	8	18
YCS15-02	38.1	39.62	Grey	QBS	DG		2		E863245	-1	12
YCS15-02	39.62	41.15	Grey	QBS	DG	1	1		E863246	-1	24
YCS15-02	41.15	42.67	Grey	QBS	DG		1		E863247	-1	27
YCS15-02	42.67	44.2	Grey	QBS	DG		2		E863248	-1	27
YCS15-02	44.2	45.72	Grey	QBS	DG		0.5	Slightly foliated. Non-magnetic.	E863249	-1	13
YCS15-02	45.72	47.24	Grey	QBS	DG		2	Pyrite occurs within the quartz veins	E863251	-1	23
YCS15-02	47.24	48.77	Grey	QBS	DG		1		E863252	-1	37

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-02	48.77	50.29	Grey	QBS	DG				E863253	-1	15
YCS15-02	50.29	51.82	Grey	QBS	DG		2		E863254	-1	15
YCS15-02	51.82	53.34	Grey	QBS	DG		1	10% of interval comprised of greenish grey quartzite	E863255	-1	18
YCS15-02	53.34	54.86	Grey	QZT	DG		0.5	20% of chips comprised of QBS	E863256	-1	21
YCS15-02	54.86	56.39	Grey	QZT	DG			30% of chips comprised of QBS. Weak pervasive chlorite alteration.	E863257	-1	14
YCS15-02	56.39	57.91	Grey	QZT	DG		1	35% of chips comprised of QBS. Very rare chips of quartzite exhibit a red brown colour interpreted to be hematite.	E863258	-1	7
YCS15-02	57.91	59.44	Grey	QZT	DG		15	Mixture of rock types. Dominated by quartzite but also ~15% quartz vein and 20% QBS. The quartzite is very white and is hard to discern from the quartzite. Quartzite typically has inclusions of biotite +/- chlorite. Rare quartzite chips contain a red-brown colour possibly reflecting hematite alteration.	E863259	-1	6
YCS15-02	59.44	60.96	Grey	CS	DG		2	Chips comprised of quartz and chlorite with chlorite content varying between 5-50%.	E863261	-1	9
YCS15-02	60.96	62.48	Green	CS	DG		1	10-15% of chips comprised of quartzite. Rock is non-magnetic.	E863262	-1	11
YCS15-02	62.48	64.01	Grey	QZT	DG			30% of chips comprised of chlorite schist.	E863263	-1	7
YCS15-02	64.01	65.53	Grey	QZT	DG		0.5	Quartzite varies from no inclusions and 100% quartz to up to 10% chlorite/biotite. Also variable green to grey. No sx anywhere. Single hematite stained chip.	E863264	-1	7
YCS15-02	65.53	67.06		QZT	DG		1	Typically all chips are grey with minor component showing a green colour. Very rare chips exhibit the red-brown colouration/staining hematite. Very rare pyrite fine grained within the quartz.	E863265	-1	15
YCS15-02	67.06	68.58	Grey	QZT	DG		1	~15% of interval comprised of CS. Rare quartzite clasts show the red-brown colouration.	E863266	-1	15
YCS15-02	68.58	70.1	Grey	QZT	DG		4	15% CS. Quartzite is variably grey to green in colour. No sulphides. Non-magnetic.	E863267	-1	10
YCS15-02	70.1	71.63		QZT	DG			All quartzite no other rock types. Trace py occurring as fine grained pinhead size crystals.	E863268	-1	20
YCS15-02	71.63	73.15	Grey	QZT	DG		0.5	Same rock as above but py content dramatically increased to ~7%. Typically fine grained and disseminated but rarely as euhedral 0.5mm crystals.	E863269	877	4140
YCS15-02	73.15	74.68	Grey	QZT	DG			As above but pyrite decreases. Rare red-brown staining. Rare chlorite chips.	E863270	117	824
YCS15-02	74.68	76.2	Grey	QZT	DG		0.5	Pyrite decreases even more to only rare quartzite chips. Fine grained and anhedral.	E863271	-1	16
YCS15-02	76.2	77.72	Grey	QZT	DG			Rare quartzite chips contain greenish stain - chlorite alteration?	E863272	-1	11
YCS15-02	77.72	79.25	Grey	QZT	DG			Majority of quartzite chips contain trace to 10% chlorite inclusions. No sulphides.	E863273	-1	10
YCS15-02	79.25	80.77	Grey	QBS	DG		1	35% of chips comprised of quartzite. No sulphides.	E863274	-1	16
YCS15-02	80.77	82.3	Grey	QBS	DG		1	~40% of chips comprised of quartzite. No sulphides.	E863275	-1	8
YCS15-02	82.3	83.82	Grey	QBS	DG			35% of chips comprised of quartzite. A subset of those quartzite chips exhibit a pistachio green staining possible due to a weak epidote alteration. Also see a rare red-brown staining possibly due to hematite.	E863276	-1	9
YCS15-02	83.82	85.34	Grey	QZT	DG			Only trace QBS. No sulphides. Subtle green colour reflects inclusion of chlorite within the quartzite chips.	E863277	-1	5
YCS15-02	85.34	86.87	Grey	QZT	DG			20% CS. Trace py within the quartzite.	E863278	-1	11
YCS15-02	86.87	88.39	Grey	QBS	DG			30% is QTZE.	E863279	-1	9
YCS15-02	88.39	89.92	Grey	QZT	DG			35% QBS	E863281	-1	6
YCS15-02	89.92	91.44	Grey	QBS	DG			~30% Quartzite	E863282	-1	10
YCS15-02	91.44	92.96	Grey	QBS	DG			~25% Quartzite	E863283	-1	11
YCS15-02	92.96	94.49	Grey	QZT	DG			~15% QBS	E863284	-1	9
YCS15-02	94.49	96.01	Grey	ARG	DG			Argillite. Black, fine-grained rock. 35% of interval comprised of quartzite. Argillite is foliated.	E863285	-1	9
YCS15-02	96.01	97.54	Grey	QBS	DG			30% Quartzite	E863286	-1	8
YCS15-02	97.54	99.06	Grey	QZT	DG			35% QBS	E863287	-1	8
YCS15-02	99.06	100.58	Grey	QBS	DG			~20% of chips are quartzite. Seems to be gradational based on biotite content; some chips have no biotite at all and chips grade to ~20-30% in the QBS.	E863288	-1	12
YCS15-02	100.58	102.11	Grey	QBS	DG		2	15% of interval is Quartzite.	E863289	-1	7
YCS15-02	102.11	103.63	Grey	QZT	DG		0.5	~35% of interval is QBS.	E863291	-1	-1
YCS15-02	103.63	105.16	Grey	QZT	DG			~3% QBS. Pyrite is fine grained and inconsistently occurring within the quartzite. Possibly occurs in very specific but rare horizons/beds.	E863292	-1	5
YCS15-02	105.16	106.68	Grey	QZT	DG			10% QBS.	E863293	-1	3
YCS15-02	106.68	108.2	Grey	QZT	DG			Pyrite is typically fine grained and pinhead size. Rare increases to 0.5mm and euhedral crystals. No QBS. 15-20% of the chips exhibit a diffuse green colour. Possibly chlorite alteration - weak.	E863294	-1	4
YCS15-02	108.2	109.73	Grey	QZT	DG		0.5	Subtle green colour reflecting minor chlorite in the quartzite. No sulphides.	E863295	-1	5
YCS15-02	109.73	111.25	Grey	QZT	DG		0.5	As above.	E863296	-1	-1
YCS15-02	111.25	112.78	Grey	QZT	DG		1	Minor greenish quartzite. Difficult to discern quartz vein from quartzite.	E863297	-1	4
YCS15-02	112.78	114.3	Grey	QZT	DG			Minor inclusions of biotite +/- chlorite in the quartzite. Pyrite is fine grained and pinhead sized and rarely is medium grained and euhedral.	E863298	-1	4
YCS15-02	114.3	115.82	Grey	QZT	DG			No quartzite with biotite or chlorite.	E863299	-1	-1
YCS15-02	115.82	117.35	Grey	QZT	DG			~20-30 of chips exhibit a green colour due to inclusions of chlorite.	E863300	-1	-1
YCS15-02	117.35	118.87	Grey	QZT	DG			Minor chlorite +/- biotite inclusions.	E863301	-1	-1
YCS15-02	118.87	120.4		QZT	DG			Trace pyrite. Minor chlorite +/- biotite in quartz grains.	E863302	5	9
YCS15-02	120.4	121.92	Grey	QZT	DG			~1-2% QBS.	E863303	-1	25
YCS15-02	121.92	123.44	Grey	QZT	DG		1	Quartz vein is massive and white while quartzite is granular and grey-green.	E863304	-1	-1

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-02	123.44	124.97	Grey	QZT	DG		0.5		E863305	-1	11
YCS15-02	124.97	126.49	Grey	QZT	DG			~20% of chips contain trace to 10% biotite inclusions.	E863306	-1	25
YCS15-02	126.49	128.02	Grey	QZT	DG			25% of chips are QBS	E863307	7	245
YCS15-02	128.02	129.54	Grey	QZT	DG			35% of chips are QBS	E863308	-1	99
YCS15-02	129.54	131.06	Grey	QBS	DG			~5% quartzite	E863309	-1	59
YCS15-02	131.06	132.59		QZT	DG			Minor biotite and chlorite inclusions in quartzite.	E863310	-1	59
YCS15-02	132.59	134.11	Grey	QZT	DG			2-4% QBS	E863311	16	74
YCS15-02	134.11	135.64	Grey	QZT	DG			Rare grains have fine grained disseminated pyrite	E863312	5	68
YCS15-02	135.64	137.16	Grey	QZT	DG			Slight increase in biotite content but very small % - still call it quartzite.	E863313	15	96
YCS15-02	137.16	138.68	Grey	QZT	DP			Light grey	E863314	13	48
YCS15-02	138.68	140.21	Grey	QBS	DP			Light grey; traces disseminated hematite interval 20% quartzite.	E863315	-1	66
YCS15-02	140.21	141.73		QZT	DP			Fine grained	E863316	6	141
YCS15-02	141.73	143.26		QZT	DP			As above except here 0.3% pyrite	E863317	10	245
YCS15-02	143.26	144.78		QZT	DP			Galena? 1 speck.	E863318	35	1170
YCS15-02	144.78	146.3		QZT	DP			Less sulphide than above.	E863319	9	470
YCS15-02	146.3	147.83		QZT	DP			Non magnetic. No po seen.	E863321	-1	117
YCS15-02	147.83	149.35		QZT	DP			As above except here py along fracture sites in addition to occurring as irregular masses. No po seen.	E863322	28	1130
YCS15-02	149.35	150.88	Grey	QZT	DP			Light brownish grey; 10% biotite.	E863323	-1	54
YCS15-02	150.88	152.4		QZT	DP				E863324	-1	36
YCS15-02	152.4	153.92		QZT	DP			As above except more Pyrite	E863325	-1	122
YCS15-02	153.92	155.45	Grey	QZT	DP			Pale green-grey; pyrite as euhedral, veinlet disseminated cubes.	E863326	-1	7
YCS15-02	155.45	156.97	Grey	QZT	DP			As above.	E863327	-1	4
YCS15-02	156.97	158.5	Grey	QZT	DP			As above.	E863328	-1	17
YCS15-02	158.5	160.02		QZT	DP			As above.	E863329	-1	12
YCS15-02	160.02	161.54	Brown	QZT	DP			Pale green- brown	E863331	-1	34
YCS15-02	161.54	163.07		QZT	DP				E863332	-1	5
YCS15-02	163.07	164.59		QZT	DP				E863333	-1	27
YCS15-02	164.59	166.12		QZT	DP			40% quartzite	E863334	-1	16
YCS15-02	166.12	167.64	Green	QZT	DP			Pale grey and green; 10% QBS	E863335	-1	24
YCS15-02	167.64	169.16	Grey	QZT	DP			Pale greenish grey	E863336	-1	12
YCS15-02	169.16	170.69	Grey	QZT	DP			Pale greenish grey; 5% QBS	E863337	-1	6
YCS15-02	170.69	172.21		QBS	DP			25% quartzite	E863338	-1	-1
YCS15-02	172.21	173.74		QBS	DP				E863339	-1	9
YCS15-02	173.74	175.26	Green	QZT	DP			Pale green, fine grained; no sulphides; 35% QBS.	E863340	-1	9
YCS15-02	175.26	176.78	Green	QZT	DP			As above except 5% QBS	E863341	-1	15
YCS15-02	176.78	178.31	Green	QZT	DP			As above except 3% QBS	E863342	-1	11
YCS15-02	178.31	179.83		QZT	DP			1% pyrite subhedral crystals to 1mm across.	E863343	-1	7
YCS15-02	179.83	181.36		QZT	DP			No sulphides seen; 3% QBS	E863344	-1	12
YCS15-02	181.36	182.88		QBS	DP			30% quartzite	E863345	-1	12
YCS15-02	182.88	184.4	Green	QZT	DP			Pale grayish green, no sulphides seen. 15% QBS.	E863346	-1	36
YCS15-02	184.4	185.93		QZT	DP			35% QBS; no sulphides seen	E863347	-1	36
YCS15-02	185.93	187.45		QZT	DP			10% QBS; 1.5% pyrite in quartzite	E863348	1300	3220
YCS15-02	187.45	188.98		QZT	DP			More pyrite than the last interval; 3% QBS	E863349	13	128
YCS15-02	188.98	190.5	White	QBS	DP	2		2% greyish white quartz veinlets. No sulphides seen.	E863351	-1	39
YCS15-02	190.5	192.02		QBS	DP			3% quartzite; no sulphides seen.	E863352	-1	58
YCS15-02	192.02	193.55		QBS	DP			3% quartzite; one subhedral pyrite grain 1mm across seen within quartz fragment.	E863353	-1	10
YCS15-02	193.55	195.07	Green	QBS	DP			3% quartzite. Quartzite weakly chlorite altered. Pale green. No sulphides seen.	E863354	7	7
YCS15-02	195.07	196.6	White	QBS	DP	1		1% greyish white vein quartz fragments which contain rare speck pyrite.	E863355	-1	25
YCS15-02	196.6	198.12	Grey	QBS	DP			Medium brownish grey; No sulphides seen	E863356	5	12
YCS15-02	198.12	199.64	Grey	QBS	DP			As above except rare vein quartz fragment.	E863357	-1	15
YCS15-02	199.64	201.17		QBS	DP			2% quartzite chips; no sulphides seen.	E863358	-1	11
YCS15-03	0	1.52		CAS					No Sample		
YCS15-03	1.52	3.05	Grey	QBS	DG			Very minor grains of quartzite. No sulphides. No veining.	E863359	-1	9
YCS15-03	3.05	4.57	Grey	QBS	DG	2		Minor quartzite chips mixed. No veining. No sulphides.	E863361	-1	25
YCS15-03	4.57	6.1	Grey	QBS	DG	2		~30% biotite.	E863362	-1	40
YCS15-03	6.1	7.62	Grey	QBS	DG	3		~25% biotite	E863363	-1	73
YCS15-03	7.62	9.14	Grey	QBS	DG	3		~25% biotite	E863364	-1	632
YCS15-03	9.14	10.67	Grey	QBS	KJ	3		~25% biotite	E863365	3900	6180
YCS15-03	10.67	12.19	Grey	QBS	KJ	3	1	~25% biotite, 20% muscovite	E863366	970	8770

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-03	12.19	13.72	Grey	QBS	KJ	4		~25% biotite, 5% muscovite	E863367	54	2240
YCS15-03	13.72	15.24	Grey	QBS	KJ	1	2	~25% biotite	E863368	12100	9780
YCS15-03	15.24	16.76	Grey	QBS	KJ	4	2	~25% biotite, small quartz grains.	E863369	183	484
YCS15-03	16.76	18.29	Grey	QBS	KJ	2		25-30% biotite	E863370	52	105
YCS15-03	18.29	19.81	Grey	QBS	KJ	1		30-35% biotite	E863371	36	38
YCS15-03	19.81	21.34	Grey	QBS	KJ	1		30% biotite, platy opposed to rounded.	E863372	13	17
YCS15-03	21.34	22.86	Grey	QBS	KJ	1		30% biotite, platy, 15% muscovite	E863373	7	9
YCS15-03	22.86	24.38	Grey	QBS	KJ	1	2	25% biotite, 20% quartz	E863374	-1	59
YCS15-03	24.38	27.43	Grey	QBS	KJ	2	1	Sample combined in the lab. 30% biotite, 5% quartz. 85-90: limonite clay.	E863375/E863376	-1	11
YCS15-03	27.43	28.96	Grey	QBS	KJ	1	2	30% biotite, 35% quartz, quartz grains with chlorite.	E863377	-1	12
YCS15-03	28.96	30.48	Grey	QBS	KJ	1	2	30% biotite, large quartz chips with chlorite.	E863378	5	15
YCS15-03	30.48	32	Grey	QBS	KJ		3	25% biotite, quartz with calcite	E863379	-1	13
YCS15-03	32	33.53	Grey	QBS	KJ	1	1	25% biotite, small quartz grains	E863381	-1	24
YCS15-03	33.53	35.05	Grey	QBS	KJ	1	1	35% biotite	E863382	-1	10
YCS15-03	35.05	36.58	Grey	QBS	KJ	4	4	35% biotite, 35% quartz	E863383	-1	14
YCS15-03	36.58	38.1	Grey	QZT	KJ	2	4	Small grains, chlorite and limonite and iron staining on quartz grains. Almost quartzite.	E863384	-1	10
YCS15-03	38.1	39.62	Grey	QZT	KJ	1	3	40% quartz, 40% biotite	E863385	-1	4
YCS15-03	39.62	41.15	Grey	QZT	KJ	3	4	Almost quartzite, 40% biotite, 40% quartz	E863386	-1	11
YCS15-03	41.15	42.67	Grey	QZT	KJ	1	4	Lots of quartz almost quartzite. 40% biotite, 40% quartz	E863387	-1	7
YCS15-03	42.67	44.2	Grey	QZT	KJ	2	5	90% quartz with biotite (10%)	E863388	-1	6
YCS15-03	44.2	45.72	Grey	QBS	KJ	1	3	40% quartz, 40% biotite	E863389	-1	7
YCS15-03	45.72	47.24	Grey	QBS	KJ	1	3	Pinkification - 40% biotite, 40% quartz. Alteration of the quartz.	E863391	-1	6
YCS15-03	47.24	48.77	Grey	QBS	KJ	2	2	20% quartz - alteration minimal	E863392	-1	6
YCS15-03	48.77	50.29	Grey	QBS	KJ	1	3	50% quartz, grains grains chloritized	E863393	-1	6
YCS15-03	50.29	51.82	Grey	QBS	KJ	1		40% quartz	E863394	-1	7
YCS15-03	51.82	53.34	Grey	QBS	KJ		2	40% biotite, quartz grains and quartz within biotite heavily chloritized. *Olivine*	E863395	-1	11
YCS15-03	53.34	54.86	Grey	QBS	KJ		1	20% quartz, quartz minimally chloritized. Olivine?	E863396	-1	10
YCS15-03	54.86	56.39	Grey	QBS	KJ		4	45% quartz, biotite with chloritized quartz veinlets; possible olivine.	E863397	-1	12
YCS15-03	56.39	57.91	Grey	QBS	KJ		4	Same as above	E863398	-1	8
YCS15-03	57.91	59.44	Grey	QBS	KJ	1	3	Biotite and smokey quartz.	E863399	-1	8
YCS15-03	59.44	60.96	Grey	QBS	KJ	1	1	10% quartz, classic QBS (no alteration)	E863400	-1	11
YCS15-03	60.96	62.48	Grey	QBS	KJ		3	quartz chloritized	E863401	-1	10
YCS15-03	62.48	64.01	Grey	QBS	KJ		3	biotite with smokey and chloritized quartz	E863402	-1	11
YCS15-03	64.01	65.53	Black	QBS	KJ		1	90% biotite, penny shaped, 10% quartz	E863403	-1	12
YCS15-03	65.53	67.06	Black	QBS	KJ		1	90% biotite, flat rounded oval chips, 10% quartz	E863404	-1	11
YCS15-03	67.06	68.58	Grey	QBS	KJ	4	5	70% quartz, many grains orange or tiger striped, quartz oxidized	E863405	-1	8
YCS15-03	68.58	70.1	Grey	QBS	KJ	1	5	80% quartz mostly smokey	E863406	-1	8
YCS15-03	70.1	71.63	Black	QBS	KJ		1	80% biotite, 10-15% quartz	E863407	-1	16
YCS15-03	71.63	73.15	Green	QBS	KJ		4	50% quartz, 50% biotite, smokey and chloritized quartz, half black/half green. Trace pyrite in the quartz.	E863408	-1	16
YCS15-03	73.15	74.68	Grey	QBS	KJ	4	4	Likely quartz vein, limonite on all quartz surfaces, some iron staining.	E863409	-1	8
YCS15-03	74.68	76.2	Black	QBS	KJ	3	3	75% biotite (black), 25% orange and white quartz, mm sized quartz vein, some minor muscovite.	E863410	-1	8
YCS15-03	76.2	77.72	Grey	QBS	KJ	5	5	Small chips, black, grey, white and orange, 70% oxidized quartz, likely numerous cm scale veins	E863411	-1	11
YCS15-03	77.72	79.25	Grey	QZT	KJ	5	5	100% oxidized quartz, 1/2 mm sized grains, 1/2 cm sized grains.	E863412	-1	6
YCS15-03	79.25	80.77	Grey	QZT	KJ	5	5	100% oxidized quartz, same as above.	E863413	-1	9
YCS15-03	80.77	82.3	Grey	QZT	KJ	4	5	100% oxidized quartz. mm + cm sized grains.	E863414	-1	8
YCS15-03	82.3	83.82	Grey	QZT	KJ	4	5	Same as above	E863415	-1	19
YCS15-03	83.82	85.34	Grey	QZT	KJ	3	4	1/2 quartz, 1/2 QBS. QBS - with chloritized quartz (black). Quartzite - same as QBS, 100% grey quartz and oxidized mm-cm scale. Vein likely ended part way through run.	E863416	-1	8
YCS15-03	85.34	86.87	Grey	QZT	KJ	1	5	Some biotite grains (10%), 90% quartz. Pinkification and iron stained pyrite in quartz.	E863417	-1	10
YCS15-03	86.87	88.39	Grey	QZT	KJ	1	5	1/2 QBS, 1/2 quartzite. QBS - minimal chlorite, almost unaltered. Quartzite - same as sample 863417.	E863418	21	8
YCS15-03	88.39	89.92	Grey	QBS	KJ	2	1	30% muscovite, 60% biotite, 10% quartz	E863419	14	17
YCS15-03	89.92	91.44	Grey	QBS	KJ	1	2	20% muscovite, 40% biotite, 20% quartz	E863421	-1	35
YCS15-03	91.44	92.96	Grey	QZT	KJ		5	5% biotite, quartz vein	E863422	14	37
YCS15-03	92.96	94.49	Grey	QZT	KJ		5	100% quartz	E863423	14	27
YCS15-03	94.49	96.01	Grey	QZT	KJ		5	100% quartz	E863424	20	23
YCS15-03	96.01	97.54	Grey	QZT	KJ		5	100% quartz	E863425	-1	16
YCS15-03	97.54	99.06	Grey	QZT	KJ		5	100% quartz	E863426	-1	7
YCS15-03	99.06	100.58	Grey	QZT	KJ		5	100% quartz	E863427	15	16
YCS15-03	100.58	102.11	Grey	QZT	KJ		5	100% quartz	E863428	-1	14

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-03	102.11	103.63	Grey	QZT	KJ		5	5% biotite	E863429	-1	11
YCS15-03	103.63	105.16	Grey	QZT	KJ		5	45% biotite, 55% quartz, muscovite	E863431	-1	15
YCS15-03	105.16	106.68		QZT	KJ		5	10% biotite, 5% muscovite	E863432	-1	4
YCS15-03	106.68	108.2		QZT	KJ		5	5% biotite, 10% muscovite	E863433	-1	13
YCS15-03	108.2	109.73		QZT	KJ		5	5% biotite	E863434	13	15
YCS15-03	109.73	111.25		QBS	KJ		1		E863435	36	122
YCS15-03	111.25	112.78		QBS	KJ		1	5% quartz	E863436	14	39
YCS15-03	112.78	114.3		QBS	KJ		1	10% quartz (small grains)	E863437	33	62
YCS15-03	114.3	115.82		QBS	KJ		1	5% quartz	E863438	16	18
YCS15-03	115.82	117.35		QBS	KJ		1	5-10% quartz	E863439	6	21
YCS15-03	117.35	118.87		QBS	KJ		1	5% quartz	E863440	19	25
YCS15-03	118.87	120.4		QBS	KJ		1	10% quartz	E863441	7	88
YCS15-03	120.4	121.92		QBS	KJ		1	Small clay fracture evident. 5% quartz.	E863442	19	197
YCS15-03	121.92	123.44	Grey	QBS	DP			3% quartzite fragments; dark brownish grey	E863443	13	76
YCS15-03	123.44	124.97		QBS	DP			1% quartzite fragments	E863444	-1	44
YCS15-03	124.97	126.49		QBS	DP			0.5% quartzite fragments	E863445	12	325
YCS15-03	126.49	128.02		QBS	DP			0.5% quartzite fragments	E863446	23	511
YCS15-03	128.02	129.54		QBS	DP				E863447	5	107
YCS15-03	129.54	131.06		QBS	DP			Pyrite as hairline veinlets along fracture surfaces	E863448	27	20
YCS15-03	131.06	132.59		QBS	DP			5% quartzite fragments, one with blue, sooty very fine grained disseminated sulphide pyrite both veinlets and irregular masses.	E863449	11	93
YCS15-03	132.59	134.11	Grey	QBS	DP			35% pale greenish grey, weakly chlorite altered quartzite	E863451	8	18
YCS15-03	134.11	135.64	Grey	QZT	DP			Light greenish grey	E863452	15	11
YCS15-03	135.64	137.16	Grey	QZT	DP			Light greenish grey	E863453	6	44
YCS15-03	137.16	138.68	Grey	QZT	DP			Light greenish grey	E863454	6	17
YCS15-03	138.68	140.21		QZT	DP			3% quartzite chips	E863455	-1	4
YCS15-03	140.21	141.73		QZT	DP			40% quartzite chips	E863456	-1	8
YCS15-03	141.73	143.26		QBS	DP			35% quartzite chips	E863457	-1	8
YCS15-03	143.26	144.78		QBS	DP			3% quartzite chips	E863458	-1	15
YCS15-03	144.78	146.3		QBS	DP			1% quartzite chips	E863459	-1	8
YCS15-03	146.3	147.83		QBS	DP			35% quartzite chips	E863461	-1	7
YCS15-03	147.83	149.35		QBS	DP			2% quartzite chips	E863462	-1	6
YCS15-03	149.35	150.88		QBS	DP			1% quartzite chips	E863463	-1	3
YCS15-03	150.88	152.4		QBS	DP			10% quartzite chips that contain traces of disseminated pyrite	E863464	-1	8
YCS15-03	152.4	153.92		QZT	DP			40% QBS fragment traces very fine disseminated pyrite in quartzite	E863465	-1	8
YCS15-03	153.92	155.45		QZT	DP			40% quartzite	E863466	-1	9
YCS15-03	155.45	156.97		QZT	DP			5% QBS fragments	E863467	7	19
YCS15-03	156.97	158.5		QZT	DP			3% QBS fragments; rare very fine specks pyrite	E863468	30	733
YCS15-03	158.5	160.02		QZT	DP			3% QBS fragments; no chlorite	E863469	24	213
YCS15-03	160.02	161.54		QZT	DP			2% QBS; no chlorite	E863470	-1	133
YCS15-03	161.54	163.07		QBS	DP		0.2	15% quartzite. One speck cpy (?) seen. One quartz veinlet 1.5mm wide	E863471	-1	35
YCS15-03	163.07	164.59		QBS	DP			10% quartzite	E863472	-1	18
YCS15-03	164.59	166.12		QBS	DP				E863473	-1	11
YCS15-03	166.12	167.64	Green	QBS	DP			QBS/CS; green, chlorite rich dust within this run. Chlorite likely part of primary rock, not alteration product.	E863474	-1	7
YCS15-04	0	1.52		CAS					No Sample		
YCS15-04	1.52	3.05	Orange	QBS	KJ	5	2	Overburden, biotite, muscovite, quartz	E863475	-1	71
YCS15-04	3.05	4.57	Orange	QZT	KJ	5	5	limonite quartz with minimal biotite	E863476	-1	49
YCS15-04	4.57	6.1	Orange	QZT	KJ	5	5	Same as above	E863477	-1	9
YCS15-04	6.1	7.62	Orange	QZT	KJ	5	5	Same as above	E863478	8	38
YCS15-04	7.62	9.14	Grey	QZT	KJ	4	5	Same as above	E863479	-1	7
YCS15-04	9.14	10.67	Grey	QZT	KJ	3	5	quartz, 5% biotite	E863481	-1	6
YCS15-04	10.67	12.19	Grey	QZT	KJ	3	5	quartz, 5% biotite	E863482	5	7
YCS15-04	12.19	13.72	Grey	QZT	KJ	3	50	10% biotite	E863483	-1	87
YCS15-04	13.72	15.24	Grey	QZT	KJ	1	3	40% biotite, 30% muscovite	E863484	-1	63
YCS15-04	15.24	16.76	Grey	QZT	KJ	3	4	40% biotite	E863485	-1	21
YCS15-04	16.76	18.29	Orange	QZT	KJ	4	5	Fresh surfaces, green/grey, old surfaces. Orange.	E863486	-1	7
YCS15-04	18.29	19.81	Orange	QZT	KJ	3	5	Fresh surfaces, green/grey, old surfaces. Orange. 30% biotite	E863487	-1	16
YCS15-04	19.81	21.34	Black	QBS	KJ		1	With small quartz grains (unaltered). 10% quartz, 10% muscovite. Quartz in biotite chlorite and sericite altered.	E863488	35	7
YCS15-04	21.34	22.86	Black	QBS	KJ		1	pyrite on surface planes, 20% muscovite, chlorite/sericite on biotite fractures	E863489	72	9

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-04	22.86	24.38	Black	QBS	KJ		1	20% muscovite, chlorite/sericite on biotite fractures	E863491	6	6
YCS15-04	24.38	25.91	Black	QBS	KJ		1	black/purple, chlorite/sericite on biotite fractures, 10% quartz, 20% muscovite	E863492	92	9
YCS15-04	25.91	27.43	Black	QBS	KJ		1	10% muscovite	E863493	8	5
YCS15-04	27.43	28.96	Black	QBS	KJ		1	15% quartz, minimal muscovite	E863494	6	7
YCS15-04	28.96	30.48	Black	QBS	KJ		1	15% quartz, minimal muscovite	E863495	-1	11
YCS15-04	30.48	32	Black	QBS	KJ		1	15% muscovite, 10% quartz.	E863496	-1	8
YCS15-04	32	33.53	Black	QBS	KJ		1	10% muscovite, 15% quartz. Chlorite/sericite salvages.	E863497	-1	6
YCS15-04	33.53	35.05	Black	QBS	KJ		1	20% muscovite, 15% quartz.	E863498	6	4
YCS15-04	35.05	36.58	Black	QBS	KJ		1	20% muscovite, 15% quartz.	E863499	-1	20
YCS15-04	36.58	38.1	Black	QBS	KJ		1	20% muscovite, 15% quartz.	E863500	-1	5
YCS15-04	38.1	39.62	Black	QBS	KJ		1	20% muscovite, 15% quartz.	L399101	-1	4
YCS15-04	39.62	41.15	Black	QBS	KJ		1	20% muscovite, 15% quartz.	L399102	-1	5
YCS15-04	41.15	42.67	Black	QBS	KJ		1	10% muscovite, 20% quartz	L399103	-1	6
YCS15-04	42.67	44.2	Black	QBS	KJ		1	10% muscovite, 20% quartz	L399104	-1	5
YCS15-04	44.2	45.72	Black	QBS	KJ		2	20% muscovite, 20% quartz.	L399105	6	5
YCS15-04	45.72	47.24	Black	QBS	KJ		2	Iron staining in quartz from biotite, 20-25% quartz.	L399106	5	-1
YCS15-04	47.24	48.77	Black	QBS	KJ		2	Iron staining in quartz from biotite, 20-25% quartz.	L399107	-1	6
YCS15-04	48.77	50.29	Black	QBS	KJ		1	15% quartz, 20% muscovite	L399108	8	5
YCS15-04	50.29	51.82	Black	QBS	KJ		1	15% quartz, 20% muscovite	L399109	-1	6
YCS15-04	51.82	53.34	Black	QBS	KJ		1	15% quartz, 15% muscovite	L399110	-1	11
YCS15-04	53.34	54.86	Black	QBS	KJ		1	15% quartz, 15% muscovite	L399111	5	23
YCS15-04	54.86	56.39	Black	QBS	KJ		2	20% quartz, 15% muscovite	L399112	-1	5
YCS15-04	56.39	57.91	Black	QBS	KJ		2	25% quartz, 10% muscovite	L399113	-1	7
YCS15-04	57.91	59.44	Black	QBS	KJ		2	25% quartz, 10% muscovite, little bit of clay in the matrix	L399114	6	9
YCS15-04	59.44	60.96	Black	QBS	KJ		2	20% quartz, 10% muscovite	L399115	7	8
YCS15-04	60.96	62.48	Black	QBS	KJ		2	20% quartz, 10% muscovite	L399116	9	11
YCS15-04	62.48	64.01	Black	QBS	KJ		2	20% quartz, 10% muscovite	L399117	14	8
YCS15-04	64.01	65.53	Black	QBS	KJ		2	20% quartz, 10% muscovite	L399118	6	11
YCS15-04	65.53	67.06	Black	QBS	KJ		3	25% quartz	L399119	7	8
YCS15-04	67.06	68.58	Black	QBS	KJ		2	20% quartz	L399121	-1	6
YCS15-04	68.58	70.1	Black	QBS	KJ		2	20% quartz	L399122	9	9
YCS15-04	70.1	71.63	Black	QBS	KJ		1	15% quartz	L399123	-1	6
YCS15-04	71.63	73.15	Black	QBS	KJ		1	20% quartz	L399124	-1	23
YCS15-04	73.15	74.68	Black	QBS	KJ		2	25% quartz all chloritized	L399125	9	5
YCS15-04	74.68	76.2	Black	QBS	KJ		4	45% chloritized quartz	L399126	-1	12
YCS15-04	76.2	77.72	Grey	QBS	KJ		4	40-40% quartz, mostly chloritized	L399127	7	28
YCS15-04	77.72	79.25	Grey	QBS	KJ		1	10% quartz	L399128	7	6
YCS15-04	79.25	80.77	Grey	QBS	KJ		1	10% quartz	L399129	-1	7
YCS15-04	80.77	82.3	Grey	QBS	KJ		1	15-20% quartz chlorite on fracture planes, some of the quartz is sugary.	L399131	-1	10
YCS15-04	82.3	83.82	Black	QBS	KJ		1	5-10% quartz	L399132	6	7
YCS15-04	83.82	85.34	Black	QBS	KJ		1	5-10% quartz	L399133	-1	10
YCS15-04	85.34	86.87	Black	QBS	KJ		1	5-10% quartz	L399134	-1	6
YCS15-04	86.87	88.39	Black	QBS	KJ		1	5-10% quartz	L399135	51	7
YCS15-04	88.39	89.92	Black	QBS	KJ		1	5-10% quartz	L399136	12	11
YCS15-04	89.92	91.44	Black	QBS	KJ		1	5-10% quartz	L399137	6	8
YCS15-04	91.44	92.96	Black	QBS	KJ		1	5-10% quartz	L399138	5	8
YCS15-04	92.96	94.49	Grey	QZT	KJ		4	chloritized to biotite, sugary texture, 60% quartz. Might be silicified QBS?	L399139	-1	7
YCS15-04	94.49	96.01	Grey	QZT	KJ		4	chloritized to biotite, sugary texture, 60% quartz. Might be silicified QBS?	L399140	5	4
YCS15-04	96.01	97.54	Black	QBS	KJ		1	10% quartz	L399141	14	11
YCS15-04	97.54	99.06	Black	QBS	KJ		2	15-20% quartz	L399142	16	8
YCS15-04	99.06	100.58	Black	QBS	KJ		1	15% quartz, quartz chloritized	L399143	9	7
YCS15-04	100.58	102.11	Black	QBS	KJ		1	5% quartz	L399144	-1	11
YCS15-04	102.11	103.63	Black	QBS	KJ		1	5% quartz	L399145	6	7
YCS15-04	103.63	105.16	Black	QBS	KJ		1	5% quartz	L399146	-1	8
YCS15-04	105.16	106.68	Black	QBS	KJ		1	10% quartz - chloritized	L399147	13	10
YCS15-04	106.68	108.2	Black	QBS	KJ		1	15% quartz, mostly chloritic	L399148	40	10
YCS15-04	108.2	109.73	Black	QBS	KJ		1	15% quartz, some fresh quartz	L399149	-1	7
YCS15-04	109.73	111.25	Black	QBS	KJ		1	15% quartz, mostly chloritized	L399151	6	9
YCS15-04	111.25	112.78	Black	QBS	KJ		1	10% quartz, mostly chloritized	L399152	-1	7

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-04	112.78	114.3	Black	QBS	KJ		1	15% quartz, mostly chloritized	L399153	-1	14
YCS15-04	114.3	115.82	Black	QBS	KJ		3	45% quartz, some fresh, some chloritized	L399154	-1	24
YCS15-04	115.82	117.35	Black	QBS	KJ		1	10% quartz	L399155	-1	15
YCS15-04	117.35	118.87	Black	QBS	KJ		1	10% quartz, 5% muscovite	L399156	-1	13
YCS15-04	118.87	120.4	Black	QBS	KJ		1	10% sugary quartz	L399157	-1	12
YCS15-04	120.4	121.92	Black	QBS	DP			pyrite specks with chlorite altered quartzite fragments that comprise 2% of interval	L399158	-1	5
YCS15-04	121.92	123.44	Black	QBS	DP			As above; greenish black	L399159	-1	6
YCS15-04	123.44	124.97	Grey	QZT	DP			Light greenish grey, weakly chlorite altered pyrite as irregular disseminations	L399161	-1	14
YCS15-04	124.97	126.49	Grey	QZT	DP			as above; chips 35% QBS	L399162	-1	25
YCS15-04	126.49	128.02	Grey	QZT	DP		0	As above; ~30% QBS. Here QBS contains traces disseminated pyrite which was not seen above.	L399163	-1	7
YCS15-04	128.02	129.54	Black	QBS	DP		0	Dark greenish grey-black; 15% quartzite fragments. Pyrite in both QBS and quartzite	L399164	-1	98
YCS15-04	129.54	131.06	Grey	QBS	DP			Dark greenish grey; 2% quartzite; 3% watery grey, unmineralized quartz veinlets	L399165	-1	24
YCS15-04	131.06	132.59	Grey	QBS	DP			Dark greenish grey; 1% quartzite; rare vein quartz chips	L399166	-1	8
YCS15-04	132.59	134.11		QBS	DP			Say 1% quartzite fragments; no sulphides	L399167	-1	7
YCS15-04	134.11	135.64		QBS	DP			As above except here rare speck pyrite	L399168	-1	23
YCS15-04	135.64	137.16		QBS	DP			As above except here rare speck pyrite	L399169	-1	5
YCS15-04	137.16	138.68		QBS	DP			3% quartzite; increasing sericite alteration with depth	L399170	-1	16
YCS15-04	138.68	140.21		QBS	DP				L399171	-1	7
YCS15-04	140.21	141.73	Grey	QBS	DP		0.5	25% light green-grey quartzite. 0.5% water grey, unmineralized quartz veinlets	L399172	-1	5
YCS15-04	141.73	143.26	Green	CS	DP		0.5	Medium greyish green; 0.5% quartz veinlets	L399173	-1	26
YCS15-04	143.26	144.78		QZT	DP		0.5	60% quartzite; 30% QBS; 10% CS	L399174	-1	17
YCS15-04	144.78	146.3		QZT	DP			20% QBS, 5% CS	L399175	-1	4
YCS15-04	146.3	147.83		QZT	DP			20% QBS	L399176	-1	7
YCS15-04	147.83	149.35	Black	QBS	DP			Greenish black; 5% quartz chips	L399177	-1	5
YCS15-04	149.35	150.88	Green	QZT	DP		0.5	Light grey-green; 5% QBS chips	L399178	-1	-1
YCS15-04	150.88	152.4	Black	QBS	DP			Greenish black (CS?)	L399179	-1	-1
YCS15-04	152.4	153.92	Green	CS	DP		0.3	Dark green; 0.3% quartz veinlets	L399181	-1	3
YCS15-04	153.92	155.45	Green	CS	DP		30	As above; 20% QBS, 10% quartzite	L399182	-1	7
YCS15-04	155.45	156.97	Green	CS	DP			Very dark green; 1% quartzite	L399183	-1	-1
YCS15-04	156.97	158.5	Green	CS	DP			As above; 3% quartzite	L399184	-1	-1
YCS15-04	158.5	160.02	Green	CS	DP			Very dark green; pyrite as irregular specks	L399185	-1	-1
YCS15-04	160.02	161.54	Green	CS	DP		0.5	Very dark green; 0.5% quartz veinlets to 1.5mm wide, unmineralized	L399186	-1	4
YCS15-04	161.54	163.07	Green	CS	DP			Very dark green	L399187	-1	3
YCS15-04	163.07	164.59	Green	CS	DP			Very dark green; no sulphides seen	L399188	-1	4
YCS15-04	164.59	166.12	Green	CS	DP			As above	L399189	-1	4
YCS15-04	166.12	167.64	Green	CS	DP	1	2	Very dark green; 2% quartz veinlets with one quartz veinlet fragment oxidized	L399191	-1	-1
YCS15-04	167.64	169.16	Green	CS	DP		1	As above except 1% quartzite veinlets; not oxidized	L399192	-1	3
YCS15-04	169.16	170.69	Green	CS	DP		1	As above	L399193	-1	3
YCS15-04	170.69	172.21	Grey	QBS	DP			dark brownish grey; pyrite as irregular blebs	L399194	-1	3
YCS15-04	172.21	173.74	Grey	QBS	DP			As above except no sulphides seen	L399195	-1	4
YCS15-04	173.74	175.26		QBS	DP			No sulphides seen	L399196	-1	-1
YCS15-04	175.26	176.78		QBS	DP			No sulphides seen	L399197	-1	-1
YCS15-04	176.78	178.31		QBS	DP			No sulphides seen	L399198	-1	3
YCS15-04	178.31	179.83		QBS	DP			One bleb pyrite 1.5mm	L399199	-1	3
YCS15-04	179.83	181.36		QBS	DP			No sulphides seen	L399200	-1	-1
YCS15-04	181.36	182.88		QBS	DP			No sulphides seen	L399201	-1	-1
YCS15-04	182.88	184.4		QBS	DP			No sulphides seen	L399202	-1	3
YCS15-04	184.4	185.93		QBS	DP			No sulphides seen	L399203	-1	3
YCS15-04	185.93	187.45		QBS	DP		1	1% quartz veinlets	L399204	-1	4
YCS15-04	187.45	188.98		QBS	DP		1	1% quartz veinlets	L399205	-1	3
YCS15-04	188.98	190.5		QBS	DP			No quartz veinlets	L399206	-1	3
YCS15-04	190.5	192.02		QBS	DP			No quartz veinlets	L399207	-1	4
YCS15-04	192.02	193.55		QBS	DP		1	1% quartz veinlets	L399208	-1	-1
YCS15-04	193.55	195.07		QBS	DP		2	2% quartz veinlets	L399209	-1	5
YCS15-04	195.07	196.6		QBS	DP		1	1% quartz veinlets	L399210	-1	4
YCS15-04	196.6	198.12		QBS	DP		0.5	0.5% quartz veinlets	L399211	-1	5
YCS15-04	198.12	199.64		QBS	DP		0.5	0.5% quartz veinlets	L399212	-1	5
YCS15-05	0	1.52		CAS				No Sample			
YCS15-05	1.52	3.05	Black	CS	KJ		1	Chlorite rich, clay when wet	L399213	-1	5

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-05	3.05	4.57	Green	CS	KJ		2	Biotite rich	L399214	-1	-1
YCS15-05	4.57	6.1	Green	CS	KJ			Biotite rich	L399215	-1	-1
YCS15-05	6.1	7.62	Green	CS	KJ		3	Biotite rich (40%), 35% quartz, quartz beige/pink in colour	L399216	-1	4
YCS15-05	7.62	9.14	Green	CS	KJ		3	As above	L399217	-1	-1
YCS15-05	9.14	10.67	Green	CS	KJ		3	As above	L399218	-1	5
YCS15-05	10.67	12.19	Black	QBS	KJ		3	Chlorite rich (40%), 30% quartz	L399219	-1	-1
YCS15-05	12.19	13.72	Grey	QBS	KJ		5	50% quartz, pinkified, secondary biotite and chlorite alteration.	L399221	-1	94
YCS15-05	13.72	15.24	Green	CS	KJ		5	45% quartz, 30% biotite	L399222	-1	20
YCS15-05	15.24	16.76	Green	CS	KJ		5	As above	L399223	-1	26
YCS15-05	16.76	18.29	Black	QBS	KJ		1	10% quartz (tipped samples over--oops, samples only half size now).	L399224	-1	14
YCS15-05	18.29	19.81	Black	QBS	KJ		1	5% quartz	L399225	-1	13
YCS15-05	19.81	21.34	Black	QBS	KJ		1	10% quartz, fresh/slightly p ink	L399226	-1	10
YCS15-05	21.34	22.86	Black	QBS	KJ		1	2% quartz	L399227	-1	8
YCS15-05	22.86	24.38	Black	QBS	KJ		1	5% quartz	L399228	-1	7
YCS15-05	24.38	25.91	Black	QBS	KJ		1	5% quartz	L399229	-1	6
YCS15-05	25.91	27.43	Black	QBS	KJ		1	5% quartz	L399231	-1	7
YCS15-05	27.43	28.96	Black	QBS	KJ		1	5% quartz, pyrite	L399232	-1	6
YCS15-05	28.96	30.48	Black	QBS	KJ		1	5% quartz, pyrite, chlorite alteration	L399233	-1	5
YCS15-05	30.48	32	Black	QBS	KJ		1	2% quartz, pyrite, chlorite alteration	L399234	-1	6
YCS15-05	32	33.53	Black	QBS	KJ		1	10% quartz with pinkification	L399235	-1	8
YCS15-05	33.53	35.05	Black	QBS	KJ		1	10-15% quartz, some limonized, some pinkified	L399236	-1	149
YCS15-05	35.05	36.58	Grey	QZT	KJ		4	45% biotite	L399237	5	535
YCS15-05	36.58	38.1	Black	QBS	KJ	1	3	25% quartz, pinkified, biotite/chlorite	L399238	-1	19
YCS15-05	38.1	39.62	Grey	QZT	KJ	1	4	15% biotite	L399239	-1	5
YCS15-05	39.62	41.15	Black	QBS	KJ		1	0% quartz	L399240	-1	7
YCS15-05	41.15	42.67	Green	QZT	KJ		5	40% biotite, pinkification and iron staining on quartz	L399241	52	1260
YCS15-05	42.67	44.2	Black	QBS	KJ		1	10% quartz	L399242	-1	13
YCS15-05	44.2	45.72	Black	QBS	KJ		2	20% quartz	L399243	-1	7
YCS15-05	45.72	47.24	Black	QBS	KJ		3	25% quartz, pinkified	L399244	-1	7
YCS15-05	47.24	48.77	Grey	QZT	KJ		4	40% biotite	L399245	-1	6
YCS15-05	48.77	50.29	Black	QBS	KJ		4	60% biotite, 40% quartz	L399246	-1	5
YCS15-05	50.29	51.82	Green	QBS	KJ		3	35% quartz pinkified	L399247	-1	5
YCS15-05	51.82	53.34	Green	QZT	KJ		5	25% biotite	L399248	-1	8
YCS15-05	53.34	54.86	Grey	QZT	KJ		5	45% biotite	L399249	-1	5
YCS15-05	54.86	56.39	Grey	QZT	KJ		5	30% biotite	L399251	-1	5
YCS15-05	56.39	57.91	Grey	QZT	KJ		5	25% biotite	L399252	-1	5
YCS15-05	57.91	59.44	Grey	QZT	KJ		5	20% biotite	L399253	-1	7
YCS15-05	59.44	60.96	Black	QBS	KJ		4	35% quartz	L399254	-1	7
YCS15-05	60.96	62.48	Grey	QBS	KJ		3	30% quartz	L399255	-1	7
YCS15-05	62.48	64.01	Grey	QBS	KJ		5	50% QBS, 50% quartz, multi coloured. QBS chloritized, QZT pinkified	L399256	-1	6
YCS15-05	64.01	65.53	Grey	QBS	KJ		5	As above	L399257	-1	9
YCS15-05	65.53	67.06	Black	QBS	KJ		1	5% quartz, pinkified	L399258	-1	8
YCS15-05	67.06	68.58	Black	QBS	KJ		1	5% quartz, pinkified, pyrite in biotite	L399259	-1	6
YCS15-05	68.58	70.1	Black	QBS	KJ		1	5% quartz, pinkified, pyrite in biotite	L399261	-1	7
YCS15-05	70.1	71.63	Black	QBS	KJ			no quartz	L399262	-1	8
YCS15-05	71.63	73.15	Black	QBS	KJ		1	5% quartz with chlorite selvages. No alteration seen	L399263	-1	11
YCS15-05	73.15	74.68	Grey	QBS	KJ		2	20% quartz	L399264	-1	8
YCS15-05	74.68	76.2	Grey	QBS	KJ		3	25% quartz (nost sulphide I've seen yet).	L399265	-1	7
YCS15-05	76.2	77.72	Grey	QBS	KJ		3	As above	L399266	-1	5
YCS15-05	77.72	79.25	Grey	QBS	KJ		3	25% quartz, mostly slightly pink pyrite primarily on biotite schist surfaces	L399267	-1	7
YCS15-05	79.25	80.77	Green	QBS	KJ		1	5% quartz, biotite is chloritized	L399268	-1	9
YCS15-05	80.77	82.3	Green	QBS	KJ		1	10% quartz	L399269	-1	6
YCS15-05	82.3	83.82	Black	QBS	KJ		1	5% quartz	L399270	-1	6
YCS15-05	83.82	85.34	Green	QBS	KJ		1	15% quartz pinkified	L399271	-1	8
YCS15-05	85.34	86.87	Black	QBS	KJ		1	10% quartz, pinkified and chloritic	L399272	-1	6
YCS15-05	86.87	88.39	Pink	QZT	KJ		5	45% QBS (unaltered), 55% quartz (pink), possible cu stain	L399273	-1	5
YCS15-05	88.39	89.92	Grey	QZT	KJ		5	45% QBS (unaltered), 55% quartz (pink and grey)	L399274	-1	5
YCS15-05	89.92	91.44	Grey	QBS	KJ	1	5	40% quartz chlorite and oxidized. Neon lime green surfaces.	L399275	-1	7
YCS15-05	91.44	92.96	Green	QBS	KJ	1	3	35% quartz (pink)	L399276	-1	7

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-05	92.96	94.49	Black	QBS	KJ		1	10% quartz (smokey)	L399277	-1	6
YCS15-05	94.49	96.01	Black	QBS	KJ		1	5% quartz (pink)	L399278	-1	5
YCS15-05	96.01	97.54	Black	QBS	KJ		1	5% quartz (white)	L399279	-1	-1
YCS15-05	97.54	99.06	Black	QBS	KJ		1	10% quartz (pink + green)	L399281	7	6
YCS15-05	99.06	100.58	Black	QBS	KJ		1	10% quartz (pink + green)	L399282	5	7
YCS15-05	100.58	102.11	Black	QBS	DP		2	2% quartz as above	L399283	-1	5
YCS15-05	102.11	103.63	Black	CS	DP		0.5	Greenish black; silt-sized fraction in drill return is medium green colour.	L399284	6	4
YCS15-05	103.63	105.16	Black	QBS	DP			(?CS). More like 330-335'. Black.	L399285	6	-1
YCS15-05	105.16	106.68	Black	QBS	DP			As above.	L399286	-1	3
YCS15-05	106.68	108.2	Green	QBS	DP		2	2% pink and green quartz as for 32-325'.	L399287	-1	3
YCS15-05	108.2	109.73	Black	QBS	DP		0.5	Greenish black; 0.5% quartz veinlets, grey and pinkish	L399288	6	6
YCS15-05	109.73	111.25	Black	QBS	DP		1	As above except 1% veinlets; less chlorite altered. Epidote (?) 0.2%	L399289	-1	4
YCS15-05	111.25	112.78	Black	QBS	DP		1	As above	L399291	5	4
YCS15-05	112.78	114.3	Black	QBS	DP			Black; unaltered; no veinlets; no sulphides seen.	L399292	5	-1
YCS15-05	114.3	115.82	Black	QBS	DP		0.2	As above except 0.2% quartz veinlets	L399293	6	5
YCS15-05	115.82	117.35		QBS	DP		0.5	0.5% quartz veinlets with chlorite altered selvages that contain rare speck pyrite	L399294	11	6
YCS15-05	117.35	118.87	Black	QBS	DP			Black, fresh.	L399295	8	3
YCS15-05	118.87	120.4	Black	QBS	DP			Black, fresh.	L399296	6	-1
YCS15-05	120.4	121.92	Black	QBS	DP		0.2	Black, fresh. 0.2% quartz veinlets	L399297	-1	4
YCS15-05	121.92	123.44	Black	QBS	DP		0.4	Black, fresh. 0.4% quartz veinlets	L399298	6	5
YCS15-05	123.44	124.97	Black	QBS	DP		0.5	Black, fresh; 0.5% quartzite (?)	L399299	7	4
YCS15-05	124.97	126.49	Black	QBS	DP		0.5	As above except no pyrite	L399300	7	3
YCS15-05	126.49	128.02	Black	QBS	DP		0.5	As above except no pyrite; 0.2% light green epidote (?) veinlets to 1.2mm	L399301	9	4
YCS15-05	128.02	129.54	Black	QBS	DP			Black; fresh; no sulphides seen	L399302	10	5
YCS15-05	129.54	131.06	Black	QBS	DP			As above	L399303	9	5
YCS15-05	131.06	132.59	Black	QBS	DP			As above	L399304	10	5
YCS15-05	132.59	134.11	Black	QBS	DP			As above	L399305	7	5
YCS15-05	134.11	135.64	Black	QBS	DP			As above, except 0.1% epidote (?) veinlets	L399306	11	4
YCS15-05	135.64	137.16	Black	QBS	DP			As above; local trace hematite	L399307	10	5
YCS15-05	137.16	138.68	Black	QBS	DP			Black, fresh; one chip pink (dusty hematite colour?) quartzite	L399308	8	5
YCS15-05	138.68	140.21	Black	QBS	DP			Black, fresh; say 0.2% light green epidote (?) Chlorite) spots	L399309	11	6
YCS15-05	140.21	141.73	Black	QBS	DP			Black, fresh; trace chlorite and rare blebs.	L399310	9	4
YCS15-05	141.73	143.26	Black	QBS	DP		0.3	Black, fresh; 0.3% quartz veinlets	L399311	10	4
YCS15-05	143.26	144.78	Black	QBS	DP			Black, fresh; no sulphides seen	L399312	8	7
YCS15-05	144.78	146.3	Black	QBS	DP		0.5	Black, fresh; 0.5% quartz veinlets	L399313	-1	8
YCS15-05	146.3	147.83	Black	QBS	DP			Black, fresh	L399314	7	10
YCS15-05	147.83	149.35	Black	QBS	DP		0.5	Black, fresh; 0.3% quartz veinlets	L399315	8	9
YCS15-05	149.35	150.88	Black	QBS	DP			Black, fresh	L399316	9	6
YCS15-05	150.88	152.4	Black	QBS	DP			Black, fresh; 3% quartzite; weakly chloritized	L399317	8	-1
YCS15-05	152.4	153.92		QBS	DP		2	2% quartz veinlets; 2% quartzite	L399318	8	-1
YCS15-05	153.92	155.45		QBS	DP			Rare speck subhedral pyrite; 5% quartzite; more chlorite altered	L399319	10	-1
YCS15-05	155.45	156.97		QBS	DP		10	10% pale orange vein (quartzite?) quartz. No sulphides seen.	L399321	7	3
YCS15-05	156.97	158.5		QZT	DP			5% QBS fragments. No sulphides seen	L399322	7	-1
YCS15-05	158.5	160.02		QZT	DP			1% QBS chips. No sulphides seen	L399323	9	3
YCS15-05	160.02	161.54		QBS	DP			45% quartz, 55% biotite	L399324	8	3
YCS15-05	161.54	163.07		QZT	DP			75% (quartz?); 25% biotite	L399325	9	4
YCS15-05	163.07	164.59		QZT	DP			35% QBS; no sulphides seen	L399326	12	4
YCS15-05	164.59	166.12		QBS	DP				L399327	11	5
YCS15-05	166.12	167.64		QBS	DP				L399328	11	-1
YCS15-05	167.64	169.16		QBS	DP		2	2% quartz veinlets	L399329	9	8
YCS15-05	169.16	170.69	Grey	QBS	DP		0.5	0.5% watery grey quartz veinlets; no sulphides seen	L399331	-1	8
YCS15-05	170.69	172.21		QBS	DP		0.5	0.5% quartz veinlets, one of which contains pyrite grain	L399332	-1	5
YCS15-05	172.21	173.74		QBS	DP		0.5	0.5% quartz veinlets; fresh rock	L399333	7	-1
YCS15-05	173.74	175.26		QBS	DP			1% quartzite grains; no sulphides	L399334	-1	5
YCS15-05	175.26	176.78		QBS	DP				L399335	-1	4
YCS15-05	176.78	178.31	Grey	QBS	DP		0.7	Dark brownish grey; 0.7% quartz veinlets	L399336	-1	3
YCS15-05	178.31	179.83	Grey	QBS	DP		0.4	Dark brownish grey; 0.4% quartz veinlets	L399337	-1	5
YCS15-05	179.83	181.36	Grey	QBS	DP		0.4	Dark brownish grey; 0.4% watery grey quartz veinlets	L399338	-1	5
YCS15-05	181.36	182.88		QBS	DP		0.4	As above	L399339	8	4

HoleID	mFrom	mTo	Colour	Rock Type	LB	Ox	Qtz Vn	Comments	SampleID	Au ppb	As ppm
YCS15-05	182.88	184.4		QBS	DP		0.2	As above except 0.2% quartz veinlets	L399340	-1	-1
YCS15-05	184.4	185.93		QBS	DP		0.2	As above; 0.2% quartz veinlets	L399341	148	-1
YCS15-05	185.93	187.45	Grey	QBS	DP			Dark brownish grey; no sulphides seen. No quartz veinlets seen	L399342	-1	4
YCS15-05	187.45	188.98	Grey	QBS	DP		1	1% pale grey quartz veinlets; rare speck pyrite	L399343	-1	3
YCS15-05	188.98	190.5		QBS	DP		0.3	0.3% quartz veinlets	L399344	-1	5
YCS15-05	190.5	192.02		QBS	DP		0.2	0.2% quartz veinlets	L399345	-1	5

APPENDIX 3
LABORATORY ASSAY CERTIFICATES,
CERTIFICATION AND ANALYTICAL
METHOD SUMMARIES
(See Attached CD)