

STRATEGIC METALS LTD.


FIGURE 4 ARCHER, CATHRO & ASSOCIATES (1981) LIMITED REGIONAL GEOLOGY KING PROPERTY

0 1 2 3 4 5 km

UTM ZONE 8, NAD 83, 115H/09

FILE: ../2015/King/F_4-Regional_Geology.wor

DATE: OCTOBER 2015



- | | |
|---|--|
| <div style="background-color: yellow; border: 1px solid black; padding: 2px; display: inline-block;">Q</div> | QUATERNARY
Q: OVERBURDEN
Unconsolidated glacial, glaciofluvial and glaciolacustrine deposits; fluvial silt, sand, and gravel, and local volcanic ash, in part with cover of soil and organic deposits |
| <div style="background-color: yellow; border: 1px solid black; padding: 2px; display: inline-block;">TQS</div> | TERTIARY AND QUATERNARY
TQS: SELKIRK GROUP
Resistant, brown weathering, columnar jointed, vesicular to massive basalt flows; minor pillow basalts; basaltic tuff and breccia |
| <div style="background-color: orange; border: 1px solid black; padding: 2px; display: inline-block;">uKC</div> | UPPER CRETACEOUS
uKC: CARMACKS GROUP
Augite-olivine basalt and breccia; hornblende-feldspar porphyry, andesitic and dacitic flows; vesicular, augite-phyric andesite and trachyte; minor sandy tuff |
| <div style="background-color: lightgreen; border: 1px solid black; padding: 2px; display: inline-block;">JL</div> | EARLY JURASSIC
JL: LABERGE GROUP
Poorly sorted, medium bedded to massive arkosic sandstone and minor shale with interbeds and thick members of resistant heterolithic pebble and boulder conglomerate; recessive, dark brown weathering, thin bedded silty shale |
| <div style="background-color: orange; border: 1px solid black; padding: 2px; display: inline-block;">EJgA</div> | EARLY JURASSIC
EJgA: AISHIHIK SUITE
Medium to coarse grained, foliated biotite-hornblende granodiorite; foliated hornblende diorite to monzonite with local potassium feldspar megacrysts |
| <div style="background-color: lightgreen; border: 1px solid black; padding: 2px; display: inline-block;">uTrP</div> | UPPER TRIASSIC
uTrP: POVOAS FORMATION
Augite- or feldspar-phyric, locally pillowed andesitic basalt flows, breccia, tuff, sandstone and argillite; local dacitic breccia and tuff with minor limestone, greenstone, chlorite schist, chlorite-augite-feldspar gneiss and amphibolite |