

MESOZOIC

JL2: LABERGE GROUP: Poorly sorted, medium-bedded to massive arkosic sandstone and minor shale with interbeds of heterolithic pebble and boulder conglomerate; recessive, dark brown to greenish, silty shale (Tanglefoot Formation)

uTrAK1: AKSALA: brown shale, black and minor red siltstone, greenish, calcareous greywacke and interbedded bioclastic, argillaceous limestone igneous or limestone-clast pebble and cobble conglomerate; lahar debris flows; rare feldspar-augite porphyry flows (Casca member)

uTrAK2: AKSALA: massive to thick bedded limestone; minor thin-bedded argillaceous to sooty limestone; coarsely crystalline, massive dolostone; minor laminated chert; massive to poorly bedded, limestone conglomerate debris flows and fanglomerate (Hancock member)

uTrP: POVOAS: augite or feldspar phyric, locally pillowed andesitic basalt flows, breccia, tuff, sandstone and argillite local dacitic breccia and tuff with minor limestone greenschist, chlorite schist, chlorite-augite-feldspar gneiss, amphibolite (Povoas Formation)

STRATEGIC METALS LTD.

FIGURE 4

ARCHER, CATHRO & ASSOCIATES (1981) LIMITED

BEDROCK GEOLOGY

POV PROPERTY

0 1,000

meters

UTM ZONE 8, NAD 83, 105E/06

FILE: ...2016/POV | DATE: SEPTEMBER 2016